**Functionalized Bispidines and Ultrasmall Silicon Nanoparticles
for Cancer Imaging**

DISSERTATION

**Zur Erlangung des akademischen Grades
Doctor rerum naturalium
(Dr.rer.nat.)**

**vorgelegt
der Fakultät Mathematik und Naturwissenschaften
der Technischen Universität Dresden**

**von
Master of Science Garima Singh
geboren am 23. Dezember 1988 in Lucknow
Indien
eingereicht am 20.03.2019**

**Die Dissertation wurde in der Zeit von Juli 2015 bis Dezember 2018 am
Helmholtz-Zentrum Dresden – Rossendorf im Institut für Radiopharmazeutische
Krebsforschung angefertigt.**

**Functionalized Bispidines and Ultrasmall Silicon Nanoparticles
for Cancer Imaging**

DISSERTATION

**To get the academic degree
Doctor rerum naturalium
(Dr.rer.nat.)**

**Submitted to
The Faculty of Mathematics and Natural Sciences
Technical University Dresden**

**Submitted by
Master of Science Garima Singh
Born on 23. December 1988 in Lucknow
India**

**Submitted on 20.03.2019
Defended on 17.05.2019**

**The dissertation was performed from July 2015 till December 2018 in the Institute
of Radiopharmaceutical Cancer Research at the Helmholtz-Zentrum – Dresden
Rossendorf.**

Gutachter: Prof. Dr. rer. nat. habil. Jörg Steinbach
(Technical University Dresden and Helmholtz-Zentrum Dresden -
Rossendorf)

Prof. Luisa De Cola
(University of Strasbourg and Karlsruhe Institute of Technology)

Eingereicht am: 20.03.2019

Datum der Verteidigung:

To my family for always believing in me.

Table of Contents

Chapter 1 Introduction.....	1
1.1 General introduction	1
1.2 Aims and objectives of the thesis	6
Chapter 2 Literature review and theoretical part	7
2.1 Molecular imaging of cancer.....	7
2.1.1 Positron Emission Tomography.....	8
2.1.2 Optical imaging	19
2.2 Multimodal molecular imaging	21
2.2.1 Silicon nanoparticles as multimodal imaging agents.....	23
2.2.2 Factors affecting the biodistribution and pharmacokinetics of NPs	27
2.2.3 Tumor targeting strategies for molecular imaging agents	30
2.2.4 Targeted functionalization of NPs and BFCAs for imaging purposes.....	32
Chapter 3 Results and Discussion	35
3.1 Synthesis, characterization and radiolabeling studies of hexadentate bispidine-based bifunctional chelating agents	37
3.1.1 Synthesis of hexadentate bispidine-based ligands	39
3.1.2 Radiochemistry of modified bispidine derivatives	59
3.1.3 Conclusions	61
3.2 Design and synthesis of Bispidine-based TATE and sd antibody bioconjugates 63	
3.2.1 Bispidine-based bioconjugates of somatostatin agonist TATE.....	63
3.2.2 Bispidine-based EGFR targeting sdAb conjugates B5-sdAb	71
3.2.3 Conclusions	80
3.3 Synthesis and characterization of amine-terminated ultrasmall Si NPs	81
3.3.1 Synthesis of amine-terminated Si NPs	81
3.3.2 Characterization of Si NPs	82
3.3.3 Surface characterization of Si NPs.....	84
3.3.4 Conclusions	85

3.4	Synthesis, characterization, and biodistribution studies of Si NP-based mono-labeled PET and optical imaging agents	86
3.4.1	Si NP-NOTA derived PET imaging agent	88
3.4.2	Synthesis and characterization of Si NP-Bispidine (Si NP-B4)	93
3.4.3	Synthesis, characterization, <i>in vitro</i> and <i>in vivo</i> evaluation of Si NP-IR800 as an optical imaging probe.....	95
3.4.5	Conclusions	104
3.5	Development of dual-labeled Si NPs and its <i>in vitro</i> and <i>in vivo</i> evaluation ...	106
3.5.1	Synthesis and characterization of Si NP-IR800-NOTA conjugate.....	107
3.5.2	Photophysical characterization of Si NP-IR800-NOTA and comparison with mono-labeled imaging agents	109
3.5.3	Radiolabeling studies of dual-labeled Si NP-IR800-NOTA	110
3.5.4	Quantitative estimation of the amount of fluorophore and the radiolabel on the Si NP conjugate.....	111
3.5.5	Cell Cytotoxicity Studies of the Si NP-IR800-NOTA	114
3.5.6	<i>In vivo</i> PET analysis and biodistribution studies, urine analysis and <i>ex vivo</i> examination of [⁶⁴ Cu]Cu-Si NP-IR800-NOTA	115
3.5.7	Comparative analysis of [⁶⁸ Ga]Ga-Si NP-IR800-NOTA and [⁶⁴ Cu]Cu-Si NP-IR800-NOTA in A431 tumor bearing mice	121
3.5.8	Conclusions	127
3.6	Investigation of mono-labeled and dual-labeled Si NPs as targeted imaging agents.....	128
3.6.1	Synthesis of Si NP-PEG-5B9 peptide conjugates.....	128
3.6.2	Characterization of the targeted Si NP conjugates	129
3.6.3	Conclusions	134
Chapter 4 Conclusions and Outlook		135
4.1	Conclusions	135
4.2	Outlook	140
Chapter 5 Experimental Section		141
5.1	Chemicals and Materials	141
5.2	Analysis, characterization and purification techniques and methods	143

5.3	Radionuclide production (^{68}Ga and ^{64}Cu)	146
5.4	Syntheses	148
5.4.1	Synthesis of bispidines derivatives (B1-B11), intermediates (B1', B3', B8', B9', B11') and bifunctional linkers (L1, L2)	148
5.4.2	Synthesis of bispidines-based bioconjugates B4-PEG-TATE and B5-sdAb-7C12	160
5.4.3	Synthesis and modification of Si NPs by hydrothermal method	161
5.5	Radiolabeling experiments	166
5.6	<i>In vitro</i> cell experiments	167
5.7	Cell binding studies	169
5.8	Animal experiments	170
Chapter 6 References		172
Chapter 7 Annex		192
7.1	Acronyms	192
7.2	Supplementary Data	195
Chapter 8 Publications, Conferences and Awards		203
Chapter 9 Acknowledgements		206
Chapter 10 Statement of declaration / Erklärungen		208

Table of Figures

Figure 1.1.1: Representation of an idealized multimodal nanoparticle for cancer diagnostic and therapeutic applications.	4
Figure 2.1.1: Electromagnetic spectrum and operating ranges of the different imaging modalities: from radiowaves (MRI) <i>via</i> near-infrared (NIR) to gamma radiation (PET/SPECT).	8
Figure 2.1.2: Illustration of the mechanism and utility of PET imaging.	9
Figure 2.1.3: The crystal structures and coordination sphere geometry of Cu(II)-NOTA, Cu(II)-TACN and Cu(II)-DMPTACN.	10
Figure 2.1.4: Derivatives of NOTA (1,4,7- triazacyclononane-1,4,7-triacetic acid) chelator with amine, isothiocyanate, maleimide and active succinimide ester functionalities.	12
Figure 2.1.5: Structure of the natural product sparteine and its resemblance to Bispidine.	13
Figure 2.1.6: Three possible conformations of bispidines (chair-chair, chair-boat and boat-boat).	14
Figure 2.1.7: Bispidines in their usual numbering system.	14
Figure 2.1.8: Conformations adapted by C2/4 substituted bicyclic bispidines.	15
Figure 2.1.9: Chemical and Cu(II)-complexed X-ray structures of N7 pyridine (1) and N7 pyridazine substituted (2) bispidines.	17
Figure 2.1.10: Radiolabeling and bioconjugation strategies for the preparation of radiolabeled BFCA–biomolecule conjugates along with the choice of radionuclides for diagnostic and therapeutic applications.	19
Figure 2.1.11: <i>In vivo</i> dual-modal, five-color lymphatic drainage imaging with the ability to visualize five distinct lymphatic drainages.	20
Figure 2.1.12: FDA and EMA-approved fluorescent dyes for <i>in vitro</i> and <i>in vivo</i> imaging.	21
Figure 2.1.13: NIRF dye derivatives (IRDye 800CW was used for this research work).	21
Figure 2.2.1: Representation of the development of medical imaging technique towards multimodal imaging.	22
Figure 2.2.2: SPECT/CT and NIR imaging for cancer tumor diagnosis.	23

Figure 2.2.3: Representation of ^{64}Cu -functionalized dextran-coated silicon quantum dots administered intravenously to the <i>in vivo</i> system and a PET image 5 min <i>p.i.</i> showing renal clearance and liver accumulation. Reprinted from [40]	24
Figure 2.2.4: Synthesis strategies of ultrasmall Si NPs.....	26
Figure 2.2.5: Schematic drawing of the structure of nanoparticle-protein complexes in plasma.	28
Figure 2.2.6: Section of kidney cortex: Typical biodistribution behavior and renal clearance of ultrasmall NPs in kidney tissue.....	29
Figure 2.2.7: Difference between the active and passive targeting of tumors.	32
Figure 2.2.8: Smallest known fully functional camelid antibody- fragment (Nanobody) derived from heavy chain only antibodies.....	33
Figure 2.2.9: Structure of EG-TATE.	34
Figure 3.1.1: The most stable double chair (cc) configuration of bispidine with endo-endo pyridyl-substitution at C2/4.	37
Figure 3.1.2: The N3 and N7 substituted pentadentate bispidine isomers showing prominent differences in their stability constants for Cu(II) complexes.....	38
Figure 3.1.3: Coordination environment around the Cu(II) ion in Bispidol.	38
Figure 3.1.4: ESI-MS spectra of B2 showing the $[\text{M}+\text{H}]^+$ ion peak at m/z 596.2.	41
Figure 3.1.5: Comparative ^1H -NMR spectra (aliphatic region) of L1, (B3'+B3) and B3 with labeled characteristic signals.	43
Figure 3.1.6: ^1H -NMR spectrum of B3 (measured in CD_3CN).	44
Figure 3.1.7: Comparative RP-HPLC chromatograms for B1–B5 with their respective retention times (R_t).	45
Figure 3.1.8: Comparative ^{13}C -NMR spectra of ligands B1–B4 with labeled characteristic NMR signals.	46
Figure 3.1.9: Comparative FTIR spectra for B1–B5 with labeled characteristic vibrational frequencies.	47
Figure 3.1.10: ESI-MS spectra of B5 showing the $[\text{M}+\text{H}]^+$ ion peak at m/z 939.1.	50
Figure 3.1.11: ESI-MS spectra of B6 showing the $[\text{M}+\text{H}-17]^+$ and $[\text{M}-17+2\text{H}]^{++}$ ion peaks at m/z 736.2 and 368.7.....	52
Figure 3.1.12: ESI-MS spectrum of B7 showing the $[\text{M}+\text{H}]^+$ ion peak at m/z 725.5.	53
Figure 3.1.13: ^1H -NMR spectrum of L2.....	54
Figure 3.1.14: ESI-MS spectrum of B9 showing the $[\text{M}+\text{H}]^+$ ion peak at m/z 822.6.	55
Figure 3.1.15: ESI-MS spectrum of B10 showing the $[\text{M}+\text{H}]^+$ ion peak at m/z 566.8. ...	57
Figure 3.1.16: ESI-MS spectra of B11 showing the $[\text{M}+\text{H}]^+$ and $[\text{M}+2\text{H}]^{++}$ ion peak at m/z 621.27 and 311.25.....	58

Figure 3.1.17: Comparative Radio-TLC chromatograms of $[^{64}\text{Cu}]\text{Cu-B1} - [^{64}\text{Cu}]\text{Cu-B5}$ (EDTA challenged) (Above); Radio-TLC of $[^{64}\text{Cu}]\text{Cu-EDTA}$ (standard) (Below).....	60
Figure 3.2.1: Representation of NOTA and NODAGA derivatives of Y3-TATE bioconjugates for PET imaging.....	64
Figure 3.2.2: Radio-HPLC chromatogram of $[^{64}\text{Cu}]\text{Cu-B4}$ reaction mixture and $[^{64}\text{Cu}]\text{CuCl}_2$ (inset).	65
Figure 3.2.3: MALDI-TOF spectrum of B4-TATE with calculated and experimental m/z values.....	67
Figure 3.2.4: MALDI-TOF spectra of B4-EG-TATE with $[\text{M}+\text{H}]^+$ at m/z 2038.5 (calculated m/z 2036.7).	70
Figure 3.2.5: Radio-HPLC chromatogram of $[^{64}\text{Cu}]\text{Cu-B4-EG-TATE}$ reaction mixture with $R_t = 14.8$ min.	71
Figure 3.2.6: General representation of Sortase-A mediated conjugation and purification strategies.	73
Figure 3.2.7: SDS-PAGE analysis of Sortase-A mediated azide-functionalization of the single-domain antibody 7C12.	75
Figure 3.2.8: MALDI-TOF mass spectra of (A) sdAb-Leu-Pro-Glu-Thr-Gly-Gly-His6 and (B) purified azide-functionalized sdAb-Leu-Pro-Glu-Thr- N_3 (calculated $m/z \sim 16.6$ kDa)	75
Figure 3.2.9: SDS-PAGE analysis of strain-promoted azide-alkyne cycloaddition between DBCO-containing bispidine and azide-functionalized single-domain antibody 7C12.	76
Figure 3.2.10: MALDI-TOF mass spectra of purified, single-conjugated sdAb functionalized with the radiometal chelator B5 (calculated $m/z \sim 17.6$ kDa).....	77
Figure 3.2.11: Radio-TLC analysis of ^{64}Cu -labeled bispidine-conjugated sdAb $[^{64}\text{Cu}]\text{Cu-sdAb-B5}$	78
Figure 3.2.12: (A) Discontinuous SDS-PAGE and (B) Autoradiography analysis of $[^{64}\text{Cu}]\text{Cu-sdAb-B5}$	78
Figure 3.2.13: Analysis of binding affinity bispidine-functionalized sdAbs to human EGFR-presenting cells.	79
Figure 3.3.1: Synthesis of amine-terminated Si NPs by hydrothermal method bottom-up strategy. Adapted from [259]	82
Figure 3.3.2: TEM and HR-TEM of Si NPs (left); size distribution graph of Si NPs with an average size of 4.3 ± 1.3 nm (right) [262]	82
Figure 3.3.3: Absorption spectrum of water dispersed Si NPs ($\lambda_{\text{abs}} = 353$ nm).....	83

Figure 3.3.4: Emission spectra of water dispersed Si NPs at different excitation wavelengths in nm (exc).....	83
Figure 3.3.5: FTIR spectra of amine-terminated Si NPs with highlighted characteristic bands.	84
Figure 3.4.1: Radiochemical purity of purified ^{64}Cu -labeled Si NP-NOTA conjugate.	89
Figure 3.4.2: Radiometric titration curve for $^{64}\text{Cu}(\text{II})$ and $\text{Cu}(\text{II})$ labeled Si NP-NOTA conjugate. (Experimental value: $\sim 0.09 \mu\text{mol}$ NOTA/mg Si NP-NOTA).....	90
Figure 3.4.3: <i>In vitro</i> cell viability and membrane integration studies of Si NP-NOTA by CyQuant assay in human embryonic kidney (HEK293) cell lines after incubation up to 72 h.....	91
Figure 3.4.4: <i>In vitro</i> toxicity evaluation of Si NP-NOTA conjugate by MTS and LDH assay after 48 hours in HEK293 cells.....	92
Figure 3.4.5: <i>In vivo</i> PET imaging of ^{64}Cu -Si NP-NOTA at different time points.	93
Figure 3.4.6: Radiometric titration curve for $^{64}\text{Cu}(\text{II})$ and $\text{Cu}(\text{II})$ labeled Si NP-B4 conjugate.	95
Figure 3.4.7: Comparative Fluorescence TLC chromatograms to detect the purity of Si NP-IR800.	97
Figure 3.4.8: A comparative absorption (A) and emission spectra (B) of Si NP-IR800 and NHS-IR 800 dye in water.....	98
Figure 3.4. 9: Quantification of IR 800 dye present on Si NP-IR800 by UV-Vis spectroscopy (Experimental value $\sim 0.013 \mu\text{mol}$ IR 800 dye per mg Si NPs).	100
Figure 3.4.10: <i>In vitro</i> cell studies of Si NP-IR800 by CyQuant assay in HEK293 cell lines.	101
Figure 3.4.11: <i>In vitro</i> toxicity evaluation of Si NP-IR800 by MTS and LDH assay after 48 h.....	101
Figure 3.4.12: <i>In vivo</i> optical imaging of Si NP-IR800 at different time points.....	102
Figure 3.4.13: <i>Ex vivo</i> optical imaging of Si NP-IR800 2 h <i>p.i.</i>	103
Figure 3.4.14: TLC analysis of urine of experimental healthy mice 1 h <i>p.i.</i> of Si NP-IR800.	104
Figure 3.4.15: Summary of synthesis, characterization and <i>in vitro</i> and <i>in vivo</i> behavior of mono-labeled Si NP conjugates.	105
Figure 3.5.1: Synthesis of dual-labeled Si NP-based imaging agents with ^{64}Cu binding NOTA for PET and IRDye 800CW for OI.....	106
Figure 3.5.2: Complementary radio-TLC and fluorescence-TLC of Si NP-IR800-NOTA.	109

Figure 3.5.3: Comparative absorption (A) and emission spectra (B) of Si NP-IR800-NOTA, Si NP-IR800 and NHS-IR 800 dye in water.....	110
Figure 3.5.4: pH dependent $^{64}\text{Cu}(\text{II})$ radiolabeling studies of Si NP-IR800-NOTA.....	111
Figure 3.5.5: Radiometric titration curve for $^{64}\text{Cu}(\text{II})$ and $\text{Cu}(\text{II})$ labeled Si NP-IR800-NOTA.....	112
Figure 3.5.6: Quantification of IR 800 dye conjugated to Si NP-IR800-NOTA (Experimental value $\sim 0.013 \mu\text{mol IR800} / \text{mg Si NP-IR800}$).	113
Figure 3.5.7: <i>In vitro</i> cell studies of Si NP-IR800-NOTA in HEK293 cell lines.	114
Figure 3.5.8: Comparative <i>in vitro</i> analysis of Si NP conjugates.....	115
Figure 3.5.9: <i>In vivo</i> PET imaging of $[^{64}\text{Cu}]\text{Cu-Si NP-IR800-NOTA}$ at different time points.	116
Figure 3.5.10: Radio-TLC analysis of the urine sample of $[^{64}\text{Cu}]\text{Cu-Si NP-IR800-NOTA}$ injected mice.	117
Figure 3.5.11: Biodistribution and autoradiography studies of $[^{64}\text{Cu}]\text{Cu-Si NP-IR800-NOTA}$ in NMRI nu/nu mice.	119
Figure 3.5.12: Comparative quantification of renal versus hepatobiliary mode of elimination for $[^{64}\text{Cu}]\text{Cu-Si NP-IR800-NOTA}$ in NMRI nu/nu mice.....	120
Figure 3.5.13: <i>Ex vivo</i> optical imaging of Si NP-IR800-NOTA at different time points <i>p.i.</i>	121
Figure 3.5.14: PET scan of $[^{68}\text{Ga}]\text{Ga-Si NP-IR800-NOTA}$ 30 min <i>p.i.</i>	123
Figure 3.5.15: PET/CT and OI/CT whole body scans of $[^{68}\text{Ga}]\text{Ga-Si NP-IR800-NOTA}$ injected mice.	124
Figure 3.5.16: Whole body PET scan of A431 right hind leg tumor bearing NMRI nu/nu mice 2 h <i>p.i.</i> of $[^{64}\text{Cu}]\text{Cu-Si NP-IR800-NOTA}$	125
Figure 3.5.17: PET and OI scans of $[^{64}\text{Cu}]\text{Cu-Si NP-IR800-NOTA}$ injected A431 tumor bearing mice.....	125
Figure 3.5.18: <i>Ex vivo</i> PET (A) and optical imaging (B) scans of excised organs for $[^{64}\text{Cu}]\text{Cu-Si NP-IR800-NOTA}$ injected A431 tumor xenograft mice 24 h <i>p.i.</i>	126
Figure 5.6.1: Synthesis of Si NP-IR800-PEG-5B9 and Si NP-IR800-NOTA-PEG-5B9.	130
Figure 5.6.2: Absorption (A) and emission spectra (B) of Si NP conjugates and IR 800 dye.	131
Figure 5.6.3: Agarose gel for Si NP-IR800-PEG-5B9 (1a, 1b) and Si NP-IR800-NOTA-PEG-5B9 (2a, 2b). IR short channel (710-730 nm) IR long channel (810-840 nm)	133
Figure 5.6.4: Agarose (left) and SDS PAGE (right) gel for Si NP-IR800-PEG-5B9 and Si NP-IR800-NOTA-PEG-5B9.	133

Figure 7.1: ESI-MS of B1' [M+H] ⁺ <i>m/z</i> 461.3 in MeOH:H ₂ O (1:1) v/v (left); ESI-MS of B1 [M+py+TFA] ⁺ <i>m/z</i> 611.6 in ACN:H ₂ O (1:1) v/v (right)	195
Figure 7.2: ¹ H NMR of B1' in CDCl ₃	195
Figure 7.3: ¹ H NMR of B1 in CDCl ₃	195
Figure 7.4: ¹ H NMR of B2 in CDCl ₃	196
Figure 7.5: ¹³ C NMR of L1 in CDCl ₃	196
Figure 7.6: ESI-MS of B3' [M+H] ⁺ <i>m/z</i> 751.8 in ACN:H ₂ O (1:1) v/v	196
Figure 7.7: ESI-MS of B3 [M+H] ⁺ <i>m/z</i> 651.4 in ACN:H ₂ O (1:1) v/v (left); ESI-MS of B4 [M+H] ⁺ <i>m/z</i> 843.3 in ACN:H ₂ O (1:1) v/v (right)	197
Figure 7.8: ¹ H NMR of B4 in CDCl ₃	197
Figure 7.9: ¹ H NMR of B5 in CDCl ₃	197
Figure 7.10: ¹ H NMR of B6 in CDCl ₃	198
Figure 7.11: ¹³ C NMR of B6 in CDCl ₃	198
Figure 7.12: ¹³ C NMR of L2 in CDCl ₃	198
Figure 7.13: ESI-MS of B8' [M+H] ⁺ <i>m/z</i> 899.1 in ACN:H ₂ O (1:1) v/v (left); ESI-MS of B8 [M+H] ⁺ <i>m/z</i> 799.5 in ACN:H ₂ O (1:1) v/v (right)	199
Figure 7.14: ESI-MS of B9' [M+H] ⁺ <i>m/z</i> 1045.4 in ACN:H ₂ O (1:1) v/v.....	199
Figure 7.15: ¹ H NMR of B9' in CDCl ₃	199
Figure 7.16: ¹ H NMR of B9 in CDCl ₃	200
Figure 7.17: ¹ H NMR of B10 in CDCl ₃	200
Figure 7.18: ¹³ C NMR of B10 in CDCl ₃	200
Figure 7.19: ESI-MS of B11' [M+H] ⁺ <i>m/z</i> 489.1 in ACN:H ₂ O (1:1) v/v.....	201
Figure 7.20: MALDI-TOF spectra of Boc-B4-PEG-TATE with [M+H] ⁺ at <i>m/z</i> 2407.1.	201
Figure 7.21: ¹ H NMR of Si NPs and trisodium citrate in D ₂ O.	201
Figure 7.22: Radiochemical purity of purified ⁶⁴ Cu-labeled Si NP-B4 conjugate.	202
Figure 7.23: TGA analysis of amine-terminated Si NPs.....	202

Chapter 1 Introduction

1.1 General introduction

Cancer is the second leading cause of death globally, and is responsible for an estimated 9.6 million deaths in 2018 [1]. After decades of research in cancer cell biology, many distinct molecular pathways involved have been discovered. This broad disease profile has prompted the tailoring of individualized cancer therapy. The novel personalized therapy has shown immense success in many cancer types for patients with access to those possibilities. According to the World Health Organization (WHO), many of these malignant tumors could be prevented with earlier diagnosis using more precise visualization tools, which could lead to a more effective and personalized therapy. Various modern imaging technologies have been widely used to monitor structural, functional, and molecular changes in cancer tissues and diagnose the tumor at an early stage [2-5].

In recent years, molecular imaging has emerged as an important medicine discipline and has had a great impact on clinics for generating non-invasive high-resolution images of the human body in real time [6-9]. These include optical imaging (bioluminescence or fluorescence), X-ray imaging (X-Ray), computed tomography (CT), magnetic resonance imaging (MRI), positron emission tomography (PET), single-photon emission computed tomography (SPECT), and ultrasound tomography (US) [9-11]. Each imaging modality has its own unique strength and intrinsic limitations, such as spatial/depth resolution and sensitivity, making the achievement of precise and reliable information at the disease site difficult. In order to circumvent the limitations of a single imaging technique and exploit their advantages synergistically, multimodal molecular imaging has been considered. The combined systems like PET-CT, PET-MRI and PET-OI have been developed to gather a more comprehensive information about the disease [12-15].

PET is an established clinical tool for whole-body imaging [13]. This imaging technology utilizes radiopharmaceuticals containing positron-emitting radionuclide. The positron as a source of annihilation results into two γ -rays emitted at an angle of $\sim 180^\circ$. This allows the reconstruction of the images to generate an imaging contrast. The most frequently used radioactive agents include ^{18}F , ^{68}Ga and ^{11}C but for scientific purposes also other positron emitting radionuclides such as ^{13}N , ^{15}O , ^{44}Sc , ^{62}Cu , ^{64}Cu , ^{72}As , ^{74}As , ^{76}Br , ^{82}Rb , ^{86}Y , ^{89}Zr , and ^{124}I are utilized [16].

Additionally to the broad range of radiotracers, nanoparticles (NPs) have emerged as a new approach for PET and other imaging techniques due to their tunable biodistribution and target site accumulation of imaging contrast agents [17]. Ultrasmall nanoparticles (USNPs), with sizes in the 1–3 nm range, exhibit unique properties distinct from those of free molecules and larger sized nanoparticles. These properties are greatly sensitive to both the composition and size of the particles. When applied in medical science, they give rise to the term of “nanomedicine” which has found more applications in preclinical, translational, and clinical research.

The research in the area of nanomedicine spans multidirectional applications varying from diagnosis, drug delivery, gene delivery, targeted therapy, biomarker mapping, to the area of our interest: molecular imaging [7, 18]. Many of these developments are starting to be translated into viable clinical products [19, 20]. The distinct intrinsic properties and facile tailorability of the NPs makes them potential targeted theranostics agents [21].

Some cancer nanomedicines have already been approved by either the Food and Drug Administration (FDA) or European Medicines Agency (EMA) and are used in the clinic to diagnose and cure the disease for over 20 years, while several systems are currently under clinical trials [19, 20, 22]. Doxil, a polyethylene glycol (PEG) functionalized liposomal doxorubicin, was the first approved (FDA 1995) cancer nanomedicine, followed by Abraxane, an albumin-bound paclitaxel NP. More recently approved products such as Marqibo, MEPACT, Feraheme, AuroLase and Cornell Dots (C-Dots) exhibit the strong presence of inorganic NPs in clinical trials [20, 23]. Inorganic NPs have been studied extensively for past decades and have paved a way in this direction [24]. Their unique material- and size-dependent physicochemical properties, the ability to modulate pharmacokinetics through surface chemistry, application of targeting mechanisms to advance the biodistribution and enhancement of contrast signals by incorporating multiple imaging moieties in a single scaffold make them attractive alternatives for imaging and ablation of malignant tissue [25-37].

Cornell Dots (C-Dots), for example, are derivatized silica NPs that are used for enhancing standard PET-optical dual-modality imaging is the nanoconstruct of our interest [23, 38]. The NP particle size of ~7 nm facilitates renal excretion, a near-infrared (NIR) cyanine 5 fluorophore to optimize fluorescence imaging, a PEG coating to avoid opsonization and subsequent internalization by macrophages in the liver. A ^{124}I -radiolabeled cRGDY peptide which functions as both, a radiolabeled molecule that facilitates PET imaging and as a targeting peptide to improve targeting and accumulation

in $\alpha_v\beta_3$ integrin expressing tumors. The first-in-man clinical trial of C-Dots for imaging of cancer in patients with metastatic melanoma were shown to be: (i) well tolerated with no reported toxic or adverse events, (ii) stable *in vivo*, (iii) provide consistent pharmacokinetic patterns, and (iv) capable of imaging lesions, including metastatic lesions, in distinct areas of the body. Effective doses of C-Dots for the clinical trials were comparable to the commonly used diagnostic radiotracers.

The ultimate biocompatibility and extraordinary pharmacokinetics revealed from these studies led to the exploitation of silicon nanoparticles (Si NPs) in the field of bioimaging and biosensing [39-41]. Si NPs featuring benign biocompatibility and unique optical properties (i.e., strong fluorescence coupled with ultrahigh photostability) are emerging as novel high-quality fluorescent nanoprobes for biological imaging analysis, particularly for tracking dynamic biological procedures in long-term and real-time manners [41-43].

The hydrothermal synthesis of the Si NPs has been optimized by several research groups to achieve large-scale synthesis of ultrasmall, hydrophilic, surface functionalized Si NPs [44]. The large surface area of the synthesized Si NPs also provides an extensive platform for combining multiple imaging modalities. These Si NPs are exploited as a core for attachment of labeling and targeting moieties to develop a multi-labeled targeted contrast agent [40, 45-56]. The high surface area and surface-functionalization of the Si NPs can accommodate a variety of substituents including drugs for therapeutic purposes, biological vectors for targeted theranostics, radionuclides for PET/SPECT imaging, fluorophores for optical imaging (OI), and linkers and spacers for antifouling properties (**Figure 1.1**) [26, 38, 41, 43, 57].

The focus of this graduation project is the development of customizable Si NPs based mono-labeled PET or OI and dual-labeled PET-OI imaging probes and examination of their *in vitro* and *in vivo* behavior.

For OI, NIR fluorophores like cyanines have been employed extensively due to their high-resolution visualization in tissue. Several studies show their optical characteristics in cells studies, biodistribution experiments and imaging applications like the tumor-targeted detection and image-guided surgery [58-61]. As a part of this thesis, we have tested the sulfo-cyanine 5 (sulfo-Cy5) and the IRDye 800 CW-Si NP conjugates for *in vitro* and *in vivo* studies.

Figure 1.1.1: Representation of an idealized multimodal nanoparticle for cancer diagnostic and therapeutic applications.

A multifunctional nanoparticle (size < 10 nm) coupled to modular ligand systems for PET and OI and targeted vectors *via* linkers for cancer diagnostics. Therapeutic agents can also be conjugated for theranostics applications. Adapted from [17]

For PET and SPECT imaging techniques, specific radionuclides are reported for diagnostic and imaging purposes with favorable half-lives and decay properties. Copper-64 (^{64}Cu ; $t_{1/2} = 12.8$ h; β^+ , 19%; β^- , 40%; and electron capture 40%) is a distinct radionuclide that is exploited for its diagnostic potential [62, 63]. It is a cyclotron-produced radionuclide, produced by $^{64}\text{Ni}(p,n)^{64}\text{Cu}$ reaction and separated from the Ni-target material with a high molar activity [64]. Generator-based Gallium-68 ($t_{1/2} = 68$ minutes, $\beta^+ = 89\%$, E_{max} , $\beta^+ = 1.92$ MeV) on the other hand, is also a common radionuclide used for assessment of shorter biochemical processes due to its short half-life [16, 65]. The aforementioned properties permit binding of the radiometal to a targeting moiety, tissue accumulation, and clinical administration *in vivo* in a manageable time frame.

The radionuclide is complexed by a specific bifunctional chelating agent (BFCA), which can be coupled to the NPs for PET imaging of the biological system. The required properties of an ideal BFCA are its ability to rapidly form a thermodynamically stable and kinetically inert complex with the radionuclide at ambient temperature under physiological conditions [66-70]. At the same time, it should allow easy linking with vector molecules. A series of BFCAs compatible to the ^{64}Cu radionuclide are available

portraying excellent chelation properties [71]. Multidentate azamacrocyclic chelators such as 1,4,7,10-tetraazacyclododecane-1,4,7,10-tetraacetic acid (DOTA), 1,4,7-triazacyclononane (TACN), and 1,4,7-triazacyclononane-1,4,7-triacetic acid (NOTA) are well-known promising chelators for PET applications [15, 72, 73].

Hexadentate 3,7-diazabicyclo[3.3.1]nonane derivatives, commonly known as bispidines, have been chosen as the main chelator system for this project [74, 75]. The ability to form different functional derivatives and kinetically inert Copper(II) complexes, bispidines behave as an optimal PET imaging agent; therefore, they were studied as potential BFCAs for targeted molecular imaging [76-78]. A series of bispidines were synthesized with varying functional moieties compatible to sensitive fluorophores and appropriate biomolecules to prepare targeted multimodal imaging modalities. Therefore, an isothiocyanate-terminated bispidine derivative was coupled to amine-terminated TATE (Tyr3-Octreotate) peptide specific to somatostatin receptor (SSTR-2) and a dibenzocyclooctyne (DBCO)-terminated bispidine to the tailor-made single-domain antibody (sdAb) to target the epidermal growth factor receptor (EGFR). These receptors are overexpressed in specific cancer malignancies and can be detected by the targeted PET imaging agents. The bispidine derivatives were examined for their effectiveness as potential targeting agents for cancer diagnosis.

1,4,7-triazacyclononane-triacetic acid (NOTA) is another suitable chelator for both ^{64}Cu and ^{68}Ga radiometals that has been used for the development of Si NP based unimodal and bimodal imaging agents [72, 79]. The biocompatibility, fast and efficient radiochelation properties and commercial availability of this chelator has led to its exploitation for PET-based analysis.

1.2 Aims and objectives of the thesis

The aim of this dissertation is the optimization and development of PET and OI agents for cancer diagnosis with high efficiency and non-invasive procedures. The major challenge is to develop a system highly biocompatible and with low side effects to translate into the clinics. Si NPs are suitable candidates due to their extraordinary intrinsic and surface properties. The ultimate goal is the development of ultrasmall multifunctionalizable Si NPs that can be modified by appropriate labels for uni and bimodal imaging. These particles are to be modified in such a way that additional functionalities such as molecules for targeting can be easily added.

The first aim focuses to determine the *in vitro* and *in vivo* behavior of the functionalized Si NPs and render them suitable for diagnostics. Bispidines were modified as a primary PET ligand and modified with diverse functionalities to permit an easy linkage to the NPs and biomolecules. For OI, commercially available fluorophore, IR 800 dye was employed. In parallel, tailor-made bifunctional bispidines were developed as targeted PET imaging agents. TATE peptide (SSTR-2) and sdAb (EGFR) were employed as biological vectors to achieve cancer targeting.

The second aim was the functionalization of Si NP core by PET/OI ligands and their in-depth characterization and evaluation of surface properties. The ultrasmall size and tunable surface properties of the Si NPs (< 5 nm) make them renally clearable, which is an advantage over large sized particles that tend to accumulate in non-targeted tissue of the *in vivo* system. Furthermore, Si NPs are biodegradable in nature, causing no harm over long-term cancer treatments. Several surface coating strategies have already been optimized to be upscaled by efficient and cost-effective techniques in shorter time durations (0.5 – 3 hours).

The third aim was to study the *in vitro* and *in vivo* behavior for the different labeled Si NPs. This includes the cytotoxicity evaluation in cells and absorption, distribution, metabolism and excretion (ADME) studies in small animals.

Altogether, this thesis aims to compile the *in vitro* and *in vivo* information gained by the synthesized mono- and dual-labeled Si NP-based (PET/OI) imaging agents and the novel bispidine-based PET imaging agents, and thus, mark its contribution towards cancer diagnostics.

Chapter 2 Literature review and theoretical part¹

2.1 Molecular imaging of cancer

Molecular imaging is the “visualization, characterization, and quantification of biological processes at the molecular and cellular levels in humans and other organisms”, as defined by the Society of Nuclear Medicine [80].

Currently, there are six major imaging modalities available to assist clinicians in diagnosing and curing human cancer: X-ray (plain film and computed tomography (CT), ultrasound (US), magnetic resonance imaging (MRI), single-photon emission computed tomography (SPECT), positron emission tomography (PET), and optical imaging (OI). These modalities cover the electromagnetic spectrum ranging from radio waves (MRI) via near-infrared (NIR) down to gamma radiation (PET/SPECT) (**Figure 2.1.1**) [4, 6, 28, 81-85]. These non-invasive techniques illustrate, characterize and measure physiological and pathophysiological processes on a molecular level in cells, tissues and living systems. Of these, only CT, MRI, SPECT, and PET are capable of three-dimensional (3D) imaging. Each technique has its own unique strength and intrinsic limitations, such as spatial depth, resolution and sensitivity, making the achievement of precise and reliable information at the disease site difficult. In order to compensate the weak aspects, multimodal molecular imaging has been considered in recent years. This technique involves the combined usage of the modalities that synergistically exploits the advantages of each individual technique combating its drawbacks.

The central theme of this thesis is to develop ultras-small (< 5 nm) probes capable of imaging cancer by a combination of PET and OI. This thesis will also discuss the research efforts made towards evaluating the suitability of developed probes for application in cancer diagnosis [86, 87].

¹ Few sections of this chapter are an excerpt of the published reviews:

Singh, G.; Gott, M.D.; Pietzsch, H.-J.; Stephan, H. **Nuclear and optical dual-labeled imaging agents**
DOI: 10.1055/s-0037-1616472.

Joshi, T.; Kubeil, M.; Nsubuga, A.; **Singh, G.**; Gasser, G., Stephan; H. **Harnessing the Coordination Chemistry of 1,4,7-Triazacyclononane for Biomimicry and Radiopharmaceutical Applications**
DOI : 10.1002/cplu.201800103.

Figure 2.1.1: Electromagnetic spectrum and operating ranges of the different imaging modalities: from radiowaves (MRI) via near-infrared (NIR) to gamma radiation (PET/SPECT).

2.1.1 Positron Emission Tomography

PET is a valuable functional imaging technique mainly used to detect the dynamic changes in the metabolic function of substances in the human body. PET imaging has an advantage of excellent tissue penetration, high detection efficiency, and superb quantitative efficiency. It is used to provide real-time, non-invasive, and quantitative monitoring of the biodistribution, pharmacokinetics and tumor-targeting efficacy in tumor diagnosis. The shortcomings of PET are low spatial resolution, limited acquisition time owing to radionuclides with short half-lives, and radiation exposure [88, 89].

PET is based on the principle of detecting of gamma (γ) rays from the body (**Figure 2.1.2**). A radiolabeled biomolecule is injected into the body, where the radionuclide emits a positron (β^+) and undergoes annihilation process interacting with an electron (e^-). The annihilation leads to a pair of 511 keV γ -rays moving in approximate opposite directions. These γ -rays may be detected by a set of scintillation crystals that converts it into photons. These photons are then detected by photomultiplier or silicon avalanche photodiodes. The detection of coincident pairs of these photons is finally recorded and transformed to a 3D PET image by a reconstruction software.

PET requires the use of a cyclotron to generate short-lived radionuclides such as ^{18}F ($t_{1/2} = 109.8$ min), ^{64}Cu ($t_{1/2} = 12.7$ h), and ^{76}Br ($t_{1/2} = 16.2$ h), or ultra short-lived radionuclides such as ^{11}C ($t_{1/2} = 20.3$ min), ^{13}N ($t_{1/2} = 10$ min), and ^{15}O ($t_{1/2} = 2.04$ min) depending upon the application and requirements. Of these, ^{64}Cu has garnered specific interest due to its favorable decay characteristics. The production method and decay characteristics of radionuclides ^{64}Cu , ^{67}Cu and ^{68}Ga are shown in **Table 2.1.1**.

Figure 2.1.2: Illustration of the mechanism and utility of PET imaging.

Table 2.1.1: Production and properties of PET radionuclides (EC = electron capture) [72, 90, 91]

Radionuclide	$t_{1/2}$	Decay mode			γ MeV (%)	Production
		β^- MeV (%)	β^+ MeV (%)	EC (%)		
^{64}Cu	12.7 h	0.573 (38.4)	0.655 (17.8)	43.8 (43.8)	0.511 (35.6) 1.35 (0.6)	Cyclotron, $^{64}\text{Ni}(p,n)^{64}\text{Cu}$
^{68}Ga	68 min	-	1.9 (87.7%)	11 (12.3)		$^{68}\text{Ge}/^{68}\text{Ga}$ $^{69}\text{Ga}(p,2n)^{68}\text{Ge}$

2.1.1.1 Ligands for radiometals for PET imaging and therapy

Metal-based radiopharmaceutical applications require chemically robust and radiolytically stable ligands, which can form radiometal complexes of high thermodynamic stability and kinetic inertness. A range of well-established ligands that follow the prerequisite of forming inert radionuclide complexes under physiological conditions, even at their nanomolar concentrations, have been reported [92-95].

Examples of commercially available radiometal chelators that have been derived from two well-known azamacrocycles cyclam (1,4,8,11-tetraazacyclotetradecane) and TACN (1,4,7-triazacyclononane), include TETA and NOTA.

The stability constants and coordination geometries of the Cu(II) complexes formed by these, and other widely used macrocyclic chelators, are reported in **Table 2.1.2**. The

crystal structures of the two frequently employed chelators Cu(II)-DMPTACN and Cu(II)-NOTA are shown in **Figure 2.1.3**. These ligands specifically are known to form mononuclear complexes with 1:1 metal-ligand stoichiometry.

Rapid radiolabeling at room temperature becomes essential when working with chelators conjugated to antibodies, which are usually temperature sensitive, or when using radionuclides with short half-life such as ^{68}Ga . Highly preorganized scaffolds like substituted cross-bridged macrocycles and their derivatives and 3,7-diazabicyclo[3.3.1]nonane (bispidines) which show optimal chelation properties are excellent examples of such biologically suited chelators. The facile synthesis of these chelators makes them one of the most exploited ligand systems for biomedical applications.

Figure 2.1.3: The crystal structures and coordination sphere geometry of Cu(II)-NOTA, Cu(II)-TACN and Cu(II)-DMPTACN.

Table 2.1.2: Macrocyclic ligands with their structure, thermodynamic stability constants ($\log K_{\text{Cu(II)-L}}$) and coordination geometry (Copper(II)-Ligand stoichiometry 1:1).

Ligand	Structure	$\log K_{\text{Cu(II)-L}}$ (Coordination geometry)	Ref
NOTA (1,4,7-triazacyclononane-1,4,7-triacetic acid)		21.6 (Distorted trigonal prism)	[96]
DMPTACN bis(2-pyridylmethyl)-1,4,7-triazacyclononane		> 25 (Distorted square pyramidal)	[97]
NODAGA 1,4,7-triazacyclononane, 1-glutaric acid-4,7-acetic acid		20 (Distorted trigonal prism)	[72]
DOTA (1,4,7,10-tetraazacyclododecane-1,4,7,10-tetraacetic acid)		22.7 (Distorted octahedron)	[98]
Cyclam (1,4,8,11-tetraazacyclotetradecane)		27.2 (Square planar)	[99]
TETA (1,4,8,11-tetraazacyclotetradecane-1,4,8,11-tetraacetic acid)		21.9 (Distorted octahedron)	[100]

As a part of this thesis, we will discuss the radiolabeling and biodistribution studies of bispidine and NOTA based ^{64}Cu -labeled PET and PET/OI dual-labeled imaging agents.

2.1.1.1.1 NOTA as a radioligand for ^{64}Cu and ^{68}Ga

Owing to the favorable binding affinity of NOTA for ^{64}Cu and ^{68}Ga radionuclides, *in vivo* studies have mostly been performed with $^{64}\text{Cu(II)}$ complexes of NOTA-attached biomolecules and, derivatives thereof [97]. For example, ^{64}Cu -NOTA was coupled to a bombesin (BBN) tetradecapeptide derivative *via* a linker to target the gastrin-releasing peptide receptor (GRPR). Similarly, ^{64}Cu -NOTA-ML (multi-leucine peptide) was developed for PACE4-targeted prostate cancer detection, ^{64}Cu -NOTA-KCCYSL (bacteriophage selected peptide) for recognizing epidermal growth factor receptor

(EGFR)-rich breast carcinoma, ^{64}Cu -NOTA-a-MSH (alpha-melanocyte stimulating hormone) for melanoma imaging and ^{64}Cu -NOTA-HsTX1 for targeting potassium channels for autoimmune diseases. Moreover, with the aim of dual targeting, ^{64}Cu -NOTA derivatives coupled to peptides that target prostate-specific membrane antigen (PSMA)-rich and GRPR-rich tumors have also been developed. NOTA has also been used for ^{64}Cu radiolabeling of A33 antibodies to evaluate a pre-targeting approach. Derivatives of NOTA with different functionalities like amines, isothiocyanates, maleimides and active NHS ester have been reported and utilized in various coupling reactions (**Figure 2.1.4**).

Figure 2.1.4: Derivatives of NOTA (1,4,7- triazacyclononane-1,4,7-triacetic acid) chelator with amine, isothiocyanate, maleimide and active succinimide ester functionalities.

2.1.1.1.2 Bispidines as ^{64}Cu based PET imaging agents

Bispidine is a bicyclic compound formed by two piperidine molecules over three carbon atoms. The highly preorganized and exceedingly rigid bidentate structure of bispidines makes them exceptional candidates to study transition metal coordination chemistry [101]. Because it resembles naturally occurring alkaloid sparteine (**Figure 2.1.5**), it is an accessible scaffold for interacting with ion channels, G-protein coupled enzymes and receptors, and the development of new radiotracers [74, 75, 102-106].

Figure 2.1.5: Structure of the natural product sparteine and its resemblance to Bispidine.

The detailed literature available on the configuration, confirmation and the radio-complexation of bispidines opens up new ventures to explore its possibilities in the field of radiopharmaceuticals for the development of potential PET imaging agents [21, 77, 103, 107-109]. A summary of established hexadentate and octadentate ligands along with the substituents is reported in **Table 2.1.3**. The tetra, penta and hexadentate ligands have been optimized for ^{64}Cu mononuclear radiometal complexes for PET imaging to study their *in vivo* targeting capability and stability, whereas the octadentate complexes which do not optimally complex ^{64}Cu but form heavy metal complexes (^{177}Lu , ^{111}In and ^{225}Ac) are currently under investigation [76, 110-113].

Table 2.1.3: Hexadentate (1–8) and octadentate (9) bispidine ligands (L) with their substituents [102]

L	R ₁ (C2/4)	R ₂ (N3)	R ₃ (N7)	R ₄ (C1/5)	X (C9)	Y (C9)
1						
2						
3						
4						
5						
6

Stereochemistry of bispidine

Bispidines are usually synthesized by a multi-component double Mannich condensation reaction. An elaborate synthesis of the bispidine core and its derivatives is reported in **Chapter 3.1**. Bispidines usually exist in three conformations, chair-chair (cc), chair-boat (cb), and boat-boat (bb) (**Figure 2.1.6**). The studies suggest that the chair-chair (cc) conformer is the most stable one; however, the introduction of substituents can cause conformational changes.

Figure 2.1.6: Three possible conformations of bispidines (chair-chair, chair-boat and boat-boat).

The diazabicyclononanes derivatives are schematically represented in their usual numbering, as shown in **Figure 2.1.7**. The modified skeleton of the bispidines, in general, includes a ketone (or a hydroxyl group) at the C9 carbon atom with two electron withdrawing ester groups flanking on the C1/5 positions (R_4). The C2/4 positions (R_1) usually incorporate electron donating aromatic groups, such as the pyridyl moieties and show strong metal ligation tendency.

Figure 2.1.7: Bispidines in their usual numbering system.

Upon modification, the bispidine skeleton can adopt one of the following three conformations: chair-chair (cc), chair-boat (cb), or boat-chair (bc). The boat-boat (bb) conformation has been found to be energetically unfavorable. Moreover, the substituents at positions C2 and C4 can independently exist in either axial or equatorial position, as endo-endo (equatorial-equatorial), endo-exo (equatorial-axial) and exo-exo (axial-axial) (**Figure 2.1.8**). This leads to three possible diastereoisomers per conformation. However, the configuration required for highly stable metal complexation is the endo-endo configuration (cis-isomer) because it constitutes the thermodynamically favored isomer.

Figure 2.1.8: Conformations adapted by C2/4 substituted bicyclic bispidines.

(The wiggly bonds represent the possible equatorial/axial orientation of the substituents).

Cu(II)-Bispidine complexes

2,4-pyridyl-substituted bispidine ligands form Cu(II) complexes with high kinetic inertness and thermodynamic stability, and show fast kinetics as well as particularly good selectivity for Cu(II) complexation. A comparison of stability constants for tetra, penta and hexadentate bispidines is summarized in **Table 2.1.4** for different metal ions. Molecular modeling calculations of pyridine-substituted hexadentate bispidines confirms Cu(II) metal ions to be the most suitable ones with ideal size and shape for complexation, in contrast to Ni(II) and Zn(II). Jahn-Teller distortions are frequently observed for Cu(II) d^9 electronic configuration, which leads to an axial elongation or tetragonal compression of the complexes forming distorted geometrical structures [102].

Table 2.1.4: Stability constants of 1:1 Ni(II), Cu(II) and Zn(II) complexes of tetra, penta and hexadentate bispidines.[76, 102]

Structure
Denticity	Tetradentate	Pentadentate	Pentadentate	Hexadentate
Stability constants (log K _{Metal-Ligand})				
Ni(II)	n.d.	6.10	9.54	5.02
Cu(II)	16.56	18.31	15.66	16.28
Zn(II)	11.37	8.28	13.57	9.18

Another interesting study of the tetrapyridyl-substituted hexadentate bispidines-Cu(II) complex reported by Comba *et al.* confirms a repulsion between the α -hydrogen atoms of the two in-plane pyridine donors in a cis configuration (**Figure 2.1.9**) [114]. This interaction generates significant strain in the complex and therefore leads to a loss in complex stability. Substituting one pyridine moiety in the system with a pyridazine relieves this stress but the lower basicity of pyridazine ($pK_a = 2.33$ vs. 5.23 for pyridine) counteracts the steric effect. Structural studies in the solid state and in solution confirmed the relief of steric distortion, but potentiometric titration as well as competition experiments and the redox potentials revealed similar stabilities for both the Cu(II) complexes. The pyridine substituents at the N3/5 and C2/4 positions of bispidines provide much-needed sites to form highly stable metal-ion coordination bonds. Therefore, an additional functional group is needed on this system if further coupling to biological vectors is desired for any potential biomedical applications.

Figure 2.1.9: Chemical and Cu(II)-complexed X-ray structures of N7 pyridine (1) and N7 pyridazine substituted (2) bispidines.

[Cu(II)-1]²⁺ (left) and [Cu(II)-2]²⁺ (right) with α -hydrogen atoms represented as red balls. (ester moieties and the C9 keto group are omitted for clarity) Reprinted and modified from [114]

Ideally, this functionality should be remote to the donor groups involved in complexation in order to avoid any kind of interference in chelation. The bispidines skeleton presents three potential sites for (two esters C1/5 and a ketone C9) that can be directly exploited for incorporating a bioconjugatable functionality [78].

Biodistribution studies of ⁶⁴Cu(II)-Bispidine bioconjugates

A reported *in vivo* biodistribution study of ⁶⁴Cu(II) labeled bispidine-bombesin derivative conjugate shows very fast clearance of the imaging agent through kidneys and urine, with very little activity remaining after 1 hour and more in organs other than the kidney [76]. The fast-renal elimination and target tissue accumulation of the biomarker is expected due to the considerable hydrophilicity and the *in vivo* stability of the bispidine complex.

2.1.1.2 Bifunctional chelating agents (BFCAs)

BFCAs are organic chelators with terminal reactive groups like amines, thiols and carboxylic acids which have been used for indirect radiolabeling of biomolecules, rendering them suitable for imaging and therapy. It serves the dual purpose of binding to

the radionuclide efficiently and forming a highly stable complex, as well as of providing a functionality and/or site for attaching targeting vectors, fluorophores (for dual imaging: combined PET (or SPECT) and optical imaging) and NPs as a core to provide a platform for multifunctionality [63, 66-71, 73, 115-117]. Common bioconjugation strategies utilize functional groups such as carboxylic acids or activated esters (e.g. N-hydroxysuccinimide (NHS)-ester, tetrafluorophenyl (TFP)-ester for coupling amino groups to generate amides and to isothiocyanates to form thiourea bonds. The thiol functional groups couple to maleimide moieties to form succinimide acid amides.

Copper-free click chemistry is gaining popularity in bioconjugate chemistry with reactions like strain-promoted azide–alkyne cycloadditions (SPAAC) [118] (e.g. dibenzocyclooctyne/azide reaction) and Diels–Alder click reactions (e.g. transcyclooctene/1,2,4,5-tetrazine) (**Figure 2.1.10**). The azide–alkyne Huisgen 1,3-dipolar cycloaddition “click” reaction cannot be used for ^{64}Cu -based PET studies for it requires copper(I) metal ion as a catalyst, which also has a tendency to complex with the ligand itself, thus, hampering its efficiency as a PET ligand.

The chelation of the radionuclides to the BFCAs can be performed by two different strategies (**Figure 2.1.10**):

1. Pre-labeling strategy: labeling of BFCAs with radionuclides followed by their attachment to the conjugates like biomolecules or fluorophores.
2. Post-labeling strategy: where the BFCAs are first decorated by the sensitive conjugates (biological vectors or fluorophores) followed by their radiolabeling.

The structure and physical properties of the radiometal-chelator complex have a large impact on the overall pharmacokinetic properties of the radiopharmaceutical construct, with many radiometal complexes being very hydrophilic and subsequently leading to rapid renal excretion when attached to small vectors like peptides and monoclonal antibodies. Linkers are often utilized to connect the biomolecule to the BFCA backbone, which helps to further tune the overall lipophilicity, charge and aqueous solubility of the radiolabeled conjugate, and thus the inclusive biodistribution and pharmacokinetic properties.

Figure 2.1.10: Radiolabeling and bioconjugation strategies for the preparation of radiolabeled BFCA–biomolecule conjugates along with the choice of radionuclides for diagnostic and therapeutic applications.

2.1.2 Optical imaging

Optical imaging is a method of obtaining biological information by signals emitted from cells, tissues and organisms using optical detection techniques combined with optical detection molecules like inorganic nanomaterials or organic fluorophores. The fluorescent report groups use excitation light to reach a higher excitation level and then emit a longer wavelength visible light to form a biological light source *in vitro* or *in vivo* for detection.

An impressive example of multimodal fluorescent probe, demonstrating the potential of nanoscale systems, has been reported by Kobayashi *et al.* A generation 6 PAMAM dendrimer was simultaneously equipped with 120 DTPA (diethylenetriaminepentaacetic acid) molecules for SPECT imaging (^{111}In) and five NIR dyes (Cy5, Alexa 660, 680, 700, 780) four molecules each. These dual-labeled probes allow the simultaneous visualization of five distinct lymphatic drainages (**Figure 2.1.11**).

Figure 2.1.11: *In vivo* dual-modal, five-color lymphatic drainage imaging with the ability to visualize five distinct lymphatic drainages.

(a) *In vivo* multiexcitation spectral fluorescence (right) and *post mortem in situ* radionuclide (left) images of a mouse injected with five distinct G6-(Bz-DTPA)119-(NIR)4-(Bz-DTPA-111In)1 nanoprobe intracutaneously into the middle digits of the bilateral upper extremities, the bilateral ears, and at the median chin, as shown in the schema (mouse 7 in Table 1). Five primary draining lymph nodes were simultaneously visualized with different colors through the skin in the *in vivo* spectral fluorescence image and are more quantitatively seen in the radionuclide image. (b) *Ex vivo* spectral fluorescence and radionuclide images of resected eight draining lymph nodes correlate well to the *in vivo* imaging. (c,d) The imaging results were consistent for all mice examined. Reprinted from [119]

In recent years, fluorescence techniques have been extensively used for studying molecular biology as well as the metabolism of small molecules in small animal *in vivo* and *ex vivo* studies [120]. The resolution of fluorescent probes is high for *in vitro* systems. The near-infrared fluorophores (NIRF) emitting at wavelengths 600 – 900 nm provide a better resolution inhibiting the autofluorescence originating from the tissues. Two fluorescent dyes, ICG Indocyanine Green (ICG) and methylene blue have thus, been approved by the FDA and EMA and have been widely used for animals and also started to be used for human applications for optical imaging of tumors (**Figure 2.1.12**).

Figure 2.1.12: FDA and EMA-approved fluorescent dyes for *in vitro* and *in vivo* imaging.

The NIRF employed in this thesis are sulfo-Cyanine 5 and IRDye CW 800 that provide the necessary penetration depth for *in vivo* studies in mice, and their emission wavelengths also does not overlap with the auto-fluorescence of the tissues (**Figure 2.1.13**) [121, 122].

Figure 2.1.13: NIRF dye derivatives (IRDye 800CW was used for this research work).

2.2 Multimodal molecular imaging

Multimodal imaging is a technique that combines two or more non-invasive molecular imaging technologies which provides a more efficient and convenient system for obtaining further information in disease diagnosis, treatment and monitoring compared to unimodal molecular imaging [123, 124]. With its rapid technological advances, pre-symptomatic detection, targeted therapy, and personalized medicine may be possible in the near future using multimodal molecular imaging [12, 125-132]. The development of multimodal imaging can be directed in two ways; either by developing exogenous techniques, which is the combination of several imaging technologies as a single identity or by developing endogenous probes which brings several imaging agents under one platform (**Figure 2.2.1**) [133].

Figure 2.2.1: Representation of the development of medical imaging technique towards multimodal imaging.

Structural imaging: X-ray, CT, MRI, US; Functional imaging: SPECT, PET, OI, MRI

The synergistic combination of PET and OI systems by combining radionuclide and fluorescent labeling to a NP platform can be utilized for pre-clinical trials to detect cancer tumors. The combination of the direct visualization of OI with the tomographic 3D properties, quantification and whole body penetration of PET can be further employed for fluorescence-guided surgeries and personalized cancer treatment. An interesting example of cancer tumor diagnosis by dual-labeled SPECT/CT and OI modalities is represented in **Figure 2.2.2**.

Functional NPs, in this regard have gained considerable attention for their ability to graft a multitude of functionalities, such as targeting units, solubility enhancers, fluorescent tags and chelating agents, onto their surface. Hence, higher sensitivity may be achieved compared to small molecules, and there is an almost infinite variability regarding the surface functionalization. Their diagnostic, sensor, imaging, vaccine and drug delivery applications provide an effective bridge to connect NPs to biological systems [85, 134-138]. Si NPs, in this regard, with their tunable size and surface properties, extreme hydrophilicity and less cytotoxicity make potential candidates for multimodal imaging.

Figure 2.2.2: SPECT/CT and NIR imaging for cancer tumor diagnosis.

Mice bearing human breast cancer xenograft (MDA-MB-231) were injected with 2 nmol of the conjugate DLIA-IL-11RR *i.v.* Representative images show results 24 h after injection. Tumors (arrows) show more conjugate uptake than muscle in both SPECT/CT and NIR images. High signal intensity in kidneys showed the excretion route of this agent even at very low injection dose. Reprinted and modified from [139]

2.2.1 Silicon nanoparticles as multimodal imaging agents²

Silicon-based NPs hold great promise for wide-ranging optoelectronic and biotechnological applications because of their several unique merits including excellent optical/mechanical/electronic properties, facile surface modification, and compatibility with imaging technologies [42, 140]. As far as biomedical applications are concerned, to date, the Si NPs have been applied in bioimaging [141-143] and in real-time immunofluorescence imaging [44, 57, 144-147]. For instance, to target cells in a specific way, Si NPs have been successfully coupled with single-stranded DNA [148-150], folic acid [151], sugars [152, 153], peptides [145] and antibodies [146, 147], and several *in vitro* studies have been performed on their cellular uptake [154-156] and cytotoxicity [157-159].

The biodistribution and PET imaging of Si NP-based imaging agents was performed for the first time by Kauzlarich *et al.* in C57BL/6 mice. The dextran-coated, manganese-doped, Si NPs were surface functionalized with a 1,4,7,10-tetraazacyclododecane-1,4,7-

² The terminologies used in the dissertation are according to the ISO/TS 80004-2:2015 (<https://www.iso.org>)

triacetic acid (DO3A) derivative for labeling with ^{64}Cu , enabling PET studies (**Figure 2.2.3**). Owing to their relatively large hydrodynamic diameter (HD) (15.1 ± 7.6 nm), the particles were rapidly cleared from the bloodstream but with considerable liver uptake. To date, only a few biodistribution studies of renal clearable NPs incorporating radiolabels high potential for *in vivo* imaging have been reported [23, 82, 160-164].

Figure 2.2.3: Representation of ^{64}Cu -functionalized dextran-coated silicon quantum dots administered intravenously to the *in vivo* system and a PET image 5 min *p.i.* showing renal clearance and liver accumulation. Reprinted from [40]

Erogbogbo *et al.* [151, 165] encapsulated Si NPs encapsulated in phospholipidic micelles and functionalized them with specific peptides for targeted *in vivo* cancer imaging and sentinel lymph node mapping. Examples of Si NP-based bimodal probes have been described by Kauzlarich *et al.* [45, 166], where iron or manganese doping enabled simultaneous MRI and OI. Blue- and red-emitting Si NPs, prepared *via* biomimetic synthesis from diatoms, were very recently reported for dual-color immunofluorescent cellular imaging as well as *in vivo* imaging in *C. elegans* worms [167]. Notably, Sailor *et al.* reported that porous Si NPs (~ 100 nm) could be biodegradable *in vivo* in a relatively short period of time (~ 8 h) and the biodegraded product, orthosilicic acid, is biocompatible to numerous tissues and is readily cleared *via* renal clearance [41, 42]. In a recent study of the biodistribution of Si NPs in a healthy mouse, Prasad *et al.* found that Si NPs of size < 15 nm (micellar and porous) are prone to degrade in the liver and spleen within two months [168].

To summarize, the most interesting features borne by the ultrasmall Si NPs to behave as promising imaging agent are:

1. Tunable intrinsic photoluminescence - depends on the morphology and surface composition [169];
2. Chemical stability - delivered by the covalent Si-C bond;
3. Functionalizable surface groups;
4. Biocompatibility, biodegradability and nontoxicity for *in vitro* and *in vivo* systems [41, 42, 170, 171].

2.2.1.1 Synthesis of ultrasmall Si NPs

Various high efficacy and workable synthesis strategies have been introduced recently for the bulk synthesis of highly fluorescent, chemically robust and photostable Si NPs. The particles can be produced either by the electrochemical (top-down) approach or the wet- chemical (bottom-up) approach [26, 172-176]. Usually, the top-down methods produce samples of higher purity but, with the bottom-up techniques the control on the size and surface termination of the NPs is much higher. In recent years, the bottom-up strategy for developing Si NPs has been widely pursued. An example for each kind of bottom-up synthesis is shown in **Figure 2.2.4**. The major modes of synthesis are the reduction of the silicon precursor, oxidation method, microwave assisted oxidation and reduction reactions and the thermal reduction of silsesquioxanes [177-193]. All these reactions involve the reduction or oxidation of a silicon precursor in the presence or the absence of a surfactant, under inert conditions, to produce hydrogen or halogen-terminated Si NPs.

In most cases, the obtained NPs are further functionalized through a hydrosilylation step, catalyzed by a platinum catalyst, UV irradiation, or through a reaction with alkyllithium reagents. The type of surfactant as well as the reducing agent used for the synthesis can dramatically influence the final size of the produced NPs.

In conclusion, bottom-up techniques provide the opportunity to functionalize the surface of the particles with different functional groups ranging from alkyl or alkenyl chains to amine, epoxy-, alkoxy-, and azide-groups. Usually the size of particles obtained with these methods is very small (< 10 nm), and the emission properties can be tuned from blue to red according to the synthetic method used.

Reduction of a Silicon precursor **Oxidation of a Silicon precursor** **Microwave-assisted synthesis** **Thermal reduction of Silsesquioxanes** **Figure 2.2.4: Synthesis strategies of ultrasmall Si NPs.**

Reduction of a silicon source (X = OMe, Cl, Br; reducing agent = LiAlH_4 , NaBH_4 , LiBH_4 or superhydride; Surfactant = Tetraoctyl ammonium bromide (TOAB), Pentaethylene glycol monododecyl ether (C12E5), Tetrabutyl ammonium bromide (TBAB), Cetyltrimethyl ammonium bromide (CTAB), Didodecyldimethyl ammonium bromide (DDAB), Dodecyltrimethyl ammonium bromide (DTAB) or Tetradodecyl ammonium bromide (TDAB); R = amine, alkyl chain, epoxy group, alkenyl chain); Oxidation of a silicon source (R = alkyl chain, alkenyl chain); Microwave-assisted synthesis; Thermal decomposition of hydrogen silsesquioxanes (R = amine, alkyl chain, epoxy group, alkenyl chain). Adapted from [26]

This also suggests that the best methods for synthesis of Si NPs applicable for bioimaging purposes would be the ones that can produce ultrasmall, highly hydrophilic NPs with small size distribution and bioconjugatable surface groups. As a part of this thesis, the microwave assisted hydrothermal synthesis was modified to produce ultrasmall, amine-terminated Si NPs with surface properties desirable for the development of mono-labeled and dual-labeled imaging agents.

2.2.2 Factors affecting the biodistribution and pharmacokinetics of NPs

2.2.2.1 General remarks

The development of NPs is a well-researched area for the last twenty years, although we still have very limited knowledge of the *in vitro* and *in vivo* behavior of this system. For the NPs to behave as perfect imaging agents, its morphology and physicochemical properties play a significant role. The *in vivo* biodistribution pattern dramatically changes upon the variation in structure, size, shape, surface charge and functionalities, and the density of the construct. These differences are also reflected in the pharmacokinetics (e.g. its circulation time, clearance, intracellular fate, and its organ accumulation), the toxicity, targetability and clearance mechanism of the NPs. The challenge lies in difficult characterization methods, synthetic reproducibility and the multiple combinations of factors such as the size and charge and surface groups on the NPs influencing the nano-bio situations [26, 194-198].

Size

The NPs of size between < 6 nm efficiently pass through the glomerulus in the kidneys and thus are cleared from the system by renal filtration while the particles with larger HDs tend to clear *via* mononuclear phagocytic cells (MPS). MPS include monocytes, which circulate in the blood, dendritic cells and macrophages, like Kupffer cells, which are present in tissues like liver, spleen, lungs, and bone marrow. These cells rapidly sequester particulate matter with HDs larger than approximately 100 nm. A large fraction of these NPs also ends up being recognized and engulfed by macrophages as a result of “opsonization”. This involves the nonspecific adsorption of various serum proteins (immunoglobulins, apolipoproteins, clotting factors, and complement proteins) to the surface of the NPs, which eventually leads to the complete loss of targeting capabilities [199-201].

Surface charge

Charged NPs are reported to induce cytotoxicity by causing a disruption in the cell membrane, thus, hampering the integrity of the cell. Such NPs also tend to interact with the serum proteins due to the electrostatic interactions. Modification strategies such as PEGylation, acetylation, succinylation or surface coating of NPs with amphiphilic

zwitterionic coatings have been applied to neutralize the surface charge, modulate the toxicity of the NPs and prevent its coating with proteins.

Protein Corona

NPs used as targeting agents, upon intravenous administration, come in contact with blood as a complex media made up of salts, sugars as well as serum proteins which mask the NPs and form a shell around it known as the 'protein corona'. The formation of this adduct can cause an increase in the HD, leading to the reversible or irreversible effects on the protein conformation, or the NP surface, its targetability and can induce toxic effects on cells/organs or even apoptosis. Various studies are being conducted to identify the protein corona, specific proteins associated with NPs, to correlate various physicochemical properties to the binding and to understand the underlying mechanism (**Figure 2.2.5**) [197, 202-210].

Figure 2.2.5: Schematic drawing of the structure of nanoparticle-protein complexes in plasma. The “core” nanoparticle is surrounded by the protein corona composed of an outer weakly interacting layer of protein (left, full red arrows) rapidly exchanging with a collection of free proteins and a “hard” slowly exchanging corona of proteins (right). Reprinted from [211]

2.2.2.2 Elimination of the NP-based imaging agents from the body

Renal Clearance

Rapidly eliminating diagnostic agents are preferred over the constructs that tend to reside in the body for longer period of time. The major and more preferred pathway for such artificial constructs is the renal elimination mechanism, which involves minimum

cellular interaction [26, 201, 212-216]. The critical size (HD) for a particle to be renally excretable is ≤ 6 nm. Particles larger than this size prefer the hepatic mode of elimination. Surface-related properties like overall charge and plasticity also play an important role in determination of the fate of the particles. **Figure 2.2.6** demonstrates the mechanism of renal clearance of the intravenously administered NPs from the body [153, 201, 214, 217, 218]. Renal clearance mechanism involves the glomerular filtration of the NPs, passing through the fenestra present on negatively charged glomerular basement membrane. The fenestrae are the nanosized pores which allow a definite sized molecule to pass through it. The glomerular basement membrane being negatively charged, facilitates a faster elimination of the neutral and positively charged molecules in general, and also positively charged NPs compared to its negatively charged equivalents. This is probably the primary reason for the kidney retention of the anionic conjugates. This was confirmed by a study reporting the renal threshold of anionic NPs to be ~ 3.7 nm when administered intravenously to the *in vivo* system. NPs larger than the threshold tend to stay in the kidneys for a long-term leading to nephrotoxicity and radiation exposure side effects in case of radiolabeled NPs [219-223].

Figure 2.2.6: Section of kidney cortex: Typical biodistribution behavior and renal clearance of ultrasmall NPs in kidney tissue.

The NPs below the renal threshold *in vivo* hydrodynamic diameter of < 6 nm are filtered through the glomerulus. Positively charged NPs with *in vivo* diameter of 6-8 nm are more readily filtered than their equally sized counterparts possessing anionic surfaces due to electrostatic repulsion by negative charges of the glomerular basement membrane. Based on the charge of the NPs, they can either be sustained in the kidney cortex or can be resorbed through the tubular fluid to the proximal tubule or filtered out to the collecting duct in the form of urine.

Hepatic Clearance

Liver is the primary hepatic organ responsible for the catabolism and biliary excretion as well as the elimination of all foreign substances in the organism. The NPs larger than the threshold to eliminate *via* renal pathway are considered for hepatobiliary mode of excretion. After systemic administration, these nanomaterials are trapped in the reticuloendothelial system organs such as liver and spleen. The bigger NPs in circulation are directed towards the liver with a dual vascular supply with arterial (hepatic artery) and well as venous (portal vein) blood mixing in the fenestrated liver capillaries called as the sinusoids. The hepatocytes (endothelial cells) or Kupffer cells (macrophages) present here engulf the NPs leading to endocytosis. NPs endocytosed by the hepatocytes are potentially processed and passed into the bile, to the intestines, and excreted in the form of feces. Other NPs that do not get taken up by these cells rejoin the systemic circulation, redistribute, and may return to the MPS organs over the subsequent pass and therefore, cleared from the blood stream over time [26, 214, 224].

Even after severe research investigations, it is still a challenge to designate an 'ideal NP size, geometry or surface functionality' for biomedical applications. It is crucial to study each set of NPs for their specific *in vivo* behavior, pharmacokinetics and interactions within the system to gather more detailed information required for developing more effective imaging agents.

2.2.3 Tumor targeting strategies for molecular imaging agents

Considering the NP-based approaches to target cancerous tumors, a series of biological vectors have been reported to diagnose and bind specifically to the tumors. According to the reported investigations, it is clear that the accumulation of the targeting agents in the tumors could be either *via* passive or active targeting.

2.2.3.1 Passive targeting (Passive accumulation)

The ability of the NPs with sizes between 30–200 nm to target solid tumors is referred to as passive targeting [26, 199, 225]. This owes to the enhanced permeability and retention (EPR) effect that occurs due to the poorly organized vasculature of the solid tumors lacking in structured fenestrations due to the high proliferation of cancer cells. This "leaky" vasculature, combined with poor lymphatic drainage, promotes the passive accumulation of NPs into it, providing the opportunity to concentrate reasonably sized

NP-based therapeutics and diagnostics within cancerous tissue (**Figure 2.2.7**) [21, 199, 226-229].

2.2.3.2 Active targeting

The two main strategies for targeted cancer imaging can be illustrated as:

- Tumor cell targeting
- Tumor vasculature targeting

Tumor cell targeting can be achieved by conjugation of specific biological vectors (e.g., antibodies, peptides, small molecules) to NPs. These conjugates undergo tumor recognition and selective binding to the receptors that are overexpressed on a certain tumor cell surface (e.g., epidermal growth factor receptor (EGFR), human epidermal growth factor receptor-2 (HER-2), somatostatin receptor-2 (SSTR-2) [122, 229].

The high surface area-to volume ratio of NPs allows a high local density of the modified NPs to enable better targeting efficiency. However, tumor cell targeting can only work if the functionalized NPs are able to effectively reach the tumor cell surfaces after extravasation across the vasculature endothelium. The general rules governing the extravasation of NPs still remain largely unknown, since NPs with near identical surface coating and charge could display surprisingly different extravasation behavior *in vivo*. Even if NPs with suitable shape, size and surface modification could efficiently extravasate, they still must overcome many biological barriers (e.g., high interstitial fluid pressure, dense collagen matrix) and must penetrate tens to hundreds of micrometers through the vasculature endothelium and cell layers before reaching the tumor cell surface and binding to the target of interest. All of these challenges explain why most tumor cell targeted NPs show highly targeting efficacy *in vitro*, but rarely exhibit significantly higher uptake *in vivo* in solid tumors when compared with the non-targeted counterparts (**Figure 2.2.7**) [21, 26, 225].

Tumor vasculature targeting

Tumor vasculature targeting is an alternative strategy, which could be applied to a wide variety of well-functionalized NPs regardless of tumor types. This method has several benefits over directly targeting the tumor cell. Firstly, all solid tumors depend on angiogenesis to receive sufficient blood flow. Therefore, targeting the tumor vasculature (i.e., targeting receptors overexpressed on the tumor vascular endothelial cells) could be an effective strategy for most solid tumors *in vivo*. Secondly, while tumor cells maybe far

away from blood vessels, tumor endothelial cells are directly exposed to circulating blood, which can greatly facilitate the binding of functionalized NPs without the barrier of extravasation [230]. Vasculature targeting of tumor angiogenesis can be achieved using biomarkers on the surface of tumor endothelial cells. Vasculature targeting moieties, including vascular endothelial growth factor (VEGF) and RGD (Arg-Gly-Asp) peptides, have been conjugated to the surface of different NP such as quantum dots (QDs), single-walled carbon nanotubes (SWNT), and graphene oxide (GO) for enhanced tumor targeted imaging and drug delivery [21, 26, 225].

Figure 2.2.7: Difference between the active and passive targeting of tumors.

(Left) In normal healthy tissue, endothelium of the vessels is quite impermeable for nanomaterials. Leaky fenestrated endothelium in the tumor bed allows the particles to diffuse and accumulate primarily due to the enhanced permeability and retention effect. (Right) In active targeting, the extravasated particles due to the presence of targeting vectors facilitate the interaction with the overexpressed receptors on the tumor cells resulting in an enhance accumulation and preferential cellular uptake through a receptor-mediated process.

As a conclusion, we can say that the contribution of NP-based passive targeting (EPR-mediated) as well as active tumor targeting (*via* active receptor-ligand interaction) leads to an effectual diagnosis of the cancer tumors during pre-clinical and clinical trials but not much is known about the biological fate of the administered NPs. The extent of tumor targeting which is majorly determined by factors like size, shape and surface functionalization of the NPs is still an area of research.

2.2.4 Targeted functionalization of NPs and BFCAs for imaging purposes

Following the active tumor targeting strategy, selected tumor biomarkers are needed to facilitate effective localization of the multi-functionalized NPs in the tumor tissue and to

minimize unspecific uptake in non-targeted organs. A variety of targeting molecules including receptor-specific peptides, aptamers, carbohydrates as well as antibodies, their fragments and other proteins have been exploited for active tumor targeting of multimodal NPs.

Camelid Single Domain Antibodies for EGFR Active Targeting

The first choice for this purpose were nanobodies or single domain antibodies (sdAb) that are based on the single variable domain of heavy chain antibodies (HCAb – VHH) devoid of light chains [96, 121, 231, 232]. These are amongst the smallest known antibody-binding fragments (13-15 kDa) that can retain full functional advantages of the conventional antibodies (**Figure 2.2.8**). They are endowed with unique properties such as low immunogenicity, high solubility, tailored shelf-life and high specificity. With a diameter of only 2.4 nm and height of 4 nm, they are excellent choice as EGFR specific cancer biomarkers with their ability to recognize uncommon or hidden epitopes, high tumor binding avidity and penetration potential followed by fast circulation kinetics [233, 234]. The sdAb are also reported to exhibit high stability in variable temperature and pH. Recently, an economic production of these nanobodies *via* microorganisms has also been reported by our laboratory [235].

Figure 2.2.8: Smallest known fully functional camelid antibody- fragment (Nanobody) derived from heavy chain only antibodies.

(VHH- variable domain of the heavy chain of heavy-chain antibodies (HcAb))

Somatostatin analogues (Octreotate, Octreotide) for Somatostatin receptor (SSTR) Active Targeting

Somatostatin receptors (SSTR) are known to be expressed in a large number of human neuroendocrine tumors and represent the basis of *in vivo* targeting in humans. The crucial conditions for the targeting of tumor cells with a high density of SSTR shapes to a sufficiently long biological half-life, the possibility of labeling with various radionuclides and, in particular, the ability of SSTR-positive tumor cells to specifically bind and internalize radiolabeled SST analogues [236-242]. Several SST analogues labeled with γ -emitters such as ^{111}In -labeled pentetreotide, ^{111}In -labeled DTPA-octreotide (Octreoscan®) or $^{99\text{m}}\text{Tc}$ -labeled depreotide (NeoTect™) are commercially available and clinically successfully used for diagnosing different types of tumors and their metastases. The therapeutic application of SST analogues labeled with β -emitting particles (^{177}Lu , ^{90}Y) is of major importance for such endocrine tumors. Alongwith its high therapeutic potential, it known to be accompanied by occurrence of delayed radiation-induced toxicity to kidney, bone marrow and liver [242-244].

The research idea was to improve the pharmacological behavior by preparing bispidine-based TATE (Tyr3-Octreotate) conjugates that can behave as a targeted PET diagnostic agent. This may ensure a positive tumor detection as well as a rapid renal elimination from the *in vivo* system minimizing the chances of nephrotoxicity and low accumulation of the biomarker in the vital organs of the system. A TATE peptide modified with oxyethylene groups to increase hydrophilicity for tumor targeting is shown in **Figure 2.2.9**.

Figure 2.2.9: Structure of EG-TATE.³

³ EG in this context is an acronym for diethylene glycol.

Chapter 3 Results and Discussion

The experimental work done for this dissertation can be conveniently divided into following sections:

Chapter 3.1

This section constitutes the design and synthesis of novel bispidine-based BFCAs. It involves the modification of the bispidine core at the C9-position, allowing the conjugation to a series of sensitive biomolecules and to the Si NPs. The optimized synthetic procedures, purification and characterization data of the novel derivatives has been reported in this section. The ^{64}Cu complexation kinetics to determine the radiolabeling efficiency of the bispidines is also stated here.

Chapter 3.2

This section ascribes the bioconjugation of isothiocyanate-functionalized and dibenzocyclooctyne-functionalized bispidines with receptor-specific TATE peptide and sdAb 7C12 respectively, for active cancer tumor targeting *via* PET imaging. The synthetic protocols, purification strategies and characterization studies followed by preliminary radiolabeling and cytotoxicity studies of the bispidine bioconjugates has been encompassed in this section.

Chapter 3.3⁴

This section is dedicated to the synthesis of multifunctional amine-terminated Si NPs. The hydrothermal synthesis of ultrasmall Si NPs along with the characterization data is described in this section.

Chapter 3.4

Towards the development to mono-labeled imaging agents, this section deals with the modification of the Si NP surface with PET or OI modular ligands. PET imaging agents were developed by coupling the Si NPs to the isothiocyanate-terminated bispidines and NOTA BFCAs respectively. OI agents, on the other hand, were produced by conjugating the fluorophores, taking advantage of the active NHS-ester chemistry. The synthesized

⁴ The synthesis and characterization of the Si NPs were performed in the group of Prof. Luisa De Cola (University of Strasbourg).

Si NP conjugates were characterized by unique methods and evaluated for their cytotoxic and biodistribution behavior.

Chapter 3.5

In this section, the amine-terminated Si NPs were modified to develop a distinct dual-labeled imaging agent. Si NPs were conjugated to the IR 800 fluorophore and NOTA BFCA concurrently. The reaction synthesis was optimized by applying two different synthesis strategies: one pot method and stepwise synthesis. The Si NP-IR800-NOTA was meticulously characterized and further evaluated by cell and biodistribution studies. PET and OI molecular imaging techniques were used for the *in vitro* and *in vivo* examination.

Chapter 3.6

In this section, both, mono-labeled and dual-labeled fluorescent Si NPs were coupled to a chimeric anti-platelet factor 4, 5B9 peptide. The synthesis of the targeted Si NPs bioconjugates *via* a bifunctional linker (PEG) to produce Si NP-IR800-PEG-5B9 and Si NP-IR800-NOTA-PEG-5B9 is reported here.

Chapter 3 Results and Discussion

3.1 Synthesis, characterization and radiolabeling studies of hexadentate bispidine-based bifunctional chelating agents

Bispidine-based scaffolds have been extensively explored for radiopharmaceutical applications because they can form radiometal complexes with high kinetic inertness and thermodynamic stability [102, 113, 245]. In particular, there continues to be considerable interest in studying the coordination chemistry of bispidine-derived ligands with a view to developing them as suitable chelators in particular for radiocopper, as these can be used for both imaging (^{64}Cu) and therapy (^{67}Cu) [71, 77, 78, 102, 103, 113, 114, 208, 218, 245, 246]. Recently, novel octadentate picolinic acid-based bispidine ligands have been also proposed for radiopharmaceutical applications using ^{111}In , ^{177}Lu and ^{225}Ac [113]. The literature serves a good amount of data for such bispidine cores, confirming their facile synthesis, large synthetic variability, as well as the ease of derivatization with biological vectors [247-250]. Unless hindered by a bulky substituent, the bispidine core also adopts the most stable double chair configuration, with *endo-endo* being the favored orientation for the tethered substituent (**Figure 3.1.1**).

Figure 3.1.1: The most stable double chair (cc) configuration of bispidine with *endo-endo* pyridyl-substitution at C2/4.

An interesting example delineating the impact of substituents on Cu(II)-binding efficiency of bispidines compares two isomeric pentadentate derivatives, with pyridyl groups positioned at the N3 and N7 positions respectively (**Figure 3.1.2**) [102]. The pyridyl-substitution at N7 results in higher stability of the Cu(II) complex in comparison to the one featuring N3-positioned pyridyl arm ($\log K = 18.31$ vs 15.66 for the N3 derivative). A probable reason for this chelation behavior is that the Cu(II) coordination sphere remains unaffected by the N3 substituent which contributes minimally for the radiometal chelation.

Although, the experimental and computational studies confirm that for biological applications, the hexadentate (N3 and N7 dipyridyl-substituted) ligands are a preferred choice due to better encapsulation of the metal ion in the coordination sphere under physiological conditions.

Figure 3.1.2: The N3 and N7 substituted pentadentate bispidine isomers showing prominent differences in their stability constants for Cu(II) complexes.

Interestingly, synthetic modification of the bispidine scaffold at the C9 position remains largely unexplored. This is despite the knowledge in hand that the C9-positioned substituents bear no direct influence on the Cu(II) coordinating environment. The X-ray crystal structure of tetrapyridyl-bispidol (C2/4, N3/7 substituted) confirms that the C9 position is remote to the coordination site of the metal atom and does not interfere with any electron-donating group participating in the chelating process (**Figure 3.1.3**) [218].

Figure 3.1.3: Coordination environment around the Cu(II) ion in Bispidol.

Thermal ellipsoids drawn on a 30% probability level. Gray spheres denote carbon, red spheres oxygen, blue spheres nitrogen, and yellow spheres copper atoms. The overlay with the space-filled model shows the efficient encapsulation of the copper ions by the ligand. Reprinted from [76]

A discrete example for C9-modified bispidine involves the reduction of the keto group to form the corresponding bispidol, which was then linked to a boron-dipyrromethene (BODIPY) derivative, propargylamine or a bombesin (BBN) peptide sequence using a 4-nitrophenyl bispidine carbonate derivative. Continued interest in bispidine-based bifunctional chelators prompted us to explore the possibility of developing hexadentate bispidines, modified at C9 position to provide a site for optional attachment of vector molecules and/or fluorescence tags.

This chapter describes the synthesis and detailed spectroscopic characterization of a series of hexadentate bispidine derivatives and their bioconjugates, together with the assessment of their ability to form complexes with copper radionuclides for potential application in cancer diagnosis (PET imaging) and therapy (**Chapter 3.1 and 3.2**).

3.1.1 Synthesis of hexadentate bispidine-based ligands

3.1.1.1 Synthesis and characterization of Piperidone (**B1'**), Bispidone (**B1**) and Bispidol (**B2**) precursors⁵

The synthesis of bispidine scaffold follows two consecutive Mannich reactions (**Experimental section 5.4.1**). A slightly modified synthetic route was applied for synthesizing hexadentate tetrapyridyl-bispidines (**Scheme 3.1.1**). In the first step, 2-picolylamine (a) and 2-pyridinecarboxyaldehyde (b) react to generate an imine intermediate. The α -hydrogen of the dimethyl-1,3-acetonedicarboxylate (c) facilitates its keto-enol tautomerism, which reacts with the freshly formed imine. The reaction occurs twice for primary amines and furnishes the product Piperidone (**B1'**), which is a β -amino-carbonyl compound and also known as the Mannich base. The second reaction enables the formation of a rigid cyclonane ring when after Piperidone is attacked by the new imine formed by the reaction of formaldehyde (d) and 2-picolylamine (a). The product formed after the completion of two Mannich reactions is termed as Bispidone (**B1**) with a ketone at C9 position. The ¹³C NMR and FTIR spectra of **B1** is reported as **Figure 3.1.8** and **Figure 3.1.9**.

⁵ The ESI-MS and ¹H NMR spectra of **B1'**, **B1** and **B2** are reported as supporting information Annex (**Figure 7.1-7.4**)

Scheme 3.1.1: The synthetic scheme of Bispidone (B1) via double Mannich reaction with Piperidone (B1') as an intermediate.

(a) 2-picoylamine, (b) 2-pyridinecarboxaldehyde, (c) dimethyl-1,3 acetonedicarboxylate, (d) formaldehyde).

The bispidone derivative once formed has a tendency to undergo a retro-Mannich reaction that leads to the formation of epimers by the ring cleavage and rotation of C-N3 bond as represented in **Scheme 3.1.2**.

Scheme 3.1.2: Retro-Mannich reaction caused by the carbonyl at C9 that might lead to the formation of epimers.

To avoid this situation, the carbonyl group at the C9 was subjected to NaBH₄-mediated reduction to form the corresponding bispidol (**B2**), prior to any further chemical modification. The reduction procedure was slightly modified from the literature data. Product **B2** was purified by recrystallization as a white crystalline powder (**Scheme 3.1.3**).

Scheme 3.1.3: Reduction of the C9 carbonyl of **B1** to secondary hydroxyl group in **B2**.

The reduction of keto group at C9 position was confirmed by appearance of a signal for the hydroxyl carbon at 72 ppm in the ^{13}C NMR spectra, along with the disappearance of carbonyl resonance at 203 ppm (**Figure 3.1.8**). Moreover, a peak at m/z 596.2 (calculated m/z 594.26) corresponding to $[\text{M}+\text{H}]^+$ ion was the observed in the ESI-MS of **B2** (**Figure 3.1.4**). As expected, the FTIR spectra showed a strong O-H stretch for the hydroxyl group of **B2** at 3500 cm^{-1} in **Figure 3.1.9**.

The literature corroborates **B2** as a hexadentate ligand for $^{64}\text{Cu}(\text{II})$ radiotracers and its applicability for PET. The stability constant for the $\text{Cu}(\text{II})$ -**B2** complex has been experimentally determined and reported to be $\log K$ 16.28 [102].

Figure 3.1.4: ESI-MS spectra of **B2** showing the $[\text{M}+\text{H}]^+$ ion peak at m/z 596.2.

3.1.1.2. Synthesis and characterization of C9-modified novel B2 derivatives B3–B5

Bispidine amine derivative B3⁶

B2 as a precursor was modified with a halo-alkyl amine linker at the C9 position to create an amine-functionalized bispidine-based bifunctional chelating agent, which can be conjugated to isothiocyanate or active NHS-ester terminated biological vector and can be utilized as targeted PET imaging agents.

Propyl amine was the linker of choice to modify the hydroxyl group at C9 position of **B2**. The terminal amine of the bromo-propyl amine was protected by *tert*-butoxycarbonyl (Boc) protecting group to develop **L1** (**Figure 3.1.4**). A white crystalline product obtained in 70% yield after column chromatography was characterized by NMR. The ¹H NMR spectra displayed a characteristic singlet at δ 1.3 ppm for the *tert*-butyl protons of **L1** (**Figure 3.1.5**).

Scheme 3.1.4: The two-step synthesis scheme of the B3.

(i) Reaction of **B2** to **L1** to form Boc-protected **B3'**, (ii) The deprotection reaction leading to the amine-terminated **B3**.

The reaction between **B2** and **L1** lead to the formation of a Boc-intermediate (**B3'**) as a result of nucleophilic substitution reaction. The hydroxyl group at C9 of **B2**, under basic conditions, attacks the brominated carbon, which is electrophilic in nature. Bromide ion, a good leaving group quickly cleaves off furnishing the **B3'**. The work-up of the crude product was performed to extract the product in chloroform by liquid-liquid extraction

⁶ The synthetic optimization of the precursor **B3** was performed by Sebastian Hunoldt and Carmen Rühl at Helmholtz-Zentrum Dresden - Rossendorf.

against water. The organic layer was evaporated to extract a brown-oily product in 35% crude yield. The intermediate **B'** was analyzed by NMR and mass spectroscopy.

The ^1H of crude **B3'** depicted two peaks for the *tert*-butyl protons from **L1** and **B3'** between δ 1.4–1.3 ppm (**Figure 3.1.5**). ESI-MS confirmed the identity of **B3'** with a $[\text{M}+\text{H}]^+$ ion at $m/z = 751.8$ (calculated m/z 751.37) (**Annex Figure 7.6**).

Figure 3.1.5: Comparative ^1H -NMR spectra (aliphatic region) of **L1**, **(B3'+B3)** and **B3** with labeled characteristic signals.

B3' was deprotected under acidic conditions by trifluoroacetic acid (TFA) to furnish **B3**. Mass spectrometry with a $[\text{M}+\text{H}]^+$ ion at $m/z = 651.4$ (calculated m/z 651.32) confirmed

the complete deprotection of the amine and the identity of product **B3** (**Annex Figure 7.7**). The product was purified by preparative reverse-phase HPLC (**“Method B” Section 5.2**) ($R_t = 11.5$ min) (**Figure 3.1.7**) and characterized by ^1H and ^{13}C NMR and FTIR spectroscopy.

A comparative NMR spectra of **A**, **B3'** and **B3** is displayed in **Figure 3.1.5**. ^1H NMR spectra of **B3** shows no peaks corresponding to *tert*-butyl protons from the Boc-group confirming a successful deprotection of **B3'**. An appearance of a singlet at δ 4.3 ppm for **B3'** supports the functionalization *via* C9 hydroxyl moiety and its characteristic shift to δ 4.75 ppm for **B3** confirms the deprotection of the primary amine. The protons from the propyl chain on **B3** appear in the range of δ 3.7–1.9 ppm (**Figure 3.1.6**).

Figure 3.1.6: ^1H -NMR spectrum of **B3** (measured in CD_3CN).

The FTIR spectrum depicts a strong N-H stretch between 3500 to 2900 cm^{-1} . The carbonyl vibration of esters ($\text{O}=\text{C}-\text{O}$) at C1 and C5 positions is recorded between 1800 to 1750 cm^{-1} . N-H bending peaks are observed from 1160 to 1130 cm^{-1} and the ether group (C-O stretch) is observed at 1200 cm^{-1} (**Figure 3.1.9**). This data thus confirms the successful synthesis of **B3**.

The efficiency to encapsulate ^{64}Cu and conjugate to biological vectors with compatible functionalities like active succinimide esters, isothiocyanates and the carboxylic acids makes **B3** a potential BFCA for imaging and radiopharmaceutical applications. Therefore, after the successful synthesis and analysis of **B3**, it was used as a precursor for developing additional derivatives **B4-B9**, with varying functional moieties

complementary to the functionalities present usually on the delicate biomolecules like the antibodies and peptides (**Scheme 3.1.5**).

Figure 3.1.7: Comparative RP-HPLC chromatograms for B1–B5 with their respective retention times (R_t).

(RP-HPLC (Method B) Gradient 30% to 70% B in 30 min; Mobile phase: A = H_2O + 0.1 Vol. % TFA, B = CH_3CN + 0.1 Vol. % TFA; Flow rate = 1 mL/min, λ = 220 nm).

Figure 3.1.8: Comparative ^{13}C -NMR spectra of ligands B1–B4 with labeled characteristic NMR signals.

Figure 3.1.9: Comparative FTIR spectra for B1–B5 with labeled characteristic vibrational frequencies.

Scheme 3.1.5: Modification and functionalization of B3 to develop bispidine-derivatives B4–B9 with different bioconjugatable functionalities.

Bispidine isothiocyanate derivative B4

Scheme 3.1.6: The synthesis of isothiocyanate-terminated B4 by reacting B3 with *para*-phenylene diisothiocyanate.

B3 when reacted with *para*-phenylene diisothiocyanate under basic conditions, the nucleophilic amine attacks the electrophilic carbon of the isothiocyanate, resulting in a stable thiourea linkage (**Scheme 3.1.6**). No leaving group is involved in this reaction. Due to solubility differences, product **B4** precipitates in diethyl ether and can conveniently be extracted. Although, **B4** was also purified by RP-HPLC (**“Method B” Section 5.2**) with $R_t = 26.1$ minutes for a high purity content because of its subsequent applications in biological studies.

The $[M+H]^+$ ion peak at m/z 843.3 (calculated m/z 843.30) confirmed the identity of the isothiocyanate derivative **B4** (**Annex Figure 7.7**). The ^{13}C NMR of **B4** showed an influx

of aromatic carbons in the range of 155–125 ppm and the characteristic resonance isothiocyanate carbon was observed at δ 182 ppm (**Figure 3.1.8**). The FTIR of the **B4** showed the characteristic N=C=S stretch at 2140 cm^{-1} (**Figure 3.1.9**).

Bispidine dibenzocyclooctyne derivative B5

Scheme 3.1.7: Modification of B3 to B5 by using NHS-DBCO reagent.

Cyclooctynes are useful in strain-promoted copper-free click-chemistry based cycloaddition reactions as they can react with azide-functionalized biomolecules without the need for a Cu(I) catalyst to result in a stable triazole linkage. The absence of exogenous metal catalysts like Cu ions make the reactions suitable for the *in vivo* applications of bioorthogonal chemistry. For instance, Cu ions have a tendency to bind to bispidine that hampers its $^{64}\text{Cu(II)}$ radiolabeling efficiency which is a crucial factor for PET-based imaging.

Cyclooctyne-functionalized bispidine **B5** was synthesized during overnight room temperature stirring of **B3** with a commercially available active NHS-ester analogue of dibenzocyclooctyne (DBCO-NHS ester) (**Scheme 3.1.7**). The crude product **B5** was purified by HPLC (**“Method B” Section 5.2**) with a retention time (R_t) of 26.1 minutes. The high retention time of **B5** compared to the precursor **B3** (R_t = 11.5 minutes) signifies an increase in the lipophilicity of **B5** due to the addition of the aromatic DBCO moiety in the structure.

The mass spectrometric characterization specifies the formation of **B5** by $[\text{M}+\text{H}]^+$ ion peak at m/z 939.1 (calculated m/z 938.41) (**Figure 3.1.10**). The ^1H NMR of **B5** shows a cluster of aromatic peaks between 7.6–7.2 ppm characteristic for protons of the dibenzyl group (**Annex Figure 7.9**).

Figure 3.1.10: ESI-MS spectra of B5 showing the [M+H]⁺ ion peak at m/z 939.1.

3.1.1.3. Synthesis and characterization of C9 modified novel derivatives B6–B9

Thiolated bispidine derivatives B6 and B7

Thiol-terminated BFCA are essential for bioapplications due to their compatibility with maleimide-terminated biological vectors. The coupling occurs *via* Michael type addition reaction. The synthetic scheme for two-sulfhydryl derivatives **B6** and **B7** is reported as **Scheme 3.1.8**.

Scheme 3.1.8: The synthesis of **B6** (one step) and **B7** (two steps) from the precursor **B3**.

B6 was prepared in a single-step by reacting **B3** with Traut's reagent (2-iminothiolane) under basic conditions. The reaction proceeds with the 2-iminothiolane ring opening, leading to the formation of a terminal thiol group with an amidine bond. An interesting result was obtained by mass spectrometry analysis of **B6**. The mass spectrum depicts an intramolecular cyclization side reaction occurring within the molecule **B6** (Scheme 3.1.9).

Scheme 3.1.9: Representation of intramolecular cyclization reaction of **B6** and its determination by mass spectrometry.

The ionization of **B6** eventually leads to the loss of an immonium ion fragment (m/z 17) from the molecule as shown in the mass spectra (Figure 3.1.11). This hypothesis is supported by the literature and also confirms that this side reaction does not affect the coupling tendency of thiol-bearing **B6** to maleimide-based compounds [251].

Figure 3.1.11: ESI-MS spectra of B6 showing the [M+H-17]⁺ and [M-17+2H]⁺⁺ ion peaks at m/z 736.2 and 368.7.

The second thiolated derivative **B7** is synthesized by two consecutive coupling and deprotection reactions. **B7'** intermediate was obtained by stirring **B3** with *N*-succinimidyl-S-acetylthioacetate (SATA) at room temperature for 2.5 hours leading to stable amide bond formation. SATA is a thiolating reagent ensuring the covalent modification of primary amines to protected yet exposable sulfhydryl groups.

B7' with a protected thiol can be stored safely for long period (upto 8 months) and can be treated with hydroxylamine to generate **B7** with the labile sulfhydryl group when needed for further conjugation reactions. The deprotection reaction is performed under aqueous hydroxylamine/EDTA/sodium bicarbonate solution stirring for 3 hours.

The product was purified by HPLC ("**Method D**" **Section 5.2**), followed by freeze drying of the pure fractions to obtained **B7** as a colorless solid in 68% yield. The mass spectrometry of the [M+H]⁺ ion at m/z of 725.5 (calculated m/z 725.30) confirms the identity of **B7** (**Figure 3.1.12**).

Figure 3.1.12: ESI-MS spectrum of B7 showing the $[M+H]^+$ ion peak at m/z 725.5.

Bispidine phenyl amine derivative B8

The idea behind the development of an aromatic amine-derivative of the bispidine is to increase the blood retention times of its targeted bioconjugate (coupled to a peptide or monoclonal antibody) due to a lipophile attached to the amine-terminated bispidine. Shorter blood half-lives of the conjugates and rapid elimination is favored for diagnostic purposes but considering the therapeutic applications, slight body retention is a preferred choice [198, 252].

Scheme 3.1.10: The derivatization of B3 to B8 via a two-step synthesis.

B8 derivative was produced by reacting precursor B3 to a freshly prepared linker L2, in small amount to test the feasibility of the two-step synthesis. The synthesis scheme of

L2 is shown in **Figure 3.1.10**. It was then purified by column chromatography and characterized by NMR spectroscopy. The ^1H NMR of **L2** (**Figure 3.1.13**) displayed a singlet at δ 1.5 ppm which corresponds to the 9 *tert*-butyl protons. The aromatic multiplets were observed between δ 7.5-7.3 ppm.

Figure 3.1.13: ^1H -NMR spectrum of **L2**.

The coupling of the **L2** to **B3** was performed by a 24 hour stirring of the reaction at 85 $^{\circ}\text{C}$. The crude product was confirmed by mass spectrometry and the deprotection reaction was performed *in situ*. The intermediate **B8'** and product **B8** were identified by ESI-MS spectrometry as $[\text{M}+\text{H}]^+$ ion at m/z 899.3 (calculated m/z 899.43) and 799.5 (calculated m/z 799.38) respectively (**Annex figure 7.13**).

Bispidine triglycine derivative B9

The amine-terminated precursor **B3** behaves as a potential BFCA to develop a proficient bioconjugate **B9**, for non-invasive, *in vivo* PET imaging and therapy. The aim was to perform sortase-A mediated site-specific labeling of single-domain antibody-fragments (sdAbs) by the triglycine-functionalized end of the conjugate **B9** (mechanism explained in **Chapter 3.2**) [232]. Radiolabeling and cytotoxicity studies of this conjugate are not a part of this work.

Scheme 3.1.11: Synthesis of bioconjugate **B9 via a two-step synthesis from **B3**.**

The bioconjugate **B9** was prepared by a two-step synthesis strategy (**Scheme 3.1.11**). The first step involves the conjugation of **B3** to a commercially available Fmoc-triglycine peptide by the formation of a stable amide bond. The intermediate **B9'** was purified by RP-HPLC (**"Method C"** **Section 5.2**) and identified by mass spectrometry as $[M+H]^+$ ion at m/z 1045.4 (calculated m/z 1045.4) (**Annex Figure 7.14**).

In the second step, **B9'** was deprotected under basic conditions using 20% piperidine in DMF to obtain the final bioconjugate **B9**. It was also purified by RP-HPLC (**"Method C"** **Section 5.2**). Mass spectrometry confirmed the identity of the **B9** with the $[M+H]^+$ ion peak at m/z 822.6 (calculated m/z 822.38) (**Figure 3.1.14**).

Figure 3.1.14: ESI-MS spectrum of **B9 showing the $[M+H]^+$ ion peak at m/z 822.6.**

3.1.1.4. Synthesis and characterization of C1/5 modified derivatives **B10** and **B11**

The bispidine-derivatives **B10** and **B11** showcase the modifications at C1/5 positions, contrary to the previous C9 modified derivatives (**B3–B9**). These modifications allows the study of the influence of different side chains on the physiological properties of the conjugates for *in vivo* experiments.

Bispidine carboxylic acid derivative B10

The C1/5 dicarboxylate derivative **B10** was hydrolyzed by dissolving **B3** in a cesium hydroxide-tetrahydrofuran solution and stirred for 24 hours at room temperature (**Scheme 3.1.12**). The white emulsion obtained was then acidified by 1M HCl and the solvent was evaporated. Crude **B10** obtained was then purified by RP-HPLC (**Method D” Section 5.2**) and characterized by NMR spectroscopy.

Scheme 3.1.12: Hydrolysis of the C1/5 esters of **B2** to yield dicarboxylic acid derivative **B10**.

In ^1H NMR, a loss of the characteristic 6 proton singlet of the methyl esters confirms the basic hydrolysis of **B3** precursor (**Annex Figure 7.17**). The $[\text{M}+\text{H}]^+$ ion peak at an m/z value of 566.8 (calculated m/z 566.23) confirms the identity of the product (**Figure 3.1.15**).

Figure 3.1.15: ESI-MS spectrum of B10 showing the $[M+H]^+$ ion peak at m/z 566.8.

Bispidine ethyl ester derivative B11

Derivatives B11' and B11 were synthesized to analyze the effect ethyl esters on the pharmacokinetics of the system compared to the usual dimethyl ester substituents present at C1/5 positions of all possible C9-modified derivatives. **B11'** and **B11** were synthesized by using diethyl-1,3-acetonedicarboxylate following the double Mannich reactions (dimethyl-1,3-acetonedicarboxylate used **B1'** and **B1**) (**Scheme 3.1.13**). The preliminary characterization of the product **B11** was performed by ESI-MS with $[M+H]^+$ ion peak at an m/z value of 621.2 (calculated m/z 620.27) (**Figure 3.1.16**).

Scheme 3.1.13: Synthesis of B11' and B11 via double Mannich reactions.
The red bonds represent the difference from derivative B1' and B1.

Figure 3.1.16: ESI-MS spectra of B11 showing the $[M+H]^+$ and $[M+2H]^{++}$ ion peak at m/z 621.27 and 311.25.

Table 3.1.1: Representation of the mass spectrometry calculated vs experimental m/z values of derivatives B1–B11 (including the intermediates B1', B3', B8', B9', B11').

Compound	MS $[M+H]^+$	MS $[M+H]^+$
	Calculated m/z	Experimental m/z
B1'	460.17	461.3
B1	592.24	611.6
B2	594.26	596.2
B3'	751.37	751.8
B3	651.32	651.4
B4	843.30	843.3
B5	938.41	939.1
B6	753.35	736.2
B7	725.50	725.3
B8'	899.43	899.3
B8	799.38	799.5
B9'	1044.45	1045.4
B9	822.38	822.6
B10	566.23	566.8
B11'	488.21	489.1
B11	620.27	621.2

3.1.2 Radiochemistry of modified bispidine derivatives

3.1.2.1 Radiolabeling of precursors **B1** and **B2**, and BFCAs **B3 – B5**

The conjugation of the synthesized bispidine-based BFCA to a targeting agent is a pivotal step in the generation of high quality imaging tracers and therapeutic agents. To ensure this, the investigation of the radiolabeling properties and distribution behavior of the BFCAs are performed prior to its bioconjugation. The precursors **B1** and **B2** were examined on these parameters to obtain comparative radiochemical data and study the effects of derivatization on compounds **B3 – B5**.

As discussed earlier, bispidines are excellent ^{64}Cu radionuclide binding agents. The complexation of bispidines with radiocopper occurs at ambient temperature and physiological pH, to form thermodynamically stable and kinetically inert complexes, which makes them an exceptional radiochelator of choice. Another peculiar feature of the complexation is that the radiolabeling is independent of the different kinds of buffer system (acetate, succinate, citrate, and 2-[*N*-morpholino] ethanesulfonic acid (MES) buffer). MES buffer was used for this set of experiments to eliminate any traces of complex formed between $^{64}\text{Cu}(\text{II})$ and the buffer salts in the reaction solution.

The synthesized bispidine derivatives **B1-B5** (Scheme 3.1.5, Table 3.1.2) were radiolabeled using $[^{64}\text{Cu}]\text{CuCl}_2$ in 0.1 M MES/NaOH buffer (pH 5.5) at room temperature. The radioligands $[^{64}\text{Cu}]\text{Cu-B1}$, $[^{64}\text{Cu}]\text{Cu-B2}$, $[^{64}\text{Cu}]\text{Cu-B3}$, $[^{64}\text{Cu}]\text{Cu-B4}$ and $[^{64}\text{Cu}]\text{Cu-B5}$ showed a high radiochemical yield (RCY) of $\geq 99\%$ within 2 minutes of the reaction which is a characteristic feature of $^{64}\text{Cu}(\text{II})$ -bispidine complexes. The aforementioned complexes also demonstrated high resistance towards transchelation and demetalation processes which was confirmed by performing challenge experiments against 10,000 folds excess of competitive ligands such as EDTA ($\log K_{\text{Cu}(\text{II})-\text{L}} = 18.8$) or cyclam ($\log K_{\text{Cu}(\text{II})-\text{L}} = 27.2$) for up to 24 hours.

The radiolabeling and competition reactions of the complexes $[^{64}\text{Cu}]\text{Cu-B1-B5}$ were monitored *via* radio-TLC system (“Method II” section 5.3). A comparative radio-TLC of the $^{64}\text{Cu}(\text{II})$ - bispidine complexes for the challenge experiments is displayed in Figure 3.1.17 (above). The $^{64}\text{Cu}(\text{II})$ -bispidine complexes move on the TLC plate with a retention factor (R_f) of 0.6–0.7. EDTA also complexes any excess or non-specifically bound $^{64}\text{CuCl}_2$ radionuclide to the bispidines. From the TLC analysis it was clear that negligible amount ($< 5\%$) of $[^{64}\text{Cu}]\text{Cu-EDTA}$ ($R_f = 0$) were detected in all the five complex solutions, which supports the characteristic feature of high radiolabeling efficiency, radiochemical

purity and lack of transchelation for $[^{64}\text{Cu}]\text{Cu-B1-B5}$. A reference TLC for $[^{64}\text{Cu}]\text{Cu-EDTA}$ is shown in **Figure 3.1.17** (below).

Figure 3.1.17: Comparative Radio-TLC chromatograms of $[^{64}\text{Cu}]\text{Cu-B1}$ – $[^{64}\text{Cu}]\text{Cu-B5}$ (EDTA challenged) (Above); Radio-TLC of $[^{64}\text{Cu}]\text{Cu-EDTA}$ (standard) (Below).

TLC plotted after 30 min reaction, (TLC system: neutral Alox plates, Mobile phase: 2M $\text{NH}_4\text{OAc}:\text{MeOH}$ (1:1, v/v), (Method II) Retention factor values (R_f).

3.1.3 Conclusions

Working towards the aim of developing bispidine-based radiocopper chelating agents, this chapter dealt with the synthetic optimization of a series of novel C9-functionalized BFCA **B3–B9**. A summary of the synthesized ligands with the respective substituents is presented in **Table 3.1.2**. The derivatives were prepared *via* facile synthetic routes in high chemical yields. The reverse phase preparative HPLC technique was employed to isolate the ligands in high purity suitable for biological applications. The structure elucidation, identification, and characterization of the compounds were done by techniques like NMR, FTIR and mass spectrometry. The synthesized bispidines form hexadentate mononuclear coordination complexes with ^{64}Cu with high thermodynamic stability and kinetic inertness. Like reported bispidine systems, these conjugates also undergo fast complexation (< 5 minutes) with the radiometal at ambient temperature. Copper-64 based experimental radiochemical studies of the conjugates **B1–B5** determined the radiolabeling efficiency and radiochemical purity of the complexes. A detailed description of the bioconjugation of ligands **B4** and **B5** is reported in the next chapter (**Chapter 3.2**).

Table 3.1.2: Summary of hexadentate bispidine derivatives B1–B11 and their substituents. (C9 modified B2–B9 and C1/5 modified B10–B11).

L	R ₁ (C2/4)	R ₂ = R ₃ (N3/7)	R ₄ (C1/5)	X (C9)	Y (C9)
B1					
B2					
B3					
B4					
B5					
B6					
B7					
B8					
B9					
B10					
B11					

3.2 Design and synthesis of Bispidine-based TATE and sd antibody bioconjugates

Bispidines being exceptional radiochelators with excellent radiolabeling efficiency and biocompatibility have been studied in detail over the years, for their development as potential PET imaging agents. Although, very limited data is available in the literature of bioconjugated bispidines for *in vivo* cancer tumor targeting. An *in vivo* biodistribution study of bispidine-bombesin (BBN) peptide bioconjugate was reported by Stephan, Comba and coworkers and showed significant amount of active tumor accumulation in human prostate tumor (PC-3) bearing mice. Recent studies report that bispidines have also been conjugated to amino acids (lysine) [77] and fluorophores (BODIPY) to aid disease targeting and imaging. However, there are no animal experiment data available that can be used to assess the success with such conjugates *in vivo*.

This chapter deals with the synthesis of bispidine bioconjugates designed for ^{64}Cu -based targeted PET imaging of cancerous tumors. Specifically, the isothiocyanate- and dibenzocyclooctyne-terminated bispidines, **B4** and **B5** (**Chapter 3.1**) have been suitably bioconjugated to allow targeting of the somatostatin (SST) and epidermal growth factor (EGF) receptors employing TATE peptide and sdAb respectively. These receptors are highly overexpressed in neuroendocrine cancer tumors and numerous epithelial cancer cells.

3.2.1 Bispidine-based bioconjugates of somatostatin agonist TATE

Somatostatin receptors (SSTR) are known to be overexpressed in a variety of human neuroendocrine tumors. Out of the five receptor subtypes, somatostatin receptor subtype 2 (SSTR-2) is found in several types of malignancies and has become a popular receptor target for modified somatostatin analogues radiolabeled probes for disease detection by nuclear imaging techniques [238, 240, 253]. TATE (Tyr3-Octreotate or Y3-TATE) is a derivative of somatostatin with a long biological half-life and high affinity for SSTR-2 receptors.

The choice of BFCA and radiometal has shown to have a direct effect on the pharmacokinetics of SSTR-targeted imaging agents. Lin *et al.* showed that ^{68}Ga -labeled TATE modified with a NOTA analogue bearing three carboxylates has greater accumulation in SSTR-positive xenograft, and shows superior pharmacokinetics than the analogues with more carboxylates (**Figure 3.2.1**) [239]. Likewise, NOTA analogues also demonstrated superior pharmacokinetics and a higher liver clearance when compared with the DOTA analogue.

Fani *et al.* compared the effect of radionuclides ^{64}Cu and ^{68}Ga on the *in vivo* biodistribution behavior using the radiolabel-modified somatostatin antagonist, NODAGA-LM3 [253]. The ^{64}Cu and ^{68}Ga -labeled agents exhibited similar accumulation in the SSTR-positive xenograft, however, there were significant differences in the clearance from the kidneys and pancreas.

An instructive comparison of commercially available chelators, p-SCN-Bn-NOTA and NODAGA, conjugated to TATE and radiolabeled with ^{64}Cu and ^{68}Ga , respectively, has also been reported. According to the data, the size of linkers and method of conjugation had a significant impact on the performance of the SSTR2-targeted PET imaging agents [237, 253, 254]. For example, directly conjugated BFCA demonstrated superior *in vivo* performance for both ^{64}Cu - as well as ^{68}Ga -labeled analogues compared to chelators conjugated to Y3-TATE with PEG linkages. The choice of radiometal used for the experiments, in this case, was determined to be insignificant.

Figure 3.2.1: Representation of NOTA and NODAGA derivatives of Y3-TATE bioconjugates for PET imaging.

Radiochemistry of **B4**

Prior to the bioconjugation of **B4**, the molar activity and radiochemical purity were assessed to ensure the stability of the radiolabeled complex [^{64}Cu]Cu-**B4**.

For the determination of molar activity, five samples of 0.1, 0.5, 1, 5 and 10 μg of **B4** were dissolved in 100 μL MES buffer. To these samples, [^{64}Cu]CuCl₂ (~100 MBq) was added and the reaction mixture was stirred for 60 minutes at room temperature. The reaction was followed by radio-TLC (“**Method II**” **Section 5.3**) at 5, 30 and 60 minute time points. The complex [^{64}Cu]Cu-**B4** was purified by reverse phase-HPLC system with

a RCY of > 95% (**“Method A” Section 5.2**). The HPLC chromatogram for [^{64}Cu]Cu-**B4** (**Figure 3.2.2**) displayed the elution of radiolabeled complex at a retention time (R_t) of 18.5 minutes. The molar activity of **B4** was calculated to be 181 GBq/ μmol which is optimum for a chelator to be employed for bioimaging purposes.

Figure 3.2.2: Radio-HPLC chromatogram of [^{64}Cu]Cu-B4** reaction mixture and [^{64}Cu]CuCl₂ (inset).** (HPLC system: Gradient 0% to 70% B in 20 min (A = H₂O + 0.1 Vol. % TFA, B = CH₃CN + 0.1 Vol. % TFA) flow rate = 1 mL/min, λ = 220 nm; [^{64}Cu]Cu-**B4** R_t = 18.5 min and [^{64}Cu]CuCl₂ R_t = 6.6 min (**Method A**).

Distribution ratio ($\log D_{o/w}$) of **B4**⁷

The distribution ratio helps in determining the relative hydrophilicity of the compounds, which govern the factors like biodistribution and pharmacokinetic properties of the complex at physiological pH, which is a crucial parameter to be considered prior to associated biological applications. The experiments performed in 1-octanol/water system at pH 7.2, 7.4 and 7.6 gives also an idea about the aqueous solubility of the complex. The obtained negative $\log D$ values for the [^{64}Cu]Cu-**B1** and [^{64}Cu]Cu-**B4** reported in **Table 3.2.1** confirm the hydrophilic behavior of the complexes.

⁷ The radiolabeling experiments were performed by Ms. Karin Landrock, Institute of Radiopharmaceutical Cancer Research at HZDR.

Table 3.2.1: Distribution ratio ($\log D_{o/w}$) values of $[^{64}\text{Cu}]\text{Cu-B1}$ and $[^{64}\text{Cu}]\text{Cu-B4}$ complexes at different pH values.

pH	$\log D_{o/w}$ $[^{64}\text{Cu}]\text{Cu-B1}$	$\log D_{o/w}$ $[^{64}\text{Cu}]\text{Cu-B4}$
7.2	-2.90	-1.25
7.4	-2.88	-1.27
7.6	-2.85	-1.40

3.2.1.1 Synthesis and characterization of B4-TATE bioconjugate

To synthesize the bispidine-based TATE bioconjugate, commercially available Boc-*tert*-butyl-protected TATE peptide with a free terminal amine of the phenylalanine amino acid was coupled with the isothiocyanate-functionalized bispidine **B4**. The two-step synthesis of the conjugate is displayed as **Scheme 3.2.1**.

Scheme 3.2.1: Scheme representing two-step synthesis of B4-TATE bioconjugate.

Step i) Peptide coupling reaction yielding Boc-*tert*-butyl-protected B4-TATE, ii) TFA deprotection reaction yielding B4-TATE.

The TATE peptide precursor solution was activated by DIPEA prior to the addition of **B4** for conjugation. After the addition of **B4**, the reaction solution was stirred overnight at room temperature. The deprotection of the Boc-TATE-**B4** bioconjugate was performed

in situ. TFA was employed to cleave the *tert*-butoxycarbonyl groups on the amine and the tertiary butyl residues on hydroxyl groups. Tri-isopropylsilane (TIPS) was used as a scavenging agent to provide milder conditions for the deprotection reaction to form **B4-TATE**.

Purification and Characterization of **B4-TATE**

The crude **B4-TATE** was purified by RP-HPLC (**“Method B” Section 5.2**) in 25% yield and analyzed by MALDI-TOF mass spectroscopy. The $[M+H]^+$ ion peak at m/z 1921.4 confirms the identity of the product **B4-TATE** (**Figure 3.2.3**). Interestingly, along with the molecular ion peak for **B4-TATE**, an additional peak at m/z 903.4 (**Figure 3.2.3**) was also observed in the spectrum. This indicated the presence of an unexpected secondary product in the sample isolated after HPLC purification.

Figure 3.2.3: MALDI-TOF spectrum of **B4-TATE** with calculated and experimental m/z values.

It was inferred that the thiourea bond formed in the **B4-TATE** bioconjugate is unstable under acidic conditions. The reason for the instability is the placement of the thiourea bond directly on the α -amine of peptide that leads to Edman's degradation of the product,

resulting in removal of the conjugated amino acid of the peptide (**Scheme 3.2.2**) [240]. Of note, in general, such degradation requires acidic conditions for it to occur.

Scheme 3.2.2: Mechanism of Edman degradation of B4-TATE conjugate leading to the loss of phenylalanine.

B4-TATE in this case also undergoes Edman's degradation reaction, which leads to the loss of phenylalanine amino acid conjugated directly to **B4** and the formation of a thiohydantoin derivative (experimental m/z 988.8; calculated m/z 990.3) of the bispidine. The mechanism of the degradation reaction is displayed in **Scheme 3.2.2**.

The movement of the thiourea away from the α -amino group can increase the stability of the thiourea when conjugated to a peptide. Banks and Paquette demonstrated that the conjugation through the ϵ -amine of a lysine versus the α -amine was more stable after 10 days at 37°C . Therefore, to eliminate the possibility for Edman's degradation to take place, the **B4-TATE** construct was redesigned to avoid positioning of the thiourea bond at the α -amine of the peptide (**Section 3.2.1.2**).

3.2.1.2 Synthesis and characterization of B4-EG-TATE⁸

To avoid the Edman degradation, a short linker in the form of diethylene glycol, was then used to connect TATE and **B4**. The commercial availability of the modified peptide EG-TATE was useful in reducing the number of reaction steps and obtaining the bioconjugate in reasonable yields. The two-step reaction between the commercially available EG-TATE and **B4** was performed under similar conditions as for **B4**-TATE using acetonitrile as solvent. The reaction was facile with a simple RP-HPLC purification of the product **B4**-EG-TATE in 33% yield (**Scheme 3.2.3**).

Scheme 3.2.3: Two-step synthesis of B4-EG-TATE bioconjugate.

Step i) Reaction of isothiocyanate terminated **B4** with glycolated TATE peptide (Yield 37%), ii) TFA deprotection to yield B4-EG-TATE (Yield 33%).

⁸ EG refers to diethylene glycol.

The MALDI-TOF mass spectrometry analysis of the **B4**-EG-TATE confirmed the reaction success. The MALDI-TOF mass spectra displayed $[M+H]^+$ ion peak at m/z 2407.1 (calculated m/z 2407.1) for the Boc-protected **B4**-EG-TATE (first step) (**Annex Figure 7.20**) and m/z 2038.5 (calculated m/z 2036.7) for the deprotected **B4**-EG-TATE (second step) (**Figure 3.2.4**). Compared to the **B4**-TATE conjugates, **B4**-EG-TATE bioconjugate showed no signs of degradation when stored under inert conditions at freezing temperature.

Figure 3.2.4: MALDI-TOF spectra of **B4**-EG-TATE with $[M+H]^+$ at m/z 2038.5 (calculated m/z 2036.7).

Radiolabeling of the $[^{64}\text{Cu}]\text{Cu-B4-EG-TATE}$

B4-EG-TATE conjugate was complexed with $[^{64}\text{Cu}]\text{CuCl}_2$ to determine the radiolabeling efficiency and stability of the $[^{64}\text{Cu}]\text{Cu-B4-EG-TATE}$ complex. For the experiment, 10 μg of the conjugate was dissolved in a solution of 10% DMSO and 185 μL 0.05 M MES buffer (pH 5.6). Roughly 30 MBq of $[^{64}\text{Cu}]\text{CuCl}_2$ were added to this solution and stirred for 20 minutes. The reaction progress was monitored by radio-TLC ("**Method II**" **Section 5.3**) and the radiochemical yield (RCY) was calculated to be ~98% with the R_f of

$[^{64}\text{Cu}]\text{Cu-B4-EG-TATE} = 0.8$ and R_f of $[^{64}\text{Cu}]\text{CuCl}_2 = 0.0$. The radiolabeled conjugate was purified by HPLC (**“Method A” Section 5.2**) with a radiochemical purity of > 95% (**Figure 3.2.5**).

Figure 3.2.5: Radio-HPLC chromatogram of $[^{64}\text{Cu}]\text{Cu-B4-EG-TATE}$ reaction mixture with $R_t = 14.8$ min.

(HPLC system: Gradient 0% to 70% B in 20 min; Mobile phase: A = $\text{H}_2\text{O} + 0.1$ Vol. % TFA, B = $\text{CH}_3\text{CN} + 0.1$ Vol. % TFA; Flow rate = 1 mL/min, $\lambda = 220$ nm; $[^{64}\text{Cu}]\text{CuCl}_2$ $R_t = 6.6$ min (Method A)).

The **B4-EG-TATE** bioconjugate is currently under investigation for the *in vivo* PET analysis, the biodistribution behavior and tumor targeting for the SSTR2 tumor bearing mice.

3.2.2 Bispidine-based EGFR targeting sdAb conjugates **B5-sdAb**⁹

For the development of bispidine-based EGFR targeting conjugates, a generic chemoenzymatic strategy of sortase-A (SrtA)-mediated site-specific labeling of camelid single-domain antibody-fragments (sdAbs) was applied. The key advantages of this system are the simple production of the recombinant enzyme in substantial amounts, its flexibility towards its substrates, and the required presence of only very short peptides in these substrates (the sortag in the target protein and a nucleophilic N-terminal oligoglycine within the biorthogonal probe). The mechanism of the two-step transpeptidation is displayed in **Figure 3.2.6**.

In the first step, the polypeptide backbone of the substrate protein is cleaved by the SrtA at the position where the SrtA recognition motif sortag (Leu-Pro-Glu-Thr-Gly: LPETG) is incorporated. This leads to the formation of an acyl-enzyme intermediate. Subsequently, a new peptide bond is formed between the threonine residue in the sortag and the N-

⁹ Results reported in Section 3.2.2 (synthesis, purification and characterization of B4-sdAb) were performed by Dr. Kristof Zarschler, Helmholtz-Zentrum Dresden – Rossendorf. The resulting sdAb derivative was used for the experiment in the frame of this work.

terminal glycine of a nucleophilic probe (for imaging purposes), containing the bioorthogonal functionality. Besides the conjugated sdAb, the reaction mixture still contains unreacted sdAb and imaging probe, SrtA, as well as acyl-enzyme intermediate and released His-tag that result from the first step of the conjugation reaction. Therefore the purification protocol consisted of (i) immobilized metal affinity chromatography (IMAC) to remove all undesired His-tagged products, (ii) EDTA competition, to prevent occupation of the chelating agents by Ca^{2+} or Ni^{2+} ions, and (iii) size-exclusion chromatography (SEC), to separate the conjugated sdAb from the unreacted imaging probe (**Figure 3.2.6 B**).

In this part of the thesis, the site-specific labeling of the EGFR-specific single domain antibody (sdAb) 7C12 with DBCO-functionalized bispidine **B5** for non-invasive cancer imaging is reported.

To develop this conjugate, the sdAb was first functionalized with an azide residue to render a bioorthogonal reaction platform to the DBCO terminated bispidine, **B5**. Therefore, a short peptide (Gly)₃-Lys-N₃, composed of three glycine residues linked through a lysine residue to a terminal azide was used (**Scheme 3.2.4**).

Figure 3.2.6: General representation of Sortase-A mediated conjugation and purification strategies.

A) SrtA-mediated conjugation is a generic conjugation method. First, SrtA recognizes the C-terminal sortag of the sdAb and cleaves the peptide bond between T and G, with release of the downstream His-tag. Next, a nucleophilic attack of the triglycine-functionalized imaging probe on the acyl-enzyme intermediate leads to the formation of a new peptide bond with the sdAb, resulting in the site-specifically conjugated sdAb.

B) Schematic overview of the purification strategy. After SrtA-mediated conjugation the reaction mixture contains, apart from the product-of-interest (conjugated sdAb), also unreacted sdAb, acyl-enzyme intermediate, imaging probe, SrtA and processed His-tags. Except for the imaging probe, all other undesired products contain a His-tag. This allows to capture all His-tagged products *via* IMAC with Ni-sepharose beads on an FPLC system in a first purification step. Next, the collected flow-through (composed of conjugated sdAb and imaging probe) was incubated with an excess of EDTA to remove Ca^{2+} ions, that were present as a cofactor of SrtA in the reaction buffer, and Ni^{2+} ions, that could leak from the Ni-sepharose beads, and prevent them from occupying the chelating agents. After sample volume reduction by concentration, the final purification step consisted of SEC on an FPLC system to separate the product-of-interest from the excess of nucleophilic imaging probe. Adapted from [232]

Scheme 3.2.4: Overview of the two-step site-specific antibody modification and conjugation using a combination of enzyme-mediated bioconjugation and cycloaddition click reaction.

The first step uses enzymatic bioconjugation with the transpeptidase sortase-A for site-specific incorporation of a functional azide group, whereas the second step involves the azide-alkyne cycloaddition click reaction. (SPAAC: Strain-promoted azide–alkyne cycloadditions)

3.2.2.1 Sortase-A-mediated azide functionalization

The chemoenzymatic functionalization was performed using EGFR-specific single domain antibody 7C12 and (Gly)₃-Lys-N₃ as a substrate. The sdAb 7C12 was produced with a terminal hexahistidine purification tag. As successful sortase-A-mediated conjugation leads to the elimination of the hexahistidine tag, this design allows the removal of the unreacted sdAb as well as of the His₆-tagged enzyme by immobilized metal affinity chromatography. The reaction was optimized by varying the molar ratios of SrtA, sdAb and (Gly)₃-Lys-N₃ (**Figure 3.2.7**). A 4 h reaction at 30°C with a molar ratio SrtA:sdAb:(Gly)₃-Lys-N₃ of 1:1:10 was identified as being ideal. After purification of the reaction mixture by affinity chromatography, the obtained azide-functionalized sdAb was analyzed by MALDI-TOF mass spectroscopy (**Figure 3.2.8**).

The mass spectra of the purified product showed a homogeneous product of a single-conjugated, azide-functionalized sdAb with an $m/z \sim 16.6$ kDa.

Figure 3.2.7: SDS-PAGE analysis of Sortase-A mediated azide-functionalization of the single-domain antibody 7C12.

The molar amount of (Gly)₃-Lys-N₃ was varied (50-500 nmol) while the amount of the sdAb as well as the enzyme was kept constant (50 nmol).

Figure 3.2.8: MALDI-TOF mass spectra of (A) sdAb-Leu-Pro-Glu-Thr-Gly-Gly-His₆ and (B) purified azide-functionalized sdAb-Leu-Pro-Glu-Thr-N₃ (calculated $m/z \sim 16.6$ kDa)

3.2.2.2 Strain-promoted azide-alkyne cycloaddition (SPAAC) [118]

The site-specifically azide-modified sdAb was reacted with the DBCO-containing Bispidine (**B5**) for up to 4 h at 25°C. After several optimization reactions, the molar ratios of modified sdAb:B5 of 1:5 yielded the desired bioconjugate in satisfactory yields (**Figure 3.2.9**). The crude sdAb-Leu-Pro-Glu-Thr-B5 (sdAb-B5) conjugate was purified by size exclusion chromatography and analyzed by MALDI-TOF mass spectroscopy (**Figure 3.2.10**). The mass spectra of the final purified product showed a homogeneous product of a single-conjugated, bispidine-functionalized sdAb (B5-sdAb) with a molecular ion peak at m/z ~17.6 kDa.

Figure 3.2.9: SDS-PAGE analysis of strain-promoted azide-alkyne cycloaddition between DBCO-containing bispidine and azide-functionalized single-domain antibody 7C12.

The molar amount of DBCO-Bispidine was varied (1-10 nmol) while the amount of the sdAb was kept constant (1 nmol) (Standards on left and right side of the gel).

Figure 3.2.10: MALDI-TOF mass spectra of purified, single-conjugated sdAb functionalized with the radiometal chelator B5 (calculated m/z ~ 17.6 kDa).

3.2.2.3 Radiolabeling of sdAb-B5 with $[^{64}\text{Cu}]\text{CuCl}_2$

sdAb-**B5** (1 nmol) was radiolabeled with ^{64}Cu with activities ranging from 0.5 MBq to 10 MBq and the extent of radiometal complexation was assessed by radio-TLC (**Figure 3.2.11**). As expected, the amount of protein radiolabeled correlates with the activity of $[^{64}\text{Cu}]\text{CuCl}_2$ added. The reaction mixture also contains free $[^{64}\text{Cu}]\text{CuCl}_2$ that migrates as $[^{64}\text{Cu}]\text{Cu-EDTA}$ complex at the solvent front which was subsequently purified by spin filtration. The $[^{64}\text{Cu}]\text{Cu-sdAb-B5}$ bioconjugate was obtained with a radiochemical purity of ~99%.

In addition to radio-TLC, ^{64}Cu -labeling of sdAb-Leu-Pro-Glu-Thr-**B5** was also confirmed by SDS-PAGE and subsequent radioluminography (**Figure 3.2.12**). A single predominant protein species corresponding to the monomeric ^{64}Cu -labeled sdAb was easily detectable in the Coomassie stained gels as well as in the autoradiographic images. Free $[^{64}\text{Cu}]\text{CuCl}_2$ was present at the origin of the gel, whereas $[^{64}\text{Cu}]\text{Cu-B5}$ migrated to the gel front.

Figure 3.2.11: Radio-TLC analysis of ^{64}Cu -labeled bispidine-conjugated sdAb [^{64}Cu]Cu-sdAb-B5 After conjugation of the sdAb with the chelating agent, the conjugate was labeled with [^{64}Cu]CuCl₂ and analyzed by radio-TLC. Radiolabeled sdAb-Leu-Pro-Glu-Thr-B5 remains at the origin ($R_f = 0$), whereas [^{64}Cu]Cu-EDTA migrates with the solvent front ($R_f = 1$). [^{64}Cu]Cu-B5 has an R_f of ~ 0.7 . Noteworthy, any excess of [^{64}Cu]CuCl₂ is complexed by the addition of EDTA.

Figure 3.2.12: (A) Discontinuous SDS-PAGE and (B) Autoradiography analysis of [^{64}Cu]Cu-sdAb-B5. After radiolabeling, aliquots of each reaction were separated using a 15% SDS-polyacrylamide gel, and prior to staining with Coomassie staining (A), an electronic autoradiographic image of the gel was recorded using imaging plates/radioluminography laser scanner (B).

3.2.2.4 Cell binding studies

In order to investigate the targeting ability of the functionalized sdAb to EGFR, binding to the human epithelial cell line A431 originating from an epidermoid carcinoma of the skin was examined. These squamous carcinoma cells express approximately 2×10^6 EGFR molecules per cell, which represents a high expression level [255, 256]. In a saturation binding assay, where the amount of radioligand required to saturate these receptors is measured, the equilibrium dissociation constant K_d and the total number of receptors expressed on the cell surface B_{max} can be determined (**Figure 3.2.13**).

Figure 3.2.13: Analysis of binding affinity bispidine-functionalized sdAbs to human EGFR-presenting cells.

Total binding was measured in the absence of and nonspecific binding in the presence of 1 mM unlabeled sdAb. Specific binding was calculated as the difference between total and nonspecific binding.

The binding studies on the A431 cells showed promising results and directed the conjugate towards the *in vivo* analysis and biodistribution studies. The active targeting and pharmacokinetics studies of sdAb-**B5** bioconjugates for the EGFR-specific A431 tumor bearing xenografts are not a part of this work.

3.2.3 Conclusions

In this chapter, the synthesis of two novel bispidine-based imaging agents has been reported, which will be employed for the active targeting of tumors. The receptor-specific PET imaging agents can potentially be a useful tool for early diagnosis of malignant tumors.

The first conjugate **B4-EG-TATE** involves the coupling of EG-TATE to **B4** in a straightforward manner leading to a stable bioconjugate in high yields with efficient radiolabeling tendency, which is a crucial feature for determining the prospect of short-lived contrast agents. The isothiocyanate-terminated radiochelator **B4** was coupled to a diethylene glycol-modified TATE peptide *via* stable thiourea bond. The radiolabeling experiments of the conjugate displayed satisfactory results and therefore, [⁶⁴Cu]**Cu-B4-EG-TATE** was directed for *in vivo* analysis targeting the SSTR-2 cancer tumor receptors.

The second conjugate, **sdAb-B5** was developed by taking advantage of the chemoenzymatic Sortase-A-mediated labeling of the sdAb followed by the strain promoted alkyne-azide coupling (SPAAC) with **B5**. The reaction resulted in a homogeneous, site-specifically single-conjugated sdAb with an uncompromised functionality. The introduction of the azide handle at the C-terminal end of the sdAb makes it complementary to the DBCO-functionalized **B5**. The cell binding studies of **sdAb-B5** were also performed to ensure their tumor cell targeting efficiency.

The strategies represent an important contribution to expand the application of the two bispidine derivatives conjugated to different targeting vectors (TATE and sdAb) for different cancer types. This successful synthesis expands the use of bispidines as BFCAs for future preclinical studies.

3.3 Synthesis and characterization of amine-terminated ultrasmall Si NPs¹⁰

The research community assumes that the Si NPs will leave great potential for application in the future due to properties such as excellent physico-chemical behavior, extraordinary biocompatibility and low cytotoxicity. The surface-termination of the Si NPs by various functionalities like amines and esters has proved to be a beneficial strategy for developing multifunctional NPs with accessible surface groups [28, 152, 257, 258] (**Chapter 2.2.1**).

This chapter focuses on the synthesis, purification, and characterization of ultrasmall (~4 nm) amine-terminated Si NPs with large surface-to-volume ratio, and various functionalization strategies for further decoration with modular ligands.

3.3.1 Synthesis of amine-terminated Si NPs

The method used in this thesis for the hydrothermal synthesis of the ultrasmall NPs is a slight modification of the protocol reported in the literature [259]. Specifically, an aqueous solution of (3-aminopropyl)-trimethoxysilane and trisodium citrate was heated under microwave conditions for about 5 hours at 160 °C. This lead to the production of blue emitting spherical Si NP in with reasonable photoluminescent quantum yields (PLQY) (~15%). The synthesis of the Si NPs is presented in **Figure 3.3.1** [260]. Although the mechanism for this reaction is not yet fully understood, it is hypothesized that it follows Ostwald ripening method [259]. A large scale synthesis (400 mg Si NPs batch) was purified by dialysis against millipore water. Si NPs were stored protected from light at 4 °C and were stable up to 2 months with almost no aggregation and minor lowering of photoluminescence.

¹⁰ The synthesis and characterization of the Si NPs was performed in collaboration with Prof. Luisa De Cola at University of Strasbourg with the assistance of Dr. Angélique Faramus and Mr. John Ddungu.

Figure 3.3.1: Synthesis of amine-terminated Si NPs by hydrothermal method bottom-up strategy.
Adapted from [259]

3.3.2 Characterization of Si NPs

Transmission electron microscopy (TEM)

TEM images displayed monodispersed Si NPs in the size range of 2 to 5 nm (**Figure 3.3.2**). According to the size distribution histogram, the average size of Si NP is 4.3 ± 1.3 nm. High resolution TEM images of a single Si NP shows a crystal lattice of silicon with a distance of 2.0 Å between fringes, which is a characteristic lattice spacing of the (220) planes of crystalline silicon [261].

Figure 3.3.2: TEM and HR-TEM of Si NPs (left); size distribution graph of Si NPs with an average size of 4.3 ± 1.3 nm (right) [262]

UV-Visible spectroscopy

The water dispersed Si NPs displayed a stable absorbance and photoluminescence (PL) mainly in the blue region of the electromagnetic spectrum with the maximum absorbance ($\lambda_{\text{abs,max}}$) at 353 nm (**Figure 3.3.3**).

Figure 3.3.3: Absorption spectrum of water dispersed Si NPs ($\lambda_{\text{abs}} = 353 \text{ nm}$).

The emission maxima (λ_{ems}) for Si NP, when excited at $\lambda = 355 \text{ nm}$ is centered at 446 nm indicating that the Si NPs possess good optical properties with clearly resolved absorption peaks and symmetrical photoluminescence peaks. A slight red shift is observed when the spectra were recorded at higher excitation wavelengths from 290 to 390 nm (**Figure 3.3.4**) [169, 263]. In comparison to previously reported red-emitting Si NPs ($\lambda_{\text{max}} = 660 \text{ nm}$), these blue emitting Si NPs exhibit distinctly shorter λ_{max} ($\sim 446 \text{ nm}$) due to quantum-size effect. Also, the fluorescence of the Si NPs (PLQY: 2–15%) is attributable to both quantum-size and ligand-related effects, upon excitation.

Figure 3.3.4: Emission spectra of water dispersed Si NPs at different excitation wavelengths in nm (exc).

Fourier-transform infrared spectroscopy (FTIR)

The FTIR spectrum (**Figure 3.3.5**) was in agreement with the vibration bands reported in the literature for amine-terminated Si NPs [54, 143, 155, 264]. The spectral region from 3342 cm^{-1} to 2965 cm^{-1} corresponds to the C-H stretching from the propyl groups and the N-H stretching from the propyl amine. The peaks observed from 1650 cm^{-1} to 1550 cm^{-1} refer to the bending vibrations of the primary amine. The peaks in the region 1400 cm^{-1} to 1270 cm^{-1} relate to the vibrational modes of Si-CH₂ bond.

The fingerprint region displayed characteristic bending vibrations of the covalent bonds. Band observed at 1568 cm^{-1} is due to N-H bending vibrations from the amine. The vibration band at 1390 cm^{-1} refers to the Si-C bond, which confirms that the carbon is linked covalently to the silicon atom. A characteristic band from extending from 1100 to 1000 cm^{-1} confirms the presence of Si-O bonds, which are also present on the surface of NPs along with Si-C bonds. Weak bands near 1100 cm^{-1} and 1000 cm^{-1} are characteristic for silicon oxide vibration modes.

Figure 3.3.5: FTIR spectra of amine-terminated Si NPs with highlighted characteristic bands.

3.3.3 Surface characterization of Si NPs

The number of available amine groups per Si NP was estimated by thermogravimetric analysis (TGA) (**Annex Figure 7.23**) and Kaiser (Ninhydrin) test which follows the Beer-Lambert's Law. These experiments revealed that approximately 162 freely accessible

amine groups (from TGA measurements) are present on the surface of one Si NPs that corresponds to 6.8 μmol of amines per mg Si NP (from Kaiser test). (**Experimental section 5.4.3**) [265].

For the NPs to be biocompatible and show a uniform biodistribution and favorable pharmacokinetics (ADME), the surface charge of the particle plays a major role. Neutrally charged NPs are a preferred choice for biological studies as they show minimum cytotoxicity and rapid elimination from the body with reduced accumulation in body organs [266]. Water-dispersed amine-terminated Si NPs can be assumed to exist as protonated ammonium species. In the presence of citric acid, they have a zeta potential of $(-1.1 \pm 3.9 \text{ mV})$ at room temperature and physiological pH 7.4. The results suggest that despite being amine-terminated, the NPs possess a zwitterionic nature at physiological pH. This can be explained by the presence of a citrate shell around the surface of the Si NPs. The cationic ammonium groups of Si NPs tend to get neutralized by the negatively charged carboxylates of the citrate present on the surface of the NPs. This explanation was confirmed by NMR spectroscopy (**Annex Figure 7.21**). Temperature-dependent ^1H NMR spectroscopy (range: 10–45 $^{\circ}\text{C}$) also showed no significant difference in the chemical shifts confirming the temperature independence on the binding of citrate to the NPs.

3.3.4 Conclusions

The hydrothermal method of synthesis leads to the successful production of ultrasmall amine-terminated, water-dispersible Si NPs. Using this method, the NPs were obtained in high yields and with a good reproducibility. The accessible amino groups on the surface will allow for further functionalization with different moieties. The hydrophilicity of the Si NPs ensures the biocompatibility for *in vivo* studies and the ultrasmall size and neutral charge possibly facilitate the rapid renal clearance from the body. Altogether, the physico-chemical and surface properties of Si NPs suggest that they can be used as platform for the development of multifunctional imaging agents for further investigation *in vivo* mouse.

3.4 Synthesis, characterization, and biodistribution studies of Si NP-based mono-labeled PET and optical imaging agents¹¹

The Si NPs have been investigated as molecular imaging probes. The literature review displays preliminary cytotoxicity and cellular studies of Si NPs that show their potential as biomedical agents. Yet, the development of a suitable Si NP conjugate for multimodal imaging *in vivo* has not been achieved. This section shows the efforts towards the construction of Si NP-based mono-labeled imaging probes to overcome the existing limitations related to their biodistribution and small-animal studies.

The good dispersibility in water and ease of functionalization of the Si NPs allowed the decoration of the NPs with ligands for either PET or OI. A summary of the mono-labeled PET/optical imaging probes synthesized in this work is presented in **Scheme 3.4.1**.

After the successful synthesis of the mono-labeled Si NPs conjugates, radiolabeling studies were performed to quantify the amount of BFCAs attached to the surface of Si NPs. Further, the conjugates were assessed for their cytotoxicity and then directed towards the preclinical studies. These included the *in vivo* biodistribution studies, followed by the *ex vivo* autoradiography studies on healthy rats as well as A431 tumor bearing mice to determine the fate of the nanoobjects.

Similarly, the OI probes were prepared by modifying the Si NPs with IR 800 fluorophore. The NP conjugates were examined for their biocompatibility and toxicity, and further subjected to *in vivo* and *ex vivo* evaluation.

¹¹ A few sections of this chapter are an excerpt of the published article:

Licciardello, N., Hunoldt S; Bergmann, R.; **Singh, G.**; Mamat, M.; Faramus, A.; Ddungu, J.L.Z.; Silvestrini, S.; Maggini, M.; Cola, L. De; Stephan, H. **Biodistribution studies of ultrasmall silicon nanoparticles and carbon dots in experimental rats and tumor mice**
DOI: 10.1039/c8nr01063c.

Scheme 3.4.1: Scheme of the synthesis of mono-labeled Si NP-based imaging agents.

3.4.1 Si NP-NOTA derived PET imaging agent¹²

3.4.1.1 Synthesis and radiolabeling studies of Si NP-NOTA

The Si NPs were reacted with the NOTA-Bn-SCN (0.1 μmol / mg Si NPs) in aqueous medium to produce Si NP-NOTA conjugates. The synthetic scheme for the production of the Si NP-NOTA conjugate is shown as **Scheme 3.4.2**. The Si NP-NOTA conjugates were purified by dialysis followed by SEC with 0.9% NaCl solution (w/v) as eluent to remove the retaining impurities. Small (~ 100 μL) fractions of the Si NP-NOTA sample were collected from SEC.

Scheme 3.4.2: Reaction of amine-terminated Si NPs with NOTA-Bn-SCN to produce Si NP-NOTA.

Si NP-NOTA conjugates were labeled with [⁶⁴Cu]CuCl₂ in MES/NaOH buffer (pH 5.2) at room temperature for 30 minutes. Radio-iTLC (radio instant thin layer chromatography) confirmed complete complexation of [⁶⁴Cu]Cu-Si NP-NOTA with an $R_f = 0$ and no trace of free ⁶⁴Cu(II) ($R_f = 1$) (**“Method III” Section 5.3**). The stability of the ⁶⁴Cu-labeled NPs was tested in challenge experiments using EDTA (**Figure 3.4.1**). Radiolabeling studies exhibited that Si NP-NOTA readily complexes ⁶⁴Cu(II), forming a thermodynamically stable and kinetically inert [⁶⁴Cu]Cu-Si NP-NOTA under mild conditions conserving its radiolabeling efficiency even after NP conjugation.

¹² The synthesis, characterization and PET studies of Si NP-NOTA were performed by Dr. Nadia Licciardello and Sebastian Hunoldt and published in [262] Licciardello, N., et al., Biodistribution studies of ultrasmall silicon nanoparticles and carbon dots in experimental rats and tumor mice. *Nanoscale*. **10** (2018) 9880-9891.

Figure 3.4.1: Radiochemical purity of purified ^{64}Cu -labeled Si NP-NOTA conjugate.

EDTA incubated and SEC purified ^{64}Cu Cu-Si NP-NOTA; Radio-TLC system: iTLC-SG in $\text{NH}_4\text{OAc}:\text{MeOH}$ 1:1 (v/v); ^{64}Cu Cu-Si NP-NOTA $R_f = 0$, ^{64}Cu Cu-EDTA $R_f = 1$ (Method III).

Quantification of the chelator NOTA on Si NPs

Radiometric titrations of Si NP-NOTA was performed to determine the amount of NOTA conjugated to the Si NPs. Radionuclides, such as ^{57}Co and ^{64}Cu , have been employed in the past to determine the number of radiometal chelating ligands attached to the biomolecules or NPs [267-269]. The major advantage of this procedure is the extremely low amount of sample (μg) required for detection. The radiometric titration method exploits the fact that the Si NP scaffold itself does not bind Cu(II) ions. To confirm this, an excess of EDTA was added after complex formation.

For this experiment, increasing amounts of a mixture of radioactive (^{64}Cu) CuCl_2 and non-radioactive (CuSO_4) Cu(II) was added to five different batches of Si NP-NOTA. The amount of the conjugate was kept the same for all the five samples. Subsequently, an excess of EDTA was added to remove any radionuclide non-specifically bound to the Si NPs. The amount of $^{64}\text{Cu(II)}$ bound to the Si NP-NOTA complex relates to the number of bound chelator per chelate (**Section 5.5**). The complexation between Cu(II) and NOTA follows a stoichiometry of 1:1 according to which only one copper can complex to each NOTA chelator. The process of chelation was followed *via* radio-TLC analysis (**“Method III” Section 5.3**). **Figure 3.4.2** illustrates the plot of the percentage of $^{64}\text{Cu(II)}$ bound against the total amount of nonradioactive Cu(II) added. 0.09 μmol NOTA chelator was calculated bound per mg Si NP-NOTA.

Figure 3.4.2: Radiometric titration curve for $^{64}\text{Cu(II)}$ and Cu(II) labeled Si NP-NOTA conjugate. (Experimental value: $\sim 0.09 \mu\text{mol}$ NOTA/mg Si NP-NOTA)

3.4.1.2 *In vitro* cytotoxicity and cell viability studies of Si NP-NOTA¹³

After the characterization of the Si NP-NOTA, it was important to test its biocompatibility. Previous studies demonstrated that the amine-terminated Si NPs prepared by the hydrothermal method have very low cytotoxicity in human epithelial cervical cancer cells (HeLa) even after 48 hours of exposure [259]. In addition, it is also reported that the toxicity of amine-terminated Si NPs increases in the presence of serum [157] and this was ascribed to the fact that serum proteins facilitate the cellular uptake of NPs and thus cause higher toxicity [270-272]. To determine the affect of Si NP-NOTA conjugates on cellular viability, the human embryonic kidney (HEK 293) cell line was chosen. The literature suggests that the preferred mode of elimination of ultrasmall NPs is renal and therefore, it was important to ensure the lack of any nephrotoxic effects of these nanoconjugates on the kidney cells.

For the cell viability and cytotoxicity measurements, the HEK 293 cells were incubated with varying concentrations of Si NP-NOTA and analyzed for up to 72 hours. The viability and cytotoxicity of the cells were plotted against the concentration of the Si NP-NOTA conjugates. From both the assays, it was inferred by visual detections (490 nm) that the modified Si NPs do not induce any substantial changes in the cellular metabolism of the experimental cell culture.

¹³ Cell experiments of Si NPs were performed by Mrs. Utta Herzog assisted by Ms. Neena Patel, Institute of Radiopharmaceutical Cancer Research, HZDR.

The CyQ assay demonstrated > 90% viability and < 5% cytotoxicity for 1, 10, 50 and 100 $\mu\text{g/mL}$ Si NP-NOTA concentrations after 24, 48 and 72 hours, respectively (**Figure 3.4.3**).

The MTS (3-(4,5-dimethylthiazol-2-yl)-5-(3-carboxymethoxyphenyl)-2-(4-sulfophenyl)-2H-tetrazolium) and LDH (lactate dehydrogenase) calorimetric assays also illustrated no negative influence of the nanoconjugates on the metabolism and viability (> 95%) of the system over a period of 72 hours (**Figure 3.4.4**). The intact cell membranes of the cells after 72 hours of incubation assured the negligible amount of nanotoxicity (< 10%) caused by the Si NP-NOTA.

Figure 3.4.3: *In vitro* cell viability and membrane integration studies of Si NP-NOTA by CyQuant assay in human embryonic kidney (HEK293) cell lines after incubation up to 72 h.

Figure 3.4.4: *In vitro* toxicity evaluation of Si NP-NOTA conjugate by MTS and LDH assay after 48 hours in HEK293 cells.

Cell viability (mitochondrial activity) of Si NP-NOTA in HEK 293 cells at different concentration. Triton-X was used as positive control. The percentage of cell viability is calculated relative to untreated cells (triplicate), which served as a negative control. The membrane integrity is measured as lactate dehydrogenase (LDH) release into the media relating to necrosis at different time points (data shown after 48 h incubation). BSA served as a negative control and a triplicate set of wells were treated with supplied lysis buffer for maximum LDH release and regarded as a positive control. A triplicate of wells with cells treated with conjugates with no MTT added was used as blank for any interference of the fluorescence. The percentage of cytotoxicity and LDH release is relative to that of the controls.

3.4.1. Biodistribution of [⁶⁴Cu]Cu-Si NP-NOTA by small animal PET

Si NP-NOTA showed rapid clearance from the body with 84.5% ID¹⁴ in the urine 5 minutes post injection (**Figure 3.4.5**). The rapidly transferred activity in the urine confirms the renal filtration of the conjugates in the *in vivo* system. The particles thus confer to be neutral, nanosized and biocompatible with favorable pharmacokinetics.[262]

¹⁴ Accumulated activity amount (% ID = % injected dose).

Standardized uptake value (SUV) = activity concentration in tissue [Bq/g] × body weight [g] / injected dose [Bq] at a certain time point p.i

Figure 3.4.5: *In vivo* PET imaging of [^{64}Cu]Cu-Si NP-NOTA at different time points.

Maximum intensity projections obtained from a PET study after a single intravenous injection of [^{64}Cu]Cu-Si NP-NOTA ($n = 2$) in A431 tumor-bearing mice after 1 min (A, E), 5 min (B, F), 30 min (C, G), and 60 min (D, H) after injection. The images A–D were scaled to the maximum activity and the images E–H were rescaled to visualize the tumor (bl = bladder, he = heart, in = intestine, ki = kidney, li = liver, tu = tumor).

The radiolabeled particles were also observed to rapidly diffuse in the tumor with visualization under optimized imaging conditions. The maximum tumor uptake is detected within 5 min of the intravenous administration. However, the activity was quickly washed out from the tumor, as is expected for such small sized particles, with the exception of NPs equipped with tumor-specific moieties, which resided longer in the tumors. This infers the passive targeting that occurs for the first 5–10 minutes. However, the size of the particles is too small (< 30 nm) to observe any significant EPR effect and therefore, the signal is again under the limit of detection. [225]

3.4.2 Synthesis and characterization of Si NP-Bispidine (Si NP-B4)

Bispidines being excellent ^{64}Cu chelators were used as a BFCA for the development of Si NP-based PET imaging agent. A general scheme for the synthesis of Si NP-B4 is represented as **Scheme 3.4.3**. For the preparation of Si NP-B4, a series of solvent mixtures were tested in order to optimize the reaction conditions. In contrast to NOTA, B4 possesses a lower solubility in aqueous medium, leading to a decrease in the extent of conjugation due to aggregate formation. Table 3.4.1 shows the different tested conditions for the conjugation reaction between B4 ($0.1 \mu\text{mol} / \text{mg}$ Si NPs) and Si NPs. Initial trial reactions with solvent mixtures (water/ethanol, water/ACN) inferred that the

reaction works best when heated up to 35 °C and run for 12 hours. Although, in some cases, the reaction turned turbid immediately after the addition of B4 ligand, which later (1-2 hours) forms a clear solution. With solvents like DMSO, a large amount of aggregation was observed in the reaction mixture. Reactions between B4 (0.1 μmol / mg Si NPs) and Si NPs in H₂O/ACN (9:1) v/v yielded Si NP-B4 when stirred at 35 °C and run for 12 hours.

Scheme 3.4.3: Reaction of amine-terminated Si NPs with B4 to produce Si NP-B4

Table 3.4.1: Optimization of the synthesis of Si NP-B4 at pH 7–7.5.

Solvents used	Observation
Water + Ethanol (9:1 v/v)	Clear
Water + 0.1% DMSO	Aggregation
Water + ACN (9:1)	Clear
Phosphate buffer + Ethanol (9:1)	Turbid
Phosphate buffer + ACN (9:1)	Turbid

For purification of the crude Si NP-B4, SEC was repeated twice to yield the product. Of note, dialysis techniques did not work well for the purification of Si NP-B4 conjugates. The reaction yield of Si NP-B4 was lower compared to other more hydrophilic conjugates (Si NP-NOTA) due to the loss of NPs in form of aggregates.

The process of radiolabeling was monitored by radio-iTLC SG using 2M NH₄OAc:MeOH 1:1(v/v) solvent system (**“Method III” Section 5.3**). The ⁶⁴Cu(II)-labeled Si NP-B4 was detected at the origin with $R_f = 0$ while the free ⁶⁴Cu(II) moves to the front with an $R_f = 1$ (**Annex Figure 7.22**).

Radiometric titration of Si NP-B4

Radiometric titration was performed to calculate the amount of bispidine conjugated per milligram of Si NP. Bispidines also forms mononuclear Cu(II) complexes, therefore, the

number of moles of nonradioactive copper can be quantified to calculate the conjugated **B4** to the Si NPs. The radiolabeling reaction was performed as mentioned in **Section 3.4.1.1** for Si NP-NOTA conjugates. The amount of conjugated **B4** to the Si NPs derived from **Figure 3.4.6** was calculated to be $\sim 0.052 \mu\text{mol/mg}$ Si NPs. Compared to Si NP-NOTA conjugate ($0.09 \mu\text{mol/mg}$ Si NPs) the amount of bispidine present per milligram of Si NP is much less, which clearly was due to the lower hydrophilicity of **B4**. The solubility differences lead to lower radiolabeling efficiency and radiochemical purity of the Si NP-**B4** conjugates. Due to these reasons, the experiments on the Si NP-**B4** conjugates were discontinued and no further studies were performed.

Figure 3.4.6: Radiometric titration curve for $^{64}\text{Cu(II)}$ and Cu(II) labeled Si NP-**B4** conjugate. Experimental value: $\sim 0.052 \mu\text{mol B4/mg}$ Si NP-**B4** determined by radio-TLC analysis (Method III)

3.4.3 Synthesis, characterization, *in vitro* and *in vivo* evaluation of Si NP-IR800 as an optical imaging probe

The Si NPs were coupled to IR 800 infrared dye to develop them as potential optical imaging agents. Presence of an optical label in the form of IR 800 dye should allow the study of *in vivo* biodistribution and pharmacokinetic behavior of the Si NP-IR800 *via* fluorescence imaging. Keeping in mind our long- term goal of developing such probes for clinical applications, IR 800 dye was chosen as it has successfully passed all animal toxicity studies and recently been approved by FDA. Its characteristic emission lies in the red region of the visible spectrum ($\lambda_{\text{em}} = 789 \text{ nm}$ when excited at $\lambda = 774 \text{ nm}$, in water) that circumvents the possibility of the overlap with the auto-fluorescence caused by the living cells.

3.4.3.1 Synthesis of Si NP-IR800

The Si NPs were reacted with the NHS-activated IR800 dye in aqueous medium to produce Si NP-IR800 conjugates (0.5 μmol NHS-IR800 per mg Si NPs). The synthetic scheme for the production of the Si NP-IR800 conjugate is shown as **Scheme 3.4.4**. Variations in reaction conditions like the reaction time and temperature showed significant differences in the efficiency of the reactions. **Table 3.4.2** shows the different reaction conditions explored along with the solvent system.

Scheme 3.4.4: Reaction of amine-terminated Si NPs with NHS-IR 800 to produce Si NP-IR800.

The reaction of Si NPs and NHS-IR800 showed better results when stirred for longer hours (~12 hours) as analyzed by UV-Vis spectroscopy. Because water and the phosphate buffer delivered similar results, it was more feasible to select water as the solvent to avoid the desalting of the sample while purification (**Table 3.4.2**).

Table 3.4.2: Optimization of the synthesis of Si NP-IR800 reaction.

Solvent	Temperature (°C)	Time (h)	pH
Water	25	6	7-7.5
Water	25	12	7-7.5
Water	35	12	7-7.5
Phosphate buffer	25	12	7.4
Phosphate buffer	35	12	7.4

3.4.3.2 Purification and characterization of Si NP-IR800

The Si NP-IR800 conjugates were purified by dialysis followed by SEC with 0.9% NaCl solution (w/v) as eluent to remove the retaining impurities. Small ($\sim 100\ \mu\text{L}$) fractions of the Si NP-IR800 sample were collected from SEC and analyzed by RP-TLC plates using methanol as eluent. The TLC was read under the IR short channel (710 – 730 nm) and IR long channel (810 – 840 nm) of the scanner (300-900 nm) that yielded similar results. A representation of the TLC chromatogram of an impure sample (containing Si NP-IR800 and IR800 dye) and a pure sample (Si NP-IR800) is represented as **Figure 3.4.7**. Optimization of the purification strategy leads to a faster analysis with reproducible uniform Si NP-NOTA. Fluorescence TLC and UV-Vis measurements effectively detected the purity of the conjugates, demanding only nanograms of the sample.

Figure 3.4.7: Comparative Fluorescence TLC chromatograms to detect the purity of Si NP-IR800.

A: TLC plate of crude Si NP-IR800 sample (left) and the chromatogram showing Si NP-IR800 ($R_f = 0$) and free IR 800 dye ($R_f = 1$) (right). B: TLC plate after dialysis and SEC purified Si NP-IR800 sample (left) and the chromatogram showing Si NP-IR800 ($R_f = 0$). TLC system RP-TLC plate, Solvent system: Methanol.

Absorption/Emission properties of Si NP-IR800

After its conjugation to the Si NPs, a huge hypsochromic shift (λ_{abs} 775 nm to 612 nm) is observed for the Si NP-IR800. The probable reasons for this massive shift in the absorbance maxima could be ascribed to the disruption in the extended conjugation of the conjugated IR800 dye (*trans* to *cis* configuration). The profile broadening observed in this case could be due to the increased heterogeneity in the sample induced by chemical interaction among the dye molecules and with the Si NPs.

Anisotropy studies, NMR and Raman spectroscopy could be employed to deduce the exact reason for this behavior which is not a part of this work [273-276]. Nonetheless, the Si NP-IR800 conjugates remain in the NIR region (600–900 nm) and thus, can be applied for the *in vitro* and *in vivo* trials as potential imaging agents.

A comparative normalized absorption and emission spectra of Si NP-IR800 and IR 800 fluorophore are displayed as **Figure 3.4.8**. The emission spectrum of the Si NP-IR800 exhibited a smaller blue shift from λ_{em} 792 nm (IR800) to 754 nm with a typical shoulder peak which is a characteristic feature of cyanine dyes.

Figure 3.4.8: A comparative absorption (A) and emission spectra (B) of Si NP-IR800 and NHS-IR 800 dye in water.

The absorption and emission maxima of Si NPs, IR 800 dye and the Si NP-IR800 conjugate is reported in **Table 3.4.3**. The photoluminescent quantum yield (PLQY) for Si NPs and Si NP-IR800 was calculated from the respective normalized absorption-emission spectra. NHS-IR 800 dye in water (PLQY = 14.2%) was used as a reference for calculating the PLQY of Si NP-IR800 conjugate.[277]

Table 3.4.3 Photophysical characterization of Si NP-IR800 vs Si NPs and IR 800 dye.

	IR800 dye (nm)	Si NPs (nm)	Si NP-IR800 (nm)
Absorbance (λ_{abs})	775	355	612
Absorbance (shoulder)	700	-	470
Fluorescence (λ_{ems})	792	456	754
PLQY	14.2%	15%	22.5%

Size and surface charge determination

Due to the extremely small size, metalloid behavior and the organic functionalization on the Si NPs, techniques like DLS to determine the size as hydrodynamic diameter are fairly challenging and do not work properly. The physicochemical properties, the surface charge studies and the functionalization, on the other hand, provide adequate details about the behavior and characterization of the conjugates.

Surface charge determination

The zeta potential (ζ) measurements were conducted to measure the surface charge of the Si NP-IR800 conjugates. A value of -5.4 ± 4.1 mV (water, pH 7) deduced from the experiments for IR800 conjugates was comparable to that of nonconjugated Si NPs (-1.1 ± 3.9 mV). This infers to fact that the IR800 functionalization of the Si NPs does not remarkably influence the surface properties for the NPs. This was done to preserve the ultrasmall and zwitterionic behavior of the conjugates, which is crucial for its biomedical applications. It was achieved by limiting the surface functionalization of the NPs by employing a small amount of dye (0.05 $\mu\text{mol/mg}$ Si NPs), which makes it a suitable optical imaging yet conserving its inherent properties.

Quantification of IR800 dye conjugated to Si NPs

The amount of dye conjugated to the Si NPs was determined by UV spectroscopy. For this, the absorbance of parent IR 800 dye was measured at different concentrations to obtain a calibration curve, which was then used to obtain the dye concentration in the Si NP-IR800 particles (**Figure 3.4.9**). A weighted sample of Si NP-IR800 was checked for the absorbance and plotted on the same graph to know the concentration of the dye attached to the Si NPs.

The experimental value was calculated as ~ 0.013 μmol IR 800 dye per mg Si NPs.

Figure 3.4. 9: Quantification of IR 800 dye present on Si NP-IR800 by UV-Vis spectroscopy (Experimental value ~ 0.013 μmol IR 800 dye per mg Si NPs).

3.4.3.3 Cell studies of Si NP-IR800

As discussed earlier, for a nanoobject to be developed for pre-clinical and clinical imaging trials, it is of utmost importance to understand its cellular behavior and biocompatibility [210, 266, 278]. The viability of the human embryonic kidney (HEK 293) cell line was determined for Si NP-IR800 over a time frame of 72 hours (**Section 3.4.1.2**). The assessment was performed using MTS calorimetric assay, LDH assay and the CyQUANT® Direct Cell Proliferation Assay. The data obtained from the CyQ assay for Si NP-IR800 presented $> 95\%$ viability of the cell lines for all the four tested concentrations over a period of 72 hours (**Figure 3.4.10**) which is similar to data obtained for Si NP-NOTA conjugates. Interestingly, negligible amount of toxicity was recorded even at the highest concentration of $100 \mu\text{g/mL}$ of Si NP-IR800, which inferred that the Si NP functionalization did not induce any substantial changes in the cellular metabolism of the *in vitro* system.

Figure 3.4.10: *In vitro* cell studies of Si NP-IR800 by CyQuant assay in HEK293 cell lines.

The MTS assay also depicted viability and proliferation values comparable to the CyQ assay. The LDH assay performed to test the necrosis and cytotoxicity, revealed that even after 48 hours of incubation of 1, 10, 50 and 100 µg/ml nanoconjugates with the cell lines, the cellular membrane remains integrated, and the release of LDH is in the range of the negative control with a value < 5% (**Figure 3.4.11**).

Figure 3.4.11: *In vitro* toxicity evaluation of Si NP-IR800 by MTS and LDH assay after 48 h.

3.4.3.4 Determination of *in vivo* behavior of Si NP-IR800 in mice¹⁵

The successful *in vitro* analysis of the Si NP-IR800 conjugates with convincing biocompatibility instigated us to evaluate the *in vivo* potential of the fluorescent NPs. The fluorophore conjugated to the Si NPs allows the imaging of the mice with acceptable resolution. For the biodistribution experiments, ~ 100 µg (200 µL) of the Si NP-IR800 sample was administered intravenously to four healthy nu/nu murine mice. The mice were analyzed at different time points (1 minute, 10 minutes and 2 hours) by whole body fluorescence scanners to detect the fluorescence intensity in each organ. Exemplary images of the mice after the intravenous injections at different time points are displayed as **Figure 3.4.12**.

The 2 h time frame is best suited for these ultrasmall Si NPs as they are anticipated to undergo a fast renal elimination within this duration. The optical image recorded immediately (1 minute) after the administration, displayed a uniform distribution of the Si NP-IR800, all over the body of the mice, *via* the blood circulation. Within 10 minutes of injection, the particles started to pass through the kidneys, reaching the bladder and rapidly eliminated *via* urine.

Figure 3.4.12: *In vivo* optical imaging of Si NP-IR800 at different time points.

Optical imaging scans of after 1 min, 10 min and 2 h of intravenous injection in healthy NMR nu/nu mice. The scans show a rapid clearance of the particles *via* kidneys into the urine after uniform distribution of the Si NP-IR800 throughout the body (R: Reference, injected sample).

¹⁵ Biodistribution studies were conducted by Dr. Ralf Bergmann and Ms. Regina Herrlich in the group of Prof. Jens Pietzsch, Institute of Radiopharmaceutical Cancer Research at HZDR.

The later time point (2 hours *p.i.*) showed that most of the fluorescence intensity from Si NP-IR800 has already been excreted from the body with traces found in the kidneys which could be visualized only with high intensity projections.

3.4.3.5 *Ex vivo* analysis of Si NP-IR800

After the whole body fluorescence scan of the Si NP-IR800 injected to mice 2 hours *p.i.*, the mice was excised for a semi-quantitative analysis. The main visceral organs of the mice along with the urine sample were scanned for fluorescence intensities and draw a conclusion regarding the fate of the NPs. The *ex vivo* analysis of the excised mice is displayed in **Figure 3.4.13**. The *ex vivo* biodistribution studies of the dissected organs confirmed that the majority of the injected sample is excreted from the body in the form of urine exhibiting maximum fluorescence and slight distribution of fluorescence in organs like kidneys, lungs and spleen.

Figure 3.4.13: *Ex vivo* optical imaging of Si NP-IR800 2 h *p.i.*

Optical imaging scans of excised body organs 1-11, after 2 h of intravenous injection of Si NP/IR800 in healthy mice.

For a comprehensive analysis of the fate of Si NP-IR800 conjugates, urine from the mice was collected and analyzed (1 hour *p.i.*). The urine samples were analyzed by TLC (eluent: methanol, water) and compared to the reference sample (injected Si NP-IR800) **Figure 3.4.14**. The TLC system displayed a single spot at the origin ($R_f = 0$) corresponding to the Si NP-IR800 and no free/detached fluorescence (IR800) was observed at the TLC ($R_f = 1$). This concludes that the Si NP conjugate was rapidly eliminated from the *in vivo* system *via* renal pathway and confirms the integrity of the Si NP-IR800 in presence of complex biological milieu.

Figure 3.4.14: TLC analysis of urine of experimental healthy mice 1 h *p.i.* of Si NP-IR800.

The urine sample collected 1 h *p.i.* displays a single fluorescent peak from Si NP-IR800 at the origin ($R_f = 0$) and no traces of cleaved IR 800 dye. TLC chromatogram of free IR800 dye at the front ($R_f = 1$) (inset) TLC system: RP-TLC plates, Solvent system: Methanol.

From the biodistribution and pharmacokinetic studies of Si NP-NOTA and Si NP-IR800, it is verified that both the mono-labeled imaging agents show similar behavior for the *in vitro* (non-cytotoxic) and *in vivo* (rapid renal elimination, uniform biodistribution) analysis. These favorable results, therefore, directed us towards the development of Si NP-based dual modal imaging agent with NOTA (PET imaging) and IR 800 (optical imaging) for the simultaneous *in vivo* analysis by the combination of two imaging modalities on a single Si NP core. The development of the dual-labeled imaging agents is illustrated in **Chapter 3.5**.

3.4.5 Conclusions

Chapter 3.4 describes an elaborate synthesis and preclinical studies of ultrasmall Si NP-based mono-labeled imaging agents. The nanohybrids can be synthesized on a large scale (~ 100 mg batches) using reliable methods. The synthesis of the nanoobjects was performed in an aqueous solution at physiological conditions. Afterwards, the purification of the mono-labeled Si NPs was optimized by dialysis and size exclusion chromatography. The physico-chemical characterization confirmed neutral surface charge (zeta potential) and hydrophilicity (distribution constant) optimal for *in vitro* and *in vivo* applications.

Cell studies of the Si NP conjugates have shown no signs of cytotoxicity. Both *in vivo* PET and OI conducted for the two nanohybrids displayed comparable results. The conjugates (Si NP-NOTA and Si NP-IR800) after intravenous administration in mice were observed to be uniformly distributed for 10 minutes and eventually eliminated from the renal pathway to the bladder and urine within 60 minutes of injection. The rapid renal clearance profile and no liver accumulation makes these Si NPs exceptional candidates for imaging applications. These results thus pave a way towards the synthesis of dual-labeled imaging agents. **Figure 3.4.15** summarizes the biological studies of the mono-labeled Si NP based imaging agents.

Figure 3.4.15: Summary of synthesis, characterization and *in vitro* and *in vivo* behavior of mono-labeled Si NP conjugates.

The mono-labeled nanoconjugates, Si NP-NOTA and Si NP-IR800 show high cell viability (top) as determined by MTS and LDH assay up to 72 h. The PET and OI scans for Si NP-NOTA and Si NP-IR800, respectively, exhibit rapid renal clearance with minor liver accumulation.

3.5 Development of dual-labeled Si NPs and its *in vitro* and *in vivo* evaluation

To design an efficient dual-labeled probe, it is of significant value to analyze the probes at unimodal level first. As reported in **Chapter 3.3** and **3.4**, Si NPs were functionalized with a chelator and dye respectively, which fulfills the prerequisites for imaging studies. Here, the aim was to decorate the Si NP platform with modular ligands for PET (NOTA-Bn-SCN) and OI (IR 800) simultaneously to attain dual-modality and exploit the properties of the bimodal nanoconjugates synergistically. **Figure 3.5.1** shows the stepwise synthesis of Si NP-IR800-NOTA.

Figure 3.5.1: Synthesis of dual-labeled Si NP-based imaging agents with ^{64}Cu binding NOTA for PET and IRDye 800CW for OI.

3.5.1 Synthesis and characterization of Si NP-IR800-NOTA conjugate

To attain bimodality, the amine-terminated Si NPs were reacted simultaneously with a BFCA and a fluorescent marker. The ligands (NOTA-Bn-SCN and NHS-IR 800) evaluated for the development of mono-labeled imaging agents (**Chapter 3.4**) with satisfactory pharmacokinetic behavior for *in vivo* systems, were a suitable choice for this synthesis. To analyze the degree of functionalization and produce uniformly functionalized Si NPs *via* a thiourea and amide linkage, respectively, two synthesis strategies were carried out: one pot synthesis and stepwise reaction.

One-pot synthesis of Si NP-IR800-NOTA conjugate

For one pot synthesis, 1 mg (500 μ L) water-dispersed amine-terminated Si NPs were adjusted to pH 7.5. The ligands NOTA ($\sim 60 \mu$ g, 0.1μ mol/mg Si NP in 20 μ L) and IR 800 ($\sim 75 \mu$ g, 0.05μ mol/mg Si NP in 20 μ L) were simultaneously added to the reaction mixture and stirred overnight at 35 $^{\circ}$ C. (**Scheme 3.5.1**). The resulting Si NP-IR800-NOTA solution was purified by dialysis and SEC. The SEC fractions were analyzed by TLC, UV spectroscopy and radiolabeling experiments as performed for mono-labeled conjugates (**Section 3.4.1.1 and 3.4.3.2**).

The straightforward unimodal approach of modification was not applicable for the bimodal Si NPs as it resulted into a mixture of SiNP-IR800-NOTA, Si NP-IR800 and Si NP-NOTA conjugates. A probable reason for this variation owes to the different reactivities of the isothiocyanate group present on NOTA-Bn-SCN and the active succinimide ester on the NHS-IR800 that compete for the amines on the Si NP surface. To this end, a new stepwise synthesis strategy was applied for synthesizing Si NP-IR800-NOTA.

Scheme 3.5.1: One-pot and stepwise synthesis method for the synthesis of Si NP-IR800-NOTA.

Stepwise synthesis of Si NP-IR800-NOTA conjugate

The stepwise synthesis of Si NP-IR800-NOTA could evade the disadvantages of the one-pot method of synthesis. In this method, the amine-terminated Si NP (pH 7.4) were first reacted to activated NHS-IR 800 (70 μg , 0.05 $\mu\text{mol}/\text{mg}$ Si NP in 100 μL) and stirred for 8 hours at 35 °C (**Scheme 3.5.1**). The unreacted IR800 dye hydrolyzes in aqueous medium and does not interfere with further reactions. NOTA-Bn-SCN (60 μg) was then added next to the Si NP-IR800 solution and stirred overnight. The second reaction was performed *in situ*. This assures the dual labeling of all the Si NPs sustaining the uniformity of the Si NP-IR800-NOTA conjugates.

Fluorescence and radiolabeling studies of the SEC fractions confirmed the covalent attachment of the IR 800 and NOTA respectively to the Si NP core (**Figure 3.5.2**).

Figure 3.5.2: Complementary radio-TLC and fluorescence-TLC of Si NP-IR800-NOTA.

A) EDTA incubated $^{64}\text{Cu}(\text{II})$ -labeled Si NP-IR800-NOTA. (Radio-iTLC SG; Mobile phase: $\text{NH}_4\text{OAc}:\text{MeOH}$ 1:1 (v/v); ^{64}Cu]Cu-Si NP-IR800-NOTA $R_f = 0$, ^{64}Cu]Cu-EDTA $R_f = 1$.) B) Fluorescence TLC of Si NP-IR800-NOTA measured under IR short channel (710 – 730 nm). (RP-TLC; Mobile phase: Methanol; Si NP-IR800-NOTA $R_f = 0$).

3.5.2 Photophysical characterization of Si NP-IR800-NOTA and comparison with mono-labeled imaging agents

A comparative analysis of the UV-Vis photophysical characterization of Si NP-IR800 and Si NP-IR800-NOTA is demonstrated in **Figure 3.5.3**. The absorbance maxima (λ_{abs}) for both Si NP-IR800 and Si NP-IR800-NOTA are observed at 611 nm with typical cyanine shoulders at 470 nm for both, which confirms the conjugation of the IR 800 dye (λ_{abs} 792 nm) to the Si NPs (**Section 3.4.3.2**). The maximum emission (λ_{em}) on the other hand, for Si NP-IR800 and Si NP-IR800-NOTA is observed at 754 nm and 748 nm respectively. The data represents that the fluorescence intensity of the mono-labeled Si NP-IR800 to be higher than the dual-labeled Si NP-IR800-NOTA conjugate. It is assumed that the difference could be due to the presence of additional ligand NOTA which tends to quench the emission to certain extent. **Table 3.5.1** presents the absorbance (λ_{abs}) and emission (λ_{em}) and the photoluminescent quantum yields calculated for the Si NPs and its conjugates. The quantum yield calculation of the dual-labeled Si NP conjugates was performed similar to mono-labeled Si NP-IR800, using standard NHS-IR800 dye as a reference. **Table 3.5.1** comprises of collective data of the photophysical behavior of the Si NP conjugates.

Figure 3.5.3: Comparative absorption (A) and emission spectra (B) of Si NP-IR800-NOTA, Si NP-IR800 and NHS-IR 800 dye in water.

Normalized spectra for comparison between the absorbance and emission of dual-labeled Si NP-IR800-NOTA and mono-labeled Si NP-IR800 using standard IR 800 dye as reference.

Table 3.5.1: Comparative analysis of photophysical properties of mono-labeled vs dual labeled Si NPs

	IR800 dye (nm)	Si NPs (nm)	Si NP-IR800 (nm)	Si NP-IR800-NOTA (nm)
Absorbance (λ_{abs})	775	355	612	611
Absorbance (shoulder)	700	-	470	470
Fluorescence (λ_{ems})	792	456	754	748
PLQY	14.2%	15%	22.5%	15.7%

Surface charge

The zeta potential (ζ) value of $-5.1 \text{ mV} \pm 3.5 \text{ mV}$ obtained for the Si NP-IR800-NOTA confirms its zwitterionic behavior of the dual-labeled conjugate. The values show slight variations from the mono-labeled Si NPs (**Table 3.5.2**) which depends on the chemical structure and the influence of functionalities present on the NOTA and IR 800 ligands. But the nearly zero values for all the Si NP conjugates owes to the limited surface functionalization performed which is detectable by imaging techniques yet not hampers the surface properties of the conjugates.

3.5.3 Radiolabeling studies of dual-labeled Si NP-IR800-NOTA

The radiolabeling experiments were performed according to the standard protocol followed in **Section 3.4.11**. The Si NP-IR800-NOTA conjugates were labeled with $[^{64}\text{Cu}]\text{CuCl}_2$ radionuclide at pH 5.2 at room temperature for 30 minutes. $[^{64}\text{Cu}]\text{Cu-Si NP-}$

IR800-NOTA was obtained in > 90% RCY with a specific activity of ~2 MBq/ 100 µg. The variation in the parameters like increasing the reaction time and slight elevation of the temperature (35 °C) did not potentially affect the radiolabeling of the Si NP conjugate. The radio-labeled Si NPs were purified by SEC to obtain ~98% radiochemical purity.

An effect of pH on the radiolabeling efficiency of Si NP-IR800-NOTA was also determined. The percentage $^{64}\text{Cu}(\text{II})$ bound to the Si NP conjugate was plotted against the pH and is represented as **Figure 3.5.4**. The results show that highest radiolabeling is achieved at pH 4–6 for ^{64}Cu -Si NP-IR800-NOTA.

Figure 3.5.4: pH dependent $^{64}\text{Cu}(\text{II})$ radiolabeling studies of Si NP-IR800-NOTA.

The extent of radiolabeling was determined by radio-iTLC after 30 min of reaction and Radio-HPLC purification. The results were plotted as pH vs percentage $^{64}\text{Cu}(\text{II})$ bound to Si NP-IR800-NOTA.

3.5.4 Quantitative estimation of the amount of fluorophore and the radiolabel on the Si NP conjugate

Radiometric titration of Si NP-IR800-NOTA

To determine the amount of NOTA chelator attached to the dual-labeled Si NP-IR800-NOTA, radiometric titrations were performed with $^{64}\text{Cu}(\text{II})$. The protocol followed was same as for Si NP-NOTA (**Figure 3.5.5**).

The calculations yielded ~0.091 µmol of NOTA conjugates per mg of Si NP-IR800-NOTA, which is almost the same as obtained for mono-labeled Si NP-NOTA conjugate (0.09 µmol/mg Si NP-NOTA).

Figure 3.5.5: Radiometric titration curve for $^{64}\text{Cu}(\text{II})$ and $\text{Cu}(\text{II})$ labeled Si NP-IR800-NOTA.

(Experimental value: $\sim 0.091 \mu\text{mol}$ of NOTA /mg Si NP-IR800-NOTA determined by radio-TLC analysis (Method III)).

Determination of lipophilicity of Si NP-IR800-NOTA

1-octanol/water distribution experiments examined the hydrophilicity of $[^{64}\text{Cu}]\text{Cu-Si NP-IR800-NOTA}$ to determine the biological behavior of the conjugate under physiological conditions. A negative $\log D_{7.4}$ value of -3.84 ± 0.10 assures the hydrophilicity and preserved biocompatibility of the dual-labeled Si NPs. The distribution ratio varies slightly from the mono-labeled Si NP-NOTA (-4.12 ± 0.04) which is probably due to the additional IR 800 ligand in the case (**Table 3.5.2**).

The studies suggest that the Si NP conjugates are highly soluble under aqueous conditions, an important requisite for further *in vitro/in vivo* studies.

Table 3.5.2: Zeta potential and distribution constants of Si NP conjugates.

(Zeta potential was measured in water and distribution ratios for 1-octanol/water system)

Si NP conjugates	Zeta potential ζ (mV)	$\log D_{7.4}$
Si NPs	$-1.1 \pm 3.9 \text{ mV}$	–
$[^{64}\text{Cu}]\text{Cu-Si NP-NOTA}$	$-2.9 \pm 4.7 \text{ mV}$	-4.12 ± 0.04
Si NP-IR800	$-5.4 \pm 4.1 \text{ mV}$	–
$[^{64}\text{Cu}]\text{Cu-Si NP-IR800-NOTA}$	$-5.1 \pm 3.5 \text{ mV}$	-3.84 ± 0.10

Estimation of amount of IR800 conjugated to Si NPs

The amount of dye conjugated to the NPs was determined by UV spectroscopy analysis based on Beer-Lambert's Law. For the experiment, known samples of free IR 800 dye were prepared and measured for their absorbance and plotted on a graph against the concentration. A known amount of Si NP-IR800-NOTA solution was then measured and compared for the detection of the amount of the conjugated IR 800 dye (**Figure 3.5.6**). The results reveal an estimate IR 800 dye as 0.009 $\mu\text{mol}/\text{mg}$ Si NP-IR800-NOTA which in this case is less compared to its mono-labeled analogue (0.013 $\mu\text{mol}/\text{mg}$ Si NP-IR800).

Figure 3.5.6: Quantification of IR 800 dye conjugated to Si NP-IR800-NOTA (Experimental value ~ 0.013 μmol IR800 / mg Si NP-IR800).

A comparative summary of the conjugated ligands (NOTA and IR 800 dye) to the surface amine groups of Si NPs (**Table 3.5.3**).

Table 3.5.3: Quantification of ligands bound to Si NP conjugates.

Si NP conjugate	Ligand	Amount ($\mu\text{mol}/\text{mg}$ Si NP conjugates)
Si NP-NOTA	NOTA	~0.09
Si NP-IR800-NOTA	NOTA	~0.091
Si NP-IR800	IR 800	0.013
Si NP-IR800-NOTA	IR 800	0.009
Si NP-B4	Bispidine (B4)	~0.052

3.5.5 Cell Cytotoxicity Studies of the Si NP-IR800-NOTA

Biocompatibility studies for Si NP-IR800-NOTA were performed to assess the cellular toxicity related to the multifunctionalization of the NPs. The *in vitro* studies were performed following the same protocol utilized for the mono-labeled Si NPs to attain a comparative cytotoxicity data. The CyQ viability assay for the HEK 293 cells incubated with Si NP-IR800-NOTA portrays a permissible value of > 95% for 24, 48 and 72 hour time points, even at the higher concentrations (100 $\mu\text{g}/\text{mL}$) of the conjugate (**Figure 3.5.7**).

Figure 3.5.7: *In vitro* cell studies of Si NP-IR800-NOTA in HEK293 cell lines.

LDH assay results of ~5% cell death demonstrated that the dual modal Si NP conjugates show negligible LDH leakage into the cell media due to cell membrane disintegration for all four concentrations up to 3 days. A value considered to be well below the toxic threshold according to the Iso Norm 10993–5:2009 was retained for the bimodal Si NP-IR800-NOTA. There was no statistical difference when comparing the toxic effects between mono-labeled and dual-labeled Si NP conjugates. The MTS viability assay supported the CyQuant and LDH results presenting the lack of any significant nephrotoxicity with $\geq 97\%$ viability of the cells after 24, 48 and 72 hours (**Figure 3.5.8**).

Figure 3.5.8: Comparative *in vitro* analysis of Si NP conjugates. The cell viability and membrane lysis studies (MTS and LDH assay) for Si NP-NOTA, Si NP-IR800 and Si NP-IR800-NOTA upon 48 h of incubation in HEK293 cells.

Based on these results, it is apparent that the dual modal conjugate does not lead to any substantial modifications in the cellular morphology and that the cellular metabolism is in the range of the negative controls. There are no toxic effects under the examined conditions. Thus, it is confirmed that the Si NPs are biocompatible and ready for the *in vivo* trials with its bimodal functionality.

3.5.6 *In vivo* PET analysis and biodistribution studies, urine analysis and *ex vivo* examination of [^{64}Cu]Cu-Si NP-IR800-NOTA

3.5.6.1 Small animal PET studies and urine analysis of dual-labeled [^{64}Cu]Cu-Si NP-IR800-NOTA

The [^{64}Cu]CuCl₂ radiolabeled and purified [^{64}Cu]Cu-Si NP-IR800-NOTA was injected intravenously (~10 MBq each) to a set of 4 NMRI nu/nu healthy mice (standard deviation = 4). The biodistribution of the dual-labeled [^{64}Cu]Cu-Si NP-IR800-NOTA was recorded by whole body PET scans at 5, 20, 30 and 60 minute time points. The **Figure 3.5.9** shows a fast clearance of the majority of the radiolabeled Si NP bioconjugates from the kidneys to bladder. The scan suggests that the bimodal Si NPs, after intravenous administration distribute uniformly in the whole body within the first 5–10 minutes and

then start eliminating from the body after glomerular filtration attributing to their ultrasmall size and zwitterionic surface charge.

Figure 3.5.9: *In vivo* PET imaging of $[^{64}\text{Cu}]\text{Cu-Si NP-IR800-NOTA}$ at different time points.

Whole body PET scans after a single intravenous injection of $[^{64}\text{Cu}]\text{Cu-Si NP-NOTA}$ ($n = 4$) in NMRI nu/nu mice after 5, 20, 30, 60 min *p.i.*

The urine from the experimental mice was collected 5 min and 1 h *p.i.* and the activity was examined by radio-TLC system (“**Method I**” **Section 5.2**). The $[^{64}\text{Cu}]\text{Cu-Si NP-IR800-NOTA}$ remains at the origin ($R_f = 0$) on the radio RP-TLC and the $[^{64}\text{Cu}]\text{Cu-EDTA}$ moves to the solvent front ($R_f = 0.9$) (**Figure 3.5.10**). The results show no significant differences in the three chromatograms with most of the radioactive compounds present at the origin corresponding to the bimodal Si NPs ($R_f = 0$). $[^{64}\text{Cu}]\text{Cu-EDTA}$ is observed in the injected sample and the urine 1 hour *p.i.* This data confirms the stability of the radiolabeled nanoconjugates in the physiological system of the mice with intact PET imaging modality. Additionally, no signs of demetalation of $^{64}\text{Cu(II)}$ or cleavage of $[^{64}\text{Cu}]\text{Cu-NOTA}$ were observed within the biological system. This proves the covalent attachment of the radioactive chelating moiety, NOTA to the Si NPs *via* metabolically stable thiourea bonds.

Figure 3.5.10: Radio-TLC analysis of the urine sample of $[^{64}\text{Cu}]\text{Cu-Si NP-IR800-NOTA}$ injected mice. EDTA incubated SEC purified $[^{64}\text{Cu}]\text{Cu-Si NP-IR800-NOTA}$ was injected in a NMRI nu/nu. The urine samples (red) 5 min and 1 h *p.i.* were analyzed by Radio-TLC. TLC system: RP-TLC plates, 2M NH_4OAc : MeOH (1:1, v:v), $[^{64}\text{Cu}]\text{Cu-Si NP-IR800-NOTA}$ $R_f = 0$.

3.5.6.2 *Ex vivo* biodistribution and autoradiography studies of $[^{64}\text{Cu}]\text{Cu-Si NP-IR800-NOTA}$

The $[^{64}\text{Cu}]\text{Cu-Si NP-IR800-NOTA}$ injected (~ 10 MBq) 4 NMRI nu/nu healthy mice were excised 5 min, 1 hour and 24 hour post injection respectively for the *ex vivo* autoradiography studies. The standardized uptake value (SUV_{BD}) and the percent injected dose (%ID) values were used to evaluate the preferred mode of elimination of the particles from the body and the accumulation of these particles in different tissues and organs. The data were also used to compare the biodistribution of the dual-labeled imaging agent to that of previously studied and published mono-labeled agent $[^{64}\text{Cu}]\text{Cu-Si NP-NOTA}$, which lacks the fluorophore in its structure.

%ID values based analysis of [^{64}Cu]Cu-Si NP-IR800-NOTA

After 5 minutes of injection, the organs distribution studies depicted that the majority of the radioactive NPs constituting ~40% [^{64}Cu]Cu-Si NP-IR800-NOTA was eliminated via the urine with kidneys, liver and intestine accounting for 5%, 5% and 3% respectively (**Figure 3.5.11**). After 1 hour, the radioactive content in the urine showed upto 85% and less than 5% was detected in the kidneys, liver and intestine. The other organs show negligible amount of activity confirming the extremely fast complete renal clearance of these hydrophilic Si NPs from the *in vivo* system within 1 hour time post injection. The 24 hour *p.i.* study showed the presence of ~ 99% activity in the urine with 1% in the kidney and liver and negligible in other body parts.

SUV based PET analysis of [^{64}Cu]Cu-Si NP-IR800-NOTA

The SUV data analysis after 5 minutes of administration of [^{64}Cu]Cu-Si NP-IR800-NOTA showed major amount of activity concentration in the kidneys whereas all the other organs showed a SUV of 0.5–2, which refers to a smooth whole body biodistribution pattern (**Figure 3.5.11**). In 1 hour, the SUV in the kidney, liver and spleen were observed to be around 1.5, 0.7 and 0.3. The SUV in other organs were substantially lower, which further went down to 0.3 and 0.1 for kidneys and liver and negligible for all the other organs.

Having a closer look at the data and evaluating the mode of elimination of the Si NPs, we compared the %ID and SUV values of the kidneys (to check renal excretion) and the liver (to check hepatobiliary mode of elimination) (**Figure 3.5.12**).

The values clearly state that the mode of elimination for [^{64}Cu]Cu-Si NP-IR800-NOTA is renal. The particles show no sign of sticking in the renal medulla and are easily passing through the glomerular pores in as short as 1 hour time post injection.

Figure 3.5.11: Biodistribution and autoradiography studies of $[^{64}\text{Cu}]\text{Cu-Si NP-IR800-NOTA}$ in NMRI nu/nu mice.

Quantitative analysis of time-activity curves as %ID (mean $\pm$ SEM) (left) and SUV (mean $\pm$ SEM) (right) ($n = 4$) A: 5 min, B: 1 h and C: 24 h after single intravenous administration. (%ID: Highest amount of activity in urine after 5 min, 1 h and 24 h).

Figure 3.5.12: Comparative quantification of renal versus hepatobiliary mode of elimination for $[^{64}\text{Cu}]\text{Cu-Si NP-IR800-NOTA}$ in NMRI nu/nu mice.

Si NP-IR800-NOTA prefer renal mode of elimination over hepatobiliary excretion.

3.5.6.3 Optical imaging *ex vivo* biodistribution studies of Si NP-IR800-NOTA

The Si NP-IR800-NOTA ($\sim 120 \mu\text{g}$) were injected intravenously in NMRI nu/nu healthy mice ($n=4$). The mice were excised after 5 min, 1 h and 24 h *p.i.* The fluorescence was studied semi-quantitatively for different organs. The results show that the majority of the Si NP conjugates were excreted into the urine *via* bladder. The maximum fluorescence intensity projections as after the elimination of the majority of the injected sample *via* urine detects only traces of fluorescence in the organs. **Figure 3.5.13** demonstrates the organs of the excised mice 5 minutes, 1 hour and 24 hour post injection of the bimodal Si NP conjugate.

In the first 5 minutes, the fluorescence is observed uniformly distributed in the whole body and also actively filtered through the kidneys which shows prominent signals for the urine sample. In 1 hour, the activity has gone down quite low and major signal is observed in the urine, stomach and intestine. For 24 hour post injection excision sample, traces of fluorescence from the Si NPs are detected majorly in the urine with traces in the stomach.

Figure 3.5.13: Ex vivo optical imaging of Si NP-IR800-NOTA at different time points *p.i.*

Optical imaging scans of excised body organs numbered 1-21, after 5 min, 1 h and 24 *p.i.* in NMRI nu/nu mice.

The complementary biocompatibility and rapid renal clearance of the Si NP-IR800-NOTA achieved by PET and optical imaging, confirms the bimodality of the Si NP system. The results also displayed excellent pharmacokinetics and extraordinary biocompatibility of the system, which is prerequisite for any NP platform adapted for *in vivo* studies. The most acceptable reason for this behavior is the minimum required surface modification performed on the Si NP surface to achieve the imaging goals without hampering the hydrodynamic diameter and surface charge of the particles.

3.5.7 Comparative analysis of [⁶⁸Ga]Ga-Si NP-IR800-NOTA and [⁶⁴Cu]Cu-Si NP-IR800-NOTA in A431 tumor bearing mice

To discover the effect of radiometals on the pharmacokinetics of the novel dual-labeled Si NP-IR800-NOTA conjugate, *in vivo* PET and biodistribution experiments were performed with two different radionuclides. NOTA being an efficient chelator for ⁶⁸Ga as well as for ⁶⁴Cu, provided us the opportunity to develop a correlation between the *in vivo* behavior and examine the effect of different radionuclides on the biodistribution pattern of the conjugates. The time frame for the two biodistribution experiments (⁶⁴Cu and ⁶⁸Ga) was set for up to 24 hours which was optimized while investigating the mono-labeled conjugates. ⁶⁴Cu ($t_{1/2} = 12.7$ hours) radionuclide has an added advantage to measure the PET for a longer period compared to ⁶⁸Ga, ($t_{1/2} = 68$ minutes) due to substantial differences in their half-lives. In this section, we present the quantitative (PET imaging) and semi-quantitative (optical imaging) results obtained after the *in vivo* biodistribution, pharmacokinetics and the *ex vivo* examination of the [⁶⁸Ga]Ga-Si NP-IR800-NOTA and

[⁶⁴Cu]Cu-Si NP-IR800-NOTA administered mice. The tumor mice were used to study the pharmacokinetics of the Si NP conjugates and to check the nonspecific tumor binding.

3.5.7.1 *In vivo* PET results of [⁶⁸Ga]Ga-Si NP-IR800-NOTA in A431 tumor bearing mice¹⁶

The Si NP-IR800-NOTA were radiolabeled in MES buffer at room temperature with ⁶⁸Ga radionuclide in ~ 95% RCY and purified by SEC Sephadex G-25 in 80% radiochemical purity, for the *in vivo* experiments. The preliminary small animal experiments were performed by intravenous administration of ~23 MBq of [⁶⁸Ga]Ga-Si NP-IR800-NOTA (~ 60 µg in 200 µL) to a set of two NMRI nu/nu mice with A431 tumor in the right hind leg (n = 2). As a minimal prerequisite, the *in vivo* studies were carried out *via* intravenous injections rather than local injection in order to get reliable pharmacokinetic profiles of the NPs.

The chosen time points for recording the PET and fluorescence imaging for [⁶⁸Ga]Ga-Si NP-IR800-NOTA was 30 minutes, 2 and 24 hours. The whole body PET image of the [⁶⁸Ga]Ga-Si NP-IR800-NOTA recorded after 30 minutes of administration is displayed in **Figure 3.5.14**. The semi-quantitative PET scan clearly portrays the maximum radioactivity from the [⁶⁸Ga]Ga-Si NP-IR800-NOTA in the heart of the mice and in the circulatory system with a little amount detected in other vital organs and tissues.

The time-activity graph (**Figure 3.5.14 A**) displays a comparative biodistribution analysis of the bimodal [⁶⁸Ga]Ga-Si NP-IR800-NOTA in the blood, heart, kidneys, muscles and tumor. The Si NPs are distributed in the whole body through the blood circulation at all points of time and eliminate *via* the designated renal pathway. No significant liver accumulation validates the renal clearance of the Si NPs and thus the biocompatible hydrodynamic diameter and the zwitterionic nature of the system. The values obtained are in accordance with the data attained for dual-labeled and mono-labeled ⁶⁴Cu(II) ligated [⁶⁴Cu]Cu-Si NP-IR800-NOTA and [⁶⁴Cu]Cu-Si NP-NOTA.

¹⁶ ⁶⁸Ga radiolabeling of Si NP-IR800-NOTA and the animal experiments were performed by Dr. Ralf Bergmann at the Institute of Radiopharmaceutical Cancer Research, HZDR.

Figure 3.5.14: PET scan of $[^{68}\text{Ga}]\text{Ga-Si NP-IR800-NOTA}$ 30 min *p.i.* Whole body PET scan, A) Time-activity curve B) Tumor to muscle ratio of $[^{68}\text{Ga}]\text{Ga-Si NP-IR800-NOTA}$ in NMRI nu/nu mice.

The high tumor to muscle ratio and tumor to blood ratio (**Figure 3.5.14 B**) suggest the non-specific targeting of the tumor tissue, occurring by the perfusion of the ultrasmall Si NPs in the tumor cells. Due to the crucial < 10 nm size of the imaging agent, the chances of EPR is eradicated. The graph shows that the maximum passive targeting was achieved 20–30 minutes post injection of the $[^{68}\text{Ga}]\text{Ga-Si NP-IR800-NOTA}$. The radioactivity tends to wash away quickly from the tumor and the particles are again in the blood vessels making their way towards the kidneys.

Another interesting and valuable data from this set of experiments was the determination of the complementarity of the dual-labeled Si NP system. The **Figure 3.5.15** displays the maximum intensity projections of the simultaneous PET and fluorescence detection of the $^{64}\text{Cu(II)}$ positrons emissions and the fluorescence from the IR 800 present on the Si NP surface. The complementary dual modal detection of $[^{68}\text{Ga}]\text{Ga-Si NP-IR800-NOTA}$ in the bladder 2 hours post injection confirmed the uniformity of the system and the presence of both the imaging modalities on a single Si NP platform.

Figure 3.5.15: PET/CT and OI/CT whole body scans of [⁶⁸Ga]Ga-Si NP-IR800-NOTA injected mice. Similar PET (A) and OI (B) images of [⁶⁸Ga]Ga-Si NP-IR800-NOTA injected in A431 tumor bearing NMRI nu/nu mice validates the dual modality of the Si NP conjugate with maximum detected activity in the urinary bladder.

3.5.7.2 *In vivo* biodistribution studies [⁶⁴Cu]Cu-Si NP-IR800-NOTA in A431 tumor bearing mice by small animal PET

The second set of experiments on A431 tumor bearing mice was performed by the intravenous injection of ⁶⁴Cu labeled Si NP-IR800-NOTA followed by the *in vivo* and *ex vivo* analysis. The whole body PET scan of the mice 2 hours post injection is presented in **Figure 3.5.16**. Similar to the other data, the [⁶⁴Cu]Cu-Si NP-IR800-NOTA conjugates in this case are also visualized passing through the kidneys and ending up in the bladder and eliminated by the urine. Due to the absence of a tumor targeting moiety, no significant tumor accumulation is observed for the mono as well as dual-labeled Si NPs as in the case of healthy mice.

A simultaneous comparative PET and fluorescence read out study of the [⁶⁴Cu]Cu-Si NP-IR800-NOTA 1, 2 and 24 hour post injection, demonstrated the complementarity of the system for the tumor bearing mice, too (**Figure 3.5.17**). The results are also in agreement with the ⁶⁸Ga labeled dual-labeled nanoconjugates with a similar pharmacokinetic behavior.

Figure 3.5.16: Whole body PET scan of A431 right hind leg tumor bearing NMRI nu/nu mice 2 h *p.i.* of $[^{64}\text{Cu}]\text{Cu-Si NP-IR800-NOTA}$.

Figure 3.5.17: PET and OI scans of $[^{64}\text{Cu}]\text{Cu-Si NP-IR800-NOTA}$ injected A431 tumor bearing mice. PET (A) and OI (B) images of $[^{64}\text{Cu}]\text{Cu-Si NP-IR800-NOTA}$ in NMRI nu/nu mice 1 h and 24 h *p.i.* The maximum activity is already eliminated within 1 h *via* bladder. MIP analysis detects traces of radioactivity and fluorescence in the kidneys.

3.5.7.3 *Ex vivo* and autoradiography studies of [^{64}Cu]Cu-Si NP-IR800-NOTA

Due to the longer half-life of ^{64}Cu , unlike ^{68}Ga , an *ex vivo* analysis was also possible for the system along with the fluorescence measurements. As expected, the maximum activity and fluorescence from [^{64}Cu]Cu-Si NP-IR800-NOTA was detected in the kidneys and the urine, followed by minor quantities in the intestine, stomach and liver (**Figure 3.5.18**). The data obtained are consistent to their [^{68}Ga]Ga-Si NP-IR800-NOTA labeled counterparts.

Figure 3.5.18: *Ex vivo* PET (A) and optical imaging (B) scans of excised organs for [^{64}Cu]Cu-Si NP-IR800-NOTA injected A431 tumor xenograft mice 24 h p.i.

3.5.8 Conclusions

The dual-labeled Si NPs were prepared by stepwise synthesis method and obtained in high purity after several optimizations. Si NP-IR800-NOTA was characterized for its surface properties which shows comparable values to its mono-labeled analogues. The cell studies also displays no signs of cytotoxicity and cell damage. Therefore, the bimodal conjugate was assessed for its *in vivo* behavior.

The most interesting conclusion drawn from the small animal experiments was that no significant effect of the radionuclide ^{64}Cu or ^{68}Ga was observed on the pharmacokinetics, elimination behavior, and fate of Si NPs. Another important aspect was the complementary data obtained from the dual imaging modalities, NOTA and IR 800, present on Si NP-IR800-NOTA, confirming the successful synthesis of a silicon-based dual-labeled system. The preclinical trials on the healthy and tumor bearing mice demonstrated pharmacokinetics similar to the mono-labeled conjugates Si NP-IR800 and Si NP-NOTA. The slight non-specific tumor accumulation of the Si NP conjugate detected was also quickly washed away by the blood circulation not leading to any significant differences in the elimination pattern. The comparable conduct of the mono-labeled as well as dual-labeled Si NP-based nanohybrids in the living system can be explained by the physicochemical properties possessed by the modified Si NPs.

Si NP-IR800-NOTA displays excellent pharmacokinetics with high biocompatibility. The low unspecific binding and fast renal clearance makes it ideal for labeled Si NPs with targeting moieties and their use for *in vivo* imaging. With these characteristic properties, the bimodal conjugate can be used as a potential diagnostic tool for cancer imaging. After the successful synthesis of the dual-labeled core, the next step was to investigate their potential with a tumor-targeting agent that is reported in **Chapter 3.6**.

3.6 Investigation of mono-labeled and dual-labeled Si NPs as targeted imaging agents

The successful synthesis of non-targeting Si NP-based mono and dual-labeled agents and the promising small animal biodistribution results directed us towards the development of targeting Si NPs. This chapter deals with the synthesis, purification, photophysical and physico-chemical characterization of the targeting Si NP-based nanohybrids.

The initial coupling reactions were performed with the peptide 5B9 to analyze the extent of fabrication and stability of the NPs as multimodal systems. The peptide 5B9 is the first known chimeric anti-platelet factor 4/heparin complexes IgG1 antibody that has a human Fc fragment. It fully mimics the effect of Heparin-induced thrombocytopenia (HIT) human antibodies. This property leads to a detailed study of this disease HIT [279-281].

The biological vector conjugations to the Si NP-IR800 and Si NP-IR800-NOTA lead to a viable analysis and characterization of the Si NP bioconjugates. Hexaethylene glycol was used as a bifunctional linker with an active-NHS ester on one end and maleimide on the other to couple amine-terminated Si NPs and thiol-terminated 5B9 peptide, respectively. The advantage of using such linkers is that it does not deeply affects the crucial properties of the NPs, like the size and surface charge and thus, the biocompatibility of the system. Ethylene glycols are renowned linkers for biological applications, as they tend to increase the blood retention time of the bioconjugates with no significant drawbacks.

3.6.1 Synthesis of Si NP-PEG-5B9 peptide conjugates¹⁷

For the synthesis of Si NP-IR800-PEG-5B9 and Si NP-IR800-NOTA-PEG-5B9, Si NP-IR800 (mono-labeled) and Si NP-IR800-NOTA (dual-labeled) (**Chapter 3.4 and 3.5**) were used as precursors. The PEG moieties further allow the modification of the Si NPs to the peptide through the thiol-maleimide coupling.

The first step in this synthesis was the addition of the linker, consisting of oxyethylene groups to the Si NPs to gain maleimide-functionalized NPs (**Figure 5.6.1**). The second reaction of the 5B9 addition was performed *in situ* for both the fluorescent conjugates. The thiol-terminal 5B9 peptide was stirred in an aqueous solution with a reducing agent,

¹⁷ PEG in Chapter 3.6 refers to Hexaethylene glycol.

TCEP to break the disulfide linkages and to get free the sulfhydryl groups. The reactions were run in aqueous environment at physiological pH for 8 hours.

Upon multifunctionalization, due to no major differences in the molecular masses of Si NP conjugates, the purification, as well as the characterization of the samples was challenging. The NPs were dialyzed against water, followed by SEC. The fractions collected from SEC were further analyzed by measuring the DLS, Zeta potential, photophysical properties and electrophoresis.

3.6.2 Characterization of the targeted Si NP conjugates

The absorption and emission properties of the purified samples, Si NP-IR800-PEG-5B9 and Si NP-IR800-NOTA-PEG-5B9 were compared to their non-targeted counterparts Si NP-IR800 and Si NP-IR800-NOTA. A comparative study of absorption and emission spectra of the conjugates is represented in **Figure 5.6.2**.

The results reveal no changes in the spectral band positions of the bioconjugates upon coupling the 5B9 peptide *via* the PEG linker. However, an increase in the PLQY of the Si NP bioconjugates is observed compared to its non-targeted analogues.

The absorption, emission and calculated photoluminescent quantum yields of the fluorescent Si NP conjugates (Si NP-IR800, Si NP-IR800-NOTA, Si NP-IR800-PEG-5B9 and Si NP-IR800-NOTA-PEG-5B9) along with Si NPs and IR 800 dye are summarized in **Table 5.6.1**.

Figure 5.6.1: Synthesis of Si NP-IR800-PEG-5B9 and Si NP-IR800-NOTA-PEG-5B9.

Figure 5.6.2: Absorption (A) and emission spectra (B) of Si NP conjugates and IR 800 dye.

Table 3.6.1: Photophysical properties of non-targeted and targeted Si NPs.

	IR800 dye (nm)	Si NPs (nm)	Si NP-IR800 (nm)	Si NP-IR800-NOTA (nm)	Si NP-IR800-PEG-5B9 (nm)	Si NP-IR800-NOTA-PEG-5B9 (nm)
λ_{abs}	775	355	612	611	619	619
λ_{abs} (shoulder)	700	-	470	470	470	470
λ_{em}	792	456	754	748	753	753
PLQY	14.2%	15%	22.5%	15.7%	29.82%	26.29%

Zeta potential measurements of the fluorescent, targeted and non-targeted Si NPs was measured in water between pH 7.2–7.5. A comparative analysis of the surface charge of Si NP conjugates is reported in **Table 3.6.2**. The zeta potential values of Si NP-IR800-PEG-5B9 (-19.1 ± 6.6) and Si NP-IR800-NOTA-PEG-5B9 (-8.3 ± 5.3) do not display huge deviations from the non-targeting Si NP conjugates. As discussed earlier, the low surface modification, helps in retaining the inherent and surface properties of the Si NPs, therefore, no huge deflections were observed from the nearly zero surface charge for all the nanoconjugates. The conservation of the surface charge and size of the modified Si NPs play a major role in the noncytotoxic behavior and favorable pharmacokinetic patterns attained by the conjugates.

Table 3.6.2: Summary of surface charge (Zeta potentials) of Si NPs and its conjugates.

Si NP conjugates	Zeta potential ζ (mV)
Si NPs	-1.1 ± 3.9
Si NP-NOTA	-2.9 ± 4.7
Si NP-IR800	-5.4 ± 4.1
Si NP-IR800-NOTA	-5.1 ± 3.5
Si NP-IR800-PEG-5B9	-19.1 ± 6.6
Si NP-IR800-NOTA-PEG-5B9	-8.3 ± 5.3

SDS-PAGE and Agarose gel analysis

The charge, size and mass dependent mobility of the synthesized conjugates was verified by two kinds of gel electrophoresis techniques, the sodium dodecyl sulfate polyacrylamide gel electrophoresis (SDS-PAGE) and agarose gel electrophoresis. Previous experiments confirmed the mobility of the Si NPs in the gel but the particle with or without 5B9 peptide could not be stained. Therefore, only fluorescence read out was possible for the preliminary studies.

The two peptide conjugates Si NP-IR800-PEG-5B9 and (1) Si NP-IR800-NOTA-PEG-5B9 (2) (**Figure 5.6.3**) were loaded on 10% agarose gel and run in an electric field. The gels were read under two wavelength categories, IR short channel (710 – 730 nm) and IR long channel (810 – 840 nm). The results were overlaid for better analysis.

The major interpretations and conclusions drawn from the two kinds of gel electrophoresis are:

The free IRDye 800CW ($\lambda_{ab}/\lambda_{em} = 775/792$ nm) is seen in the IR long wavelength detector and moves to the front in both the gels (**Figure 5.6.3**). The Si NP conjugates with higher molecular mass, show less mobility. It is also evident that the IR 800 dye bears higher charge compared to Si NP conjugates.

Figure 5.6.3: Agarose gel for Si NP-IR800-PEG-5B9 (1a, 1b) and Si NP-IR800-NOTA-PEG-5B9 (2a, 2b). IR short channel (710-730 nm) IR long channel (810-840 nm)

The peptide conjugates Si NP-IR800-PEG-5B9 (**1a** and **1b**) and Si NP-IR800-NOTA-PEG-5B9 (**2a** and **2b**) are observed in the both the IR short and IR long channels and therefore, can be observed in the overlaid images too.

An interesting finding is the presence of two consistent populations of fluorescent Si NPs in each sample which might be due to the non-uniformity of conjugation on the surface of Si NPs (**Figure 5.6.3** and **5.6.4**). The two bands differ slightly in their mobility and therefore, in the charge, mass and size. The presence of the two populations even after purification steps (dialysis and SEC) confirms that they bear similar retention times with comparable mass and size.

Figure 5.6.4: Agarose (left) and SDS PAGE (right) gel for Si NP-IR800-PEG-5B9 and Si NP-IR800-NOTA-PEG-5B9.

3.6.3 Conclusions

Fluorescent Si NP conjugates Si NP-IR800 and Si NP-IR800-NOTA were grafted by a bifunctional linker (PEG) to couple a receptor targeting peptide onto the Si NP surface. The Si NP bioconjugates were synthesized in aqueous medium and purified by dialysis and SEC. The analysis was performed by fluorescence measurements. Preliminary studies confirmed the attachment of the peptide 5B9 to the Si NPs via thioether bonds. The electrophoresis studies revealed a non-uniform surface modification of both the Si NP bioconjugates leading to two distinct populations in the solution.

The investigations of the targeting Si NPs, Si NP-IR800-PEG-5B9 and Si NP-IR800-NOTA-PEG-5B9 are currently under progress and not a part of this work.

Chapter 4 Conclusions and Outlook

4.1 Conclusions

The major goals achieved during the course of this dissertation are:

Development of novel bispidine bioconjugates as ^{64}Cu chelators for PET imaging:

The facile and cheap synthesis, excellent chelating efficiency, possibility to form highly stable complexes under mild conditions with favorable biodistribution patterns was an instigation to work towards the development of bifunctional bispidines as novel chelators for PET applications. A series of bispidine derivatives (**B4** – **B9**) with varying terminal functional groups were synthesized and thoroughly characterized.

Bispidine-Bn-SCN (**B4**) and Bispidine-DBCO (**B5**) forming excellent thermodynamically stable and kinetically inert complexes were selected for the development of targeted PET imaging agents.

The novel derivative **B4** was conjugated to amine-terminated TATE peptide *via* a two-step coupling/deprotection reaction to produce **B4**-EG-TATE (EG: diethylene glycol). TATE peptide is a somatostatin analogue and is expected to efficiently target Somatostatin type-2 receptors overexpressed in cancer malignancies of neuroendocrine origin.

B4-EG-TATE
(Bispidine-EG-TATE)

Bispidine coupled to SSTR-specific EG-TATE peptide

The [^{64}Cu]Cu-**B4**-EG-TATE complex was delivered in high RCY and radiochemical purity. This also proved the preservation of the labeling efficiency of **B4** after its bioconjugation.

A Sortase-A (Srt-A) mediated approach was applied for the site-specific labeling of single domain antibody 7C12 (sdAb 7C12) to generate an azide terminal complementary biomolecule to the dibenzocyclooctyne functionality present on **B5**. This chemoenzymatic strategy leads to a facile and convenient modification of the single domain antibody. The strain promoted azide-alkyne cycloaddition between **B5** and sdAb 7C12 produced a stable triazole derivative **B5**-sdAb-7C12 *via* Cu-free click reaction. Preliminary studies provided a high radiolabeling efficiency and stability of the complex [^{64}Cu]Cu-**B5**-sdAb-7C12. Cell binding studies performed with A431 cancer cell lines revealed a specific binding to the epidermal growth factor receptors and allowed further investigation in the direction of small animal PET experiments.

B5-sdAb-7C12
(Bispidine-Thr-Glu-Pro-Leu-sdAb)

Bispidine coupled to EGFR-specific sdAb 7C12 *via* azide-cyclooctyne click chemistry

Modification of ultrasmall silicon nanoparticles as PET/OI imaging agents:

Hydrothermal method was employed to yield extremely hydrophilic, multifunctional, zwitterionic amine-terminated Si NPs with a size of ~ 4 nm. Novel Si NP-IR800 and Si NP-bispidine conjugates were developed as positron emission tomography and optical imaging agents. A comprehensive characterization of the nanoconjugates confirmed the preservation of 'ultrasmall' hydrodynamic diameter and 'zwitterionic' surface charge of the Si NPs even upon its functionalization. After successful synthesis, analysis and characterization, it was concluded that the Si based nanoparticles; Si NP-NOTA and Si NP-IR800 (**Figure**) bear optimal intrinsic and surface properties to be applicable for cell studies and preclinical trials in mice.

Cell viability and proliferation analysis of Si NP-IR800 on human embryonic kidney cells (HEK 293) showed no cytotoxicity.

The biodistribution studies from both kinds of nanoparticles; Si NP-IR800 and $[^{64}\text{Cu}]\text{Cu-Si NP-NOTA}$ performed in NMRI nu/nu healthy mice, displayed comparable results. The Si NPs tend to distribute uniformly in the whole body within the first 5–10 min after intravenous injection. The detected activity from $[^{64}\text{Cu}]\text{Cu-Si NP-NOTA}$ and fluorescence from Si NP-IR800 showed that the Si NPs take the renal elimination pathway. Majority of the radioactive NPs ($> 90\%$) are eliminated from the body within 30 min of administration and the rest is gradually cleared over time with no liver accumulation. The *ex vivo* fluorescence analysis of Si NP-IR800 2 h *p.i.* exhibited the maximum fluorescence signals in the urine followed by kidneys. Urine analysis of Si NP-IR800

confirms the covalent attachment of the ligand to the Si NPs, which is stable even within the complex biological system.

Development of bimodal Si NPs for PET-OI:

Using mono-labeled Si NPs as a precursor, efforts were made towards the synthesis of bimodal Si NP-IR800-NOTA. The stepwise synthesis method was adapted for a reproducible synthesis after several optimizations. The characterizations were performed by adapting combined methodologies (radiolabeling to determine the chelator; fluorescence studies for fluorophores).

Si NP-IR800-NOTA

The cell studies of Si NP-IR800-NOTA exhibited extremely low cytotoxicity and thus, a high biocompatibility. The small animal experiment for complementary OI and PET distribution studies of Si NP-IR800-NOTA and its ^{64}Cu -labeled analogue delivered promising results. The ultrasized, nearly neutral Si NP conjugates portrayed excellent pharmacological properties with no accumulation in the body organs. The mode of elimination of the Si NP conjugate was renal. The full body PET and fluorescence scans infer the conjugation of the labeling agents (IR 800 and NOTA) to a single Si NP core with covalent bonds. The urine analysis of the bimodal Si NPs displayed no cleavage of the radiolabel and fluorophore, thus, confirming the stability of the dual-labeled system. Another interesting data observed was the similar pharmacokinetic behavior for both, ^{64}Cu and ^{68}Ga labeled Si NP-IR800-NOTA depicting no influence of the radionuclide on *in vivo* biodistribution and fate of Si NPs. The *ex vivo* studies also showed comparable data to the unimodal imaging agents with majority of the radioactivity and fluorescence expelled into the urine and kidneys at all time-points. A

summary of results obtained by bimodal Si NP-IR800-NOTA is shown in the **Figure** below.

Figure: **A i)** *In vivo* PET scan of $[^{64}\text{Cu}]\text{Cu-Si NP-IR800-NOTA}$ injected in healthy NMRI nu/nu mice 5, 20, 30 and 60 min *p.i.*; **A ii)** Urine analysis by radio-TLC 5 and 60 min *p.i.* **B)** Complementary PET and OI images 1 h and 24 h *p.i.* **D)** Structure of $[^{64}\text{Cu}]\text{Cu-Si NP-IR800-NOTA}$.

As a conclusion, Si NPs bear an extremely versatile and functionalizable platform. The convenient and cost-effective synthesis along with superior pharmacokinetics of the mono- and dual-labeled Si NPs make them distinguished multifunctional candidates for the development of targeting agents. These exceedingly biocompatible imaging probes can build a strong foundation in the field of multimodal molecular imaging due to their unmatched physico-chemical characteristic properties.

The results presented in the thesis, therefore, successfully addressed the scientific goals towards targeted and non-targeted molecular imaging. Bispidines and ultrasmall silicon nanoparticles can serve as potential candidates in the field of cancer diagnostics and imaging in future.

4.2 Outlook

Molecular imaging is an important and envisaged research area with a broad space for further developments. The personalized approach relying on the incorporation of appropriate molecular species to target particular cancer cells is the major aspect of the molecularly targeted therapy. In this direction, the overall outlook of the functionalized bispidines and Si NPs with variable component system looks very promising.

The novel bispidine derivatives developed in the thesis opens up new opportunities to conjugate efficiently to biological vector molecules, capable of targeting variable cancer types and progress towards the evolvement of personalized cancer theranostics. The rigid bicyclic structure with inherent biocompatibility and acquired radiochelation tendency can be exploited comprehensively in the field of multimodal imaging.

The ultrasmall silicon nanoparticles on the other hand, exhibit promising characteristics with easily accessible terminal groups, facile coupling tendency and excellent pharmacokinetics. The synthesized novel dual-labeled Si NPs, in future, can be decorated with cancer tumor specific vectors to target a diverse range of tumors for personalized therapeutics.

In summary, multimodal molecular imaging applying the NPs may have a bright future. The cost-effective, reproducible and convenient synthesis of the bispidine and Si NP-based imaging probes with high biocompatibility can definitely bring a major breakthrough in medical imaging.

Chapter 5 Experimental Section

5.1 Chemicals and Materials

All chemicals and solvents for synthesis and analysis were purchased from Fisher Scientific, Fluka, Merck, VWE, Lumiprobe and Sigma-Aldrich with reagent, analytical grade and used without any further purification. Dry solvents were purchased from Sigma-Aldrich along with corresponding drying agents. Reactions demanding inert conditions were performed under argon or nitrogen atmosphere in flame-dried glassware. Ultrapure water with resistivity 18 MΩ.cm (25°C) obtained from Millipore RiOs/Origin System (Millipore Corporation, USA) was used throughout the synthesis and analysis.

Table 1 Specifications of Chemicals and Materials used.

Chemicals and Materials	Manufacturer	Specifications
Acetonitrile	Fisher	HPLC grade
Ammonium acetate (NH ₄ OAc)	Fluka®	Bioultra
Anatop filters (0.02 µm) filter	GE Healthcare	
3-Bromopropylamine hydrobromide	Sigma-Aldrich	98.0 %
Copper nitrate pentahydrate	Sigma-Aldrich	99.0 %
DBCO-NHS Ester	Lumiprobe	
Dichloromethane	Fisher	HPLC grade
Diethyl ether	Sigma-Aldrich	99.8%
Diethyl-1,3-acetonedicarboxylate	Sigma-Aldrich	96.0%
Dimethyl-1,3-acetonedicarboxylate	Sigma-Aldrich	96.0%
1,4-Dioxane	Sigma-Aldrich	99.8%
Disodium hydrogen phosphate (Na ₂ HPO ₄)	Fluka®	Anhydrous, ≥ 99.5%
Ethanol	Fisher	HPLC grade
Ethyl acetate	Merck	
N,N-Dimethyl Formamide	Sigma-Aldrich	Anhydrous, ≥ 99.8%
N,N-Diisopropylethyl amine (DIPEA)	Sigma-Aldrich	99.0 %
Dimethyl sulfoxide (DMSO)	Sigma-Aldrich	Anhydrous, ≥ 99.7%
dMal-PEG ₄ -NHS	Celares GmbH	98.0 %
N-(3-dimethylaminopropyl)-N'-ethylcarbodiimide hydrochloride (EDC)	Sigma-Aldrich	> 99.0 %

Di- <i>tert</i> -butyl dicarbonate	Sigma-Aldrich	
Ethylene diamine Tetraacetic acid (EDTA)	Fluka®	≥ 99.0 %
Float-A-Lyzer	Spectrum labs	
Formaldehyde	Sigma-Aldrich	37.0%
HBTU (<i>N,N,N',N'</i> -Tetramethyl- <i>O</i> -(1 <i>H</i> -benzotriazol-1-yl)uranium hexafluorophosphate)	Sigma-Aldrich	≥ 98.0%
His60 Ni Column	Clontech	
Hydrochloric acid	Merck	30.0%
2-Iminothiolane Hydrochloride	Sigma-Aldrich	≥ 98.0 %
IRCW 800-NHS ester	Lumiprobe	
S-2-(4-Isothiocyanatobenzyl)-1,4,7-triazacyclononane-1,4,7-triacetic acid (p-SCN-Bn-NOTA)	Macrocyclics	≥ 94.0 %
Maleic anhydride	Merck	≥ 99.0%
Methanol	Fisher	HPLC grade
Microfilter (0.2 µm)	VWR	
2-Morpholino-ethansulfonic acid monohydrate (MES)	Applichem	≥ 99.0%
NAP-5 columns	GE Healthcare	
Octylamine	Sigma-Aldrich	99.0%
PD 10 Sephadex G25	GE Healthcare	
1,3 Phenylene diisothiocyanate	Sigma-Aldrich	97.0%
2-Picolylamine	Sigma-Aldrich	99.0%
Potassium Chloride (KCl)	Fisher	
Potassium hydroxide	Merck	≥ 88.0%
Pyridine	Sigma-Aldrich	> 99.0%
Pyridine-2-aldehyde	Sigma-Aldrich	99.0%
Sephadex G25 fine resin	GE Healthcare	
Sodium borohydride	Sigma/Aldrich	98.5%
Sodium Chloride (NaCl)	Fluka®	Analytical ≥99.5%
Sodium dodecyl Sulfate (SDS)	Sigma-Aldrich	
Sodium Carbonate	Roth	≥ 99.5%
Sodium bicarbonate (NaHCO ₃)	Sigma-Aldrich	≥ 99.5%
Sodium Hydroxide (NaOH)	Sigma-Aldrich	99.9%
Tetrahydrofuran	Fisher	HPLC grade
Triethyl amine (TEA)	Sigma-Aldrich	≥ 99.5%
Tris (2-carboxyethyl)phosphine hydrochloride (TCEP)	Sigma-Aldrich	≥ 98.0%
Trifluoroacetic acid	Fluka	99.0 %

5.2 Analysis, characterization and purification techniques and methods

Dynamic Light Scattering (DLS)

Hydrodynamic size and zeta potential measurement were performed using the dynamic light scattering (DLS) using Zetasizer NanoZS (Malvern Instruments, UK) with an angle detection of 173° at 25°C with a beam wavelength of 632.8 nm. Particle size determination as well as zeta potential measurements were conducted at a concentration of 2 mg/mL in 10 mM phosphate buffer saline (pH 7.4) in disposable plastic cuvettes with a square aperture (Malvern Instruments Ltd.). Prior to each experiment, the solutions were filtered through 0.02 μm or 0.2 μm syringe filters (GE Healthcare, Germany) and equilibrated for 2 mins. The stated value is a mean of three independent measurements. Zeta potential measurements were performed in a universal dip cell (Malvern Instruments Ltd.). The stated values are a mean of three independent measurements using the Smoluchowski model. Data were evaluated with Malvern Zetasizer Software 7.11 version.

Ultraviolet-Visible Spectroscopy (UV-Vis)

UV-Vis spectroscopy was conducted on Specord[®] S50 from Analytik Jena (Germany). The samples were measured on quartz micro cuvettes ($d=1\text{cm}$) with 400-500 μL with a wavelength range from 190-1100 nm.

The standard calibration curves and the molar extinction coefficient of dPG were calculated using the Beer-Lambert equation: $A = \epsilon \times l \times c$ where A is the absorbance at λ_{max} , ϵ denotes the extinction coefficient, c denotes the concentration and l denote the path length of the cuvette (1 cm). For quantum yield experiments, the absorbance was kept between 0.01-0.1. The measurements were set relative to the blank solution (reference). Data were processed using the WinASPECT[®] software.

Fluorescence Spectrometry (FS)

Quantum Yield (QY) and fluorescence measurements were done using a PerkinElmer LS55 using a black cuvette of 100 μL with a light path 10 x 2 mm (105.250, QS). Settings: Split with 10nm and speed record.

Fluorescence emission measurements and quantum yield experiments were done using an LS 55 (230V) luminescence spectrometer from PerkinElmer equipped with a xenon flash lamp source for excitation. Samples were measured in transparent disposable cuvettes of 2 mL with a wavelength range of 200-900 nm and the slit width of 10 nm. FL UV WinLab[®] software was used for Data analysis and acquisition

Fourier Transform Infrared Spectroscopy (FTIR)

Fourier transform infrared (FTIR) spectra were recorded on a Perkin Elmer 1600 Series FTIR spectrometer in the range 4000-400 cm^{-1} , with a resolution of 4 cm^{-1} . Solid samples were prepared as KBr disks. Oils were analyzed using NaCl discs.

Mass spectrometry (MS)

Electron spray Ionization-MS (ESI-MS) was conducted on high-performance compact mass spectrometer 'expression CMS' (Advion, Germany) in either positive or negative mode. The samples 0.1 mg/mL were dissolved in acetonitrile: water (1:1) and a diluted 5 μL sample of 1:100 dilution was injected manually with a flow rate of 0.1 mL min^{-1} . Software Mass express and data express version 3.0 were used for data analysis.

Matrix-assisted laser desorption/ionization (MALDI-TOF) was conducted to determine the PEG conjugation on the sdAbs on a Daltonic Autoflex II TOF/TOF (Bruker, Germany). 3-hydroxypicolinic acid (HPA) was used as the matrix for measurement

Nuclear magnetic resonance spectroscopy (NMR)

^1H -NMR and ^{13}C spectra were recorded on a 400 MHz and 101 MHz Varian Inova 400 spectrometer. Chemical shifts were recorded on a δ scale in parts per million (ppm). The samples were dissolved in deuterated solvents as indicated. Tetramethylsilane (TMS) was used as an internal calibration standard (δ 0 ppm). MeOD (δ 3.31 ppm), CDCl_3 (δ 7.26 ppm) or D_2O (δ 4.79 pm) were used for calibration standard for ^1H spectra. The abbreviations used to describe the resonances are: s (singlet), d (doublet), dd (doublet of doublets), t (triplet), q (quartet), m (multiplet), and br (broad). The data analysis was done using Mestronova software.

Elemental analysis (EA)

It was used to determine the content of C, N, S and H were measured on an Elemental Analyzer EuroEA 3000 from EuroVector (Milan, Italy). The used software was Callidus 5.1. As standards 2,5-(Bis(5-*tert*-butyl-2-benzo-oxazol-2-yl) thiophene (BBOT) and Sulphanilamide were used.

Dialysis

Dialysis was performed on with regenerated cellulose membranes with ultra-pure cellulose ester Float-A-Lyzer membranes from Spectrum Labs, Germany with various molecular weight cut-off (MWCO: 1 kDa, 3.5-5 kDa) against appropriate solvents as indicated in the results and discussion section.

Ultracentrifugation

Amicon® Ultra-0.5 centrifugal filter devices with a molecular weight cutoff of 3K (Amicon® Ultra 3K device, Merck-Millipore) were used to purify single domain antibodies before and after its coupling to the chelators.

Size exclusion chromatography (SEC)

Size Exclusion Chromatography was performed using different resins and columns depending on the compounds and the sample volume. The eluents used were Millipore water or 0.9% NaCl (v/v) as indicated in the experiment section.

System	SEC column	Method	Mobile phase
A	PD-10 Desalting G25 column (Sephadex, G25 medium)	Gravity flow	0.9 % NaCl or water
B	NAP-5 DNA grade column (Sephadex, G25 fine)	Gravity flow	0.9 % NaCl or water

High-Performance Liquid Chromatography (HPLC)

Purification of the bispidine derivatives was performed by the Äkta explorer chromatography system (GE Healthcare, Germany) and Knauer HPLC systems on different columns with detection at 220 and 254 nm. The columns used, injection volume, solvents, flow rates and solvent gradient and UV detection wavelength can be found in the **Table** below.

Äkta explorer uses the Unicorn 5.11 software was applied for peak analysis and Knauer system uses the ChromGate software for data processing and analysis.

Solvent System A = H₂O + 0.1 Vol. % TFA, **B** = CH₃CN + 0.1 Vol. % TFA

Method	HPLC column	Solvent gradient
Analytical HPLC		
A	Phenomenex Aqua C-18 125 Å (250 × 4.6 mm, 5 µm)	0% to 70% B in 20 min 70% to 100% B in 2 min f = 1 mL/min, λ = 220 nm
Semipreparative HPLC		
B	Phenomenex Gemini C-18 110 Å (250 10 mm, 5µm)	i) 30% to 70% B in 30 min 70% to 100% B in 15 min f = 1 mL/min, λ = 220 nm ii) 30% to 50% B in 20 min 50% to 70% B in 10 min f = 1 mL/min, λ = 220 nm

C	Synergi Hydro RP 80 Å (250 x 21,2 mm, 4µm)	60% B to 40% B in 15 min f = 10 mL/min, λ = 220 nm
Preparative HPLC		
D	Phenomenex Gemini C-18 110 Å (250 x 30 mm, 5µm)	i) 30% B to 70% B in 15 min 70% B to 100% B in 15 min f = 10 mL/min, λ = 220 nm ii) 30% B to 70% B in 40 min 70% B to 100% B in 20 min f = 10 mL/min, λ = 220 nm iii) 30% B to 70% B in 70% B to 100% B in f = 10 mL/min, λ = 220 nm

Thin Layer Chromatography (TLC)

Thin layer chromatography was conducted on various stationary phases such as silica gel (reverse phase, RP-TLC) and neutral aluminum oxide (Alox) plates using different mobile phase systems in a vertical chamber for elution. UV lamp (254 nm) and Amersham Typhoon 5 Scanner (300-900 nm) were used for analysis of fluorescence labeled nanoparticles.

5.3 Radionuclide production (^{68}Ga and ^{64}Cu)

^{68}Ga was eluted from a $^{68}\text{Ge}/^{68}\text{Ga}$ generator from iThemba LABS (Faure, South Africa) in as $[^{68}\text{Ga}]\text{GaCl}_3$ by using 0.6-0.8 M HCl. The concentration of the eluate is usually in a range of 2-4 MBq /mL depending on the age of the generator. The $[^{68}\text{Ga}]\text{GaCl}_3$ was purified by ion exchange chromatography and then concentrated.

^{64}Cu was produced as $[^{64}\text{Cu}]\text{CuCl}_2$ by the cyclotron “Cyclone 18/9” (IBA molecular) at the Helmholtz-Zentrum Dresden–Rossendorf by a $^{64}\text{Ni}(p,n)^{64}\text{Cu}$ nuclear reaction. The yields of the nuclear reaction were between 3.6–5.2 GBq (EOB) with molar activities between 150–250 GBq/µmol Cu [64].

Radio-Thin Layer Chromatography (Radio-TLC)

Radiochemical analysis of the yields and kinetics of the experiments was determined using different stationary phase and eluent systems. iTLC plates were purchased from Agilent Technologies. The air-dried TLC strips are exposed to a high-resolution phosphorimaging plate (GE Healthcare). The exposed plates are scanned with an Amersham Typhoon 5 Scanner (GE Healthcare) or read by a radioluminography laser scanner BAS-1800II (Raytest). Aida Image Analyzer version 4.0 software was used for

data analysis and conversion of data to ASCII before graph-plotting in Origin. The TLC systems used are:

Method	TLC plate	Mobile phase
I	Reverse phase silica gel (RP C-18)	2M NH ₄ OAc : MeOH (1:1, v:v)
II	Neutral aluminium oxide (Alox)	2M NH ₄ OAc : MeOH (1:1, v:v)
III	iTLC - SG	2M NH ₄ OAc : MeOH (1:1, v:v)

Sample	Method I	Method II	Method III	Method IV
[⁶⁴ Cu]CuCl ₂	0.0	0.0	1.0	0.0
[⁶⁴ Cu]Cu-Si NP conjugates	0.0	0.0	0.0	0.0
[⁶⁴ Cu]Cu-NOTA	0.0	~ 0.5	~ 0.7	1.0
[⁶⁴ Cu]Cu-EDTA	1.0	0.0	1.0	1.0
[⁶⁴ Cu]Cu-Bispidine	0.0	~ 0.6	~ 0.5	~ 0.8

Radio-Size Exclusion Chromatography (Radio-SEC)

Radio-SEC was performed using the high performances columns used for the analysis of the non-radioactive compounds on a Knauer HPLC system fitted with a UV detector and a Gabi radio-detector (Raytest). ChromeGate 3.3.2 software was used for peak analysis and interpretation.

Radio High Performance Liquid Chromatography (Radio-HPLC)

Same columns and HPLC systems as for non-radioactive samples were used but connected to a radioactive detector (Raytest, Ramona Start, and Software Chromgate 2.8)

Design Software methods

All graphs were plot in OriginLab 9.0. Chemical Structures were created using ChemBioDraw Ultra 12.0. Figures were done using Adobe Photoshop CS2, Adobe ImageReady CS2 or CorelDrawX.

5.4 Syntheses

5.4.1 Synthesis of bispidines derivatives (B1-B11), intermediates (B1', B3', B8', B9', B11') and bifunctional linkers (L1, L2)

Dimethyl-4-oxo-2,6-di(pyridin-2-yl)-1-(pyridin-2-ylmethyl)piperidine-3,5-dicarboxylate (B1')

The piperidone precursor (**B1'**) was obtained by drop-wise addition of pyridine-2-aldehyde (9.56 mL, 100 mmol) and picolylamine (5.1 mL, 50 mmol) to an ice-cold solution of dimethyl acetonedicarboxylate (7.2 mL, 50 mmol) dissolved in methanol (30 mL). After 30 min, the solidified orange solution was stored in the freezer overnight, washed with cold ethanol, and recrystallized from ethanol. [282].

Yield: 18.62 g, 81%

ESI MS (m/z): $[M+H]^+$ 461.3

Dimethyl-9-oxo-2,4-di(pyridin-2-yl)-3,7-bis(pyridin-2-ylmethyl)-3,7-diazabicyclo[3.3.1]nonane-1,5-dicarboxylate (B1)

To a solution of **B1'** (15.00 g, 32.6 mmol) in 25 mL THF, 2(aminomethyl) pyridine (4.23 g, 39.1 mmol) and formaldehyde (37%, 6.4 mL, 78.3 mmol) were added. This solution run under reflux for 4 h until a deep brown solution was formed. The solvent was removed under low pressure and the remaining dark brown solid was recrystallized from ethanol [282].

Yield: 4.81 g, 25%

ESI MS (m/z): $[M+TFA]^+$ 611.6

¹H NMR: δ 8.59 (ddd, J = 4.9, 1.9, 0.9 Hz, 1H), 8.44 – 8.36 (m, 2H), 7.86 (d, J = 7.9 Hz, 1H), 7.65 (td, J = 7.7, 1.8 Hz, 1H), 7.49 – 7.39 (m, 2H), 7.37 (td, J = 7.6, 1.9 Hz, 1H), 7.20 (ddd, J = 7.6, 4.9, 1.2 Hz, 1H), 7.04 (ddd, J = 7.5, 4.8, 1.2 Hz, 2H), 6.95 (ddd, J = 7.6, 4.9, 1.2 Hz, 1H), 6.75 (dt, J = 7.8, 1.1 Hz, 1H), 5.32 (s, 2H), 3.67 (d, J = 2.5 Hz, 6H), 3.62 (s, 2H), 3.47 – 3.40 (m, 1H), 2.87 (d, J = 11.9 Hz, 1H)

¹³C NMR: δ 203.39, 168.90, 158.08, 157.17 (d, J = 27.0 Hz), 149.29, 149.02 (d, J = 2.1 Hz), 136.27, 135.68 (d, J = 24.1 Hz), 124.92, 124.22, 123.80, 122.65, 122.28, 121.59, 70.48, 63.43, 62.98, 58.60, 56.72, 52.45

Dimethyl-9-hydroxy-2,4-di(pyridin-2-yl)-3,7-bis(pyridin-2-ylmethyl)-3,7-diazabicyclo [3.3.1]-nonane-1,5-dicarboxylate (B2)

B1 (1.5 g, 2.5 mmol) was dissolved in a 3:1 mixture of 1,4-dioxane/water (170 mL) and cooled to -5°C with a ice/acetone bath. Sodium borohydride (124.5 mg, 3.3 mmol) suspended in an ice/water mixture (20 mL) was added dropwise over 2 h. The reaction was run for 12 h. The pH of the solution was adjusted to 3 with 1 M hydrochloric acid and then to pH 8 by addition of 1 M sodium hydroxide solution. The reaction mixture was extracted with DCM. The organic layer was dried over sodium sulfate. The solvent was removed under reduced pressure and the crude product was purified by preparative RP-HPLC. B3 was obtained as a white solid [76].

Yield: 88 mg, 59 %

ESI MS (m/z): $[M+H]^+$ 596.2

¹H NMR: δ 8.66 – 8.58 (m, 1H), 8.45 – 8.34 (m, 2H), 7.87 (d, J = 7.9 Hz, 1H), 7.74 – 7.53 (m, 1H), 7.48 – 7.27 (m, 3H), 7.27 – 7.15 (m, 1H), 7.02 (ddd, J = 7.3, 4.9, 1.2 Hz, 2H), 6.98 – 6.84 (m, 1H), 6.68 (d, J = 7.7 Hz, 1H), 4.76 (s, 1H), 4.67 (d, J = 5.8 Hz, 1H), 4.17 – 3.96 (m, 1H), 3.65 (s, 1H), 3.54 (d, J = 10.9 Hz, 6H), 2.70 – 2.49 (m, 3H)

¹³C NMR: δ 172.67, 159.42, 157.74, 156.62, 149.28 (d, J = 18.0 Hz), 148.38 (d, J = 14.9 Hz), 136.09, 135.53 (d, J = 16.7 Hz), 124.73, 124.39, 124.17,

122.20 (d, $J = 18.7$ Hz), 121.59 (d, $J = 10.4$ Hz), 70.88, 65.17, 57.64, 52.98, 52.03, 49.21, 30.89

FTIR : > 3000 (O-H, N-H, C-H), 1720 (C=O), 1670 (C=N), 1210 (C-O)

***Tert*-butyl (3-bromopropyl)carbamate (L1)**

$\text{C}_8\text{H}_{16}\text{BrNO}_2$
M = 238.1 g/mol

To a solution of 3-bromopropylamine hydrobromide (2.0 g, 9.1 mmol) in DCM (30 mL), TEA (5 mL, 36.5 mmol) was added and the reaction solution was stirred at room temperature for 30 min before di-*tert*-butyl-dicarbonate (2.4 g, 10.9 mmol) was added. After 12 h of continuous stirring at room temperature under nitrogen atmosphere, 20 mL ethyl acetate and 20 mL water were added to the solution. The organic phase was separated, washed with brine and the solvent was removed under reduced pressure. The product L1 was obtained as a white solid by after purification by column chromatography (silica gel, ethyl acetate:hexane = 1:5) [283].

Yield: 1.52 g, 70%

^1H NMR: δ 3.37 (t, $J = 6.5$ Hz, 2H), 3.19 (q, $J = 4.7, 4.0$ Hz, 2H), 1.97 (p, $J = 6.6$ Hz, 2H), 1.37 (s, 9H)

^{13}C NMR: δ 155.99, 39.69, 30.74, 28.71 – 27.90 (m)

Dimethyl-9-(3-((*tert*-butoxycarbonyl)amino)propoxy)-2,4-di(pyridin-2-yl)-3,7-bis(pyridin-2-ylmethyl)-3,7-diazabicyclo[3.3.1]nonane-1,5-dicarboxylate (B3')

$\text{C}_{41}\text{H}_{49}\text{N}_7\text{O}_7$
M = 751.8 g/mol

Sodium hydride (311.2 mg, 1.3 mmol) was added to a solution of **B2** (300 mg, 504 μmol) in dry THF (50 mL) and the resulting mixture was stirred for 1.5 h at 50 °C. *Tert*-butyl (3-bromopropyl)carbamate (238.1 mg, 1.0 mmol) was then added to the solution and stirred for another 20 h at 50 °C. The reaction was quenched by adding saturated sodium hydrogen carbonate solution (50 mL, pH = 8). The aqueous layer was extracted with DCM and ethyl acetate. The organic fractions were combined, washed with brine and dried over sodium sulfate. Finally, the organic solvent was removed to obtain crude **B3'**. The product was used for the next step *in situ* [284, 285].

Yield: 131 mg, 35%

ESI MS (m/z): $[M+H]^+$ 751.8

^1H NMR: δ 8.66 (d, J = 4.9 Hz, 2H), 8.44 (dd, J = 5.0, 1.7 Hz, 2H), 8.41 – 8.35 (m, 4H), 7.89 (d, J = 7.9 Hz, 3H), 7.68 (td, J = 7.6, 1.8 Hz, 2H), 7.56 – 7.45 (m, 2H), 7.37 (dtd, J = 9.8, 7.6, 1.8 Hz, 8H), 7.28 – 7.21 (m, 3H), 6.99 (ddd, J = 15.3, 7.5, 5.0 Hz, 7H), 6.68 (d, J = 7.7 Hz, 2H), 5.26 (s, 2H), 4.90 (d, J = 6.6 Hz, 2H), 4.80 (s, 3H), 4.70 (s, 1H), 4.36 (s, 2H), 3.70 (dq, J = 6.5, 3.3, 2.9 Hz, 2H), 3.58 (s, 9H), 3.55 – 3.45 (m, 9H), 3.41 (dd, J = 14.6, 8.1 Hz, 8H), 3.23 (q, J = 6.3 Hz, 5H), 2.99 (q, J = 6.6, 6.0 Hz, 4H), 2.55 – 2.43 (m, 7H), 2.42 (s, 1H), 2.06 – 1.87 (m, 6H), 1.51 (h, J = 6.3, 5.5 Hz, 4H), 1.24 – 1.20 (m, 3H), 0.86 – 0.77 (m, 3H)

Dimethyl-9-(3-aminopropoxy)-2,4-di(pyridin-2-yl)-3,7-bis(pyridin-2-ylmethyl)-3,7-diazabicyclo[3.3.1]nonane-1,5-dicarboxylate (B3)

The Boc-protected amine (100 mg, 153 μmol) was dissolved in DCM (3 mL), to which trifluoroacetic acid (3 mL) was added. The resulting reaction mixture was stirred for 24 h at room temperature. After addition of chloroform (10 mL), the solvent was removed under reduced pressure. The product **B3** was purified by RP-HPLC and obtained as a yellow-brown oil [285].

Yield: 86.5 mg, 100 %

RP-HPLC: R_t = 11 min (Method D/i)

ESI MS (m/z): $[M+H]^+$ 651.4

^1H NMR: δ 9.12 (ddd, J = 4.9, 1.8, 0.9 Hz, 1H), 8.79 (dd, J = 6.0, 1.6 Hz, 1H), (CD₃CN) 8.15 – 8.05 (m, 3H), 8.05 – 7.97 (m, 2H), 7.82 – 7.75 (m, 3H), 7.75 – 7.67 (m, 2H), 7.63 (dt, J = 7.8, 1.0 Hz, 2H), 7.31 (dt, J = 7.8, 1.1 Hz, 2H), 7.26 – 7.18 (m, 3H), 7.11 (d, J = 8.1 Hz, 2H), 6.97 (s, 1H)

^{13}C NMR: δ 167.45, 125.12 – 150.90, 125.33, 80.74, 72.64, 65.12, 55.01, 53.20, (CD₃CN) 52.14

FTIR: > 3000 (N-H, C-H), 1727 (C=O), 1670 (C=N), 1200 (C-O), 1126 (N-H)

Dimethyl-9-(3-(3-(4-isothiocyanatophenyl)thioureido)propoxy)-2,4-di(pyridin-2-yl)-3,7-bis(pyridin-2-ylmethyl)-3,7-diazabicyclo[3.3.1]nonane-1,5-dicarboxylate (B4)

B3 (250 mg, 0.307 mmol) was dissolved in DCM (25 mL). p-Phenylene diisothiocyanate (80 mg, 416 μmol) alongwith TEA (1 mL, 7.1 mmol) was added to this solution. The reaction mixture turned yellow in 5–10 min. The reaction was stirred at room temperature for 16 h. The reaction solvent was evaporated under reduced pressure. The p-phenylene diisothiocyanate was extracted from the mixture by dissolving it in diethyl ether. The crude solid **B4** was then purified by RP-HPLC.

Yield: 194 mg, ~60%

RP-HPLC: $R_t = 19 \text{ min}$ (Method D/i)

ESI MS (m/z): $[M+H]^+ 843.3$

^1H NMR: δ 8.90 (ddd, $J = 4.8, 1.8, 0.9 \text{ Hz}$, 1H), 8.63 (d, $J = 8.1 \text{ Hz}$, 1H), 8.23 – 8.12 (m, 2H), 7.93 (ddd, $J = 4.9, 1.8, 0.8 \text{ Hz}$, 2H), 7.82 (td, $J = 7.7, 1.8 \text{ Hz}$, 1H), 7.71 – 7.53 (m, 5H), 7.45 (ddd, $J = 7.4, 5.8, 1.2 \text{ Hz}$, 2H), 7.27 – 7.17 (m, 7H), 7.15 – 7.05 (m, 4H), 4.69 – 4.60 (m, 5H), 4.01 (d, $J = 12.5 \text{ Hz}$, 2H), 3.92 (d, $J = 12.7 \text{ Hz}$, 2H), 3.80 (s, 2H), 3.66 (t, $J = 5.3 \text{ Hz}$, 4H), 3.53 (s, 6H), 1.84 (p, $J = 5.6 \text{ Hz}$, 2H)

^{13}C NMR: δ 149.34, 138.00, 126.02, 125.76, 123.87, 28.60

FTIR: > 3000 (N-H, C-H), 2140 (N=C=S), 1727–1675 (C=O), 1200 (C-O), 1140 (N-H)

EA: C 53.75%, N 11.14%, H 4.61%, S 5.46%

Dimethyl-9-(5,8(8-dibenzocyclooctyne)diamido)octoxy)-2,4-di(pyridin-2-yl)-3,7-bis(pyridin-2-ylmethyl)-3,7-diazabicyclo[3.3.1]nonane-1,5-dicarboxylate (B5)

B3 (2.5 g, 2.86 mmol) and TEA (2.5 mL, 17.8 mmol) were dissolved in DCM (40 mL). DBCO-NHS ester (1.0 g, 2.5 mmol) dissolved in 10 mL DCM was added to this solution. The reaction mixture was stirred for 2 h at room temperature. The product **B5** was concentrated under reduced pressure, and purified by RP-HPLC [286].

Yield: 1.63 g, 56 %

RP-HPLC: $R_t = 22 \text{ min}$ (Method D/i)

ESI MS (m/z): $[M+H]^+ 939.1$

^1H NMR: δ 8.89 – 8.82 (m, 1H), 8.49 (s, 1H), 8.26 (s, 1H), 8.23 (d, $J = 8.1 \text{ Hz}$, 1H), 8.11 (d, $J = 5.2 \text{ Hz}$, 2H), 8.02 (d, $J = 8.9 \text{ Hz}$, 2H), 7.97 – 7.82 (m, 2H), 7.70 – 7.60 (m, 4H), 7.60 – 7.47 (m, 4H), 7.47 – 7.40 (m, 2H), 7.40 – 7.32 (m, 5H), 7.29 (d, $J = 8.1 \text{ Hz}$, 6H), 7.14 – 7.04 (m, 3H), 5.14 (s, 1H), 5.12 – 5.02 (m, 1H), 5.00 (s, 1H), 4.73 – 4.64 (m, 2H), 4.62 (s, 1H), 4.07 (d, $J = 9.6 \text{ Hz}$, 3H), 4.00 (d, $J = 13.3 \text{ Hz}$, 2H), 3.90 (d, $J = 10.4 \text{ Hz}$, 3H), 3.68 – 3.59 (m, 3H), 3.50 (t, $J = 4.1 \text{ Hz}$, 8H), 3.27 (d, $J = 9.1 \text{ Hz}$, 1H), 3.08 (t, $J = 6.0 \text{ Hz}$, 3H), 2.80 – 2.66 (m, 8H), 2.46 (dt, $J = 14.6, 7.3 \text{ Hz}$, 5H), 2.29 (ddt, $J = 21.6, 14.4, 7.1 \text{ Hz}$, 4H), 2.15 (s, 1H), 1.95 (dt, $J = 16.6, 7.1 \text{ Hz}$, 2H), 1.58 (s, 1H), 1.52 (s, 1H), 1.24 (s, 2H), 0.82 (s, 1H).

FTIR: > 3000 (N-H, C-H), 1740-1600 (C=O), 1200 (C-O), 1140 (N-H)

Dimethyl-9-(3-(4-mercaptobutanimidamido)propoxy)-2,4-di(pyridin-2-yl)-3,7-bis(pyridin-2-ylmethyl)-3,7-diazabicyclo[3.3.1]nonane-1,5-dicarboxylate (B6)

B3 (1.2 g, 1.8 mmol) was dissolved in 10 mL acetonitrile and methanol (4 mL) to which iminothiolane hydrochloride (250 mg, 1.8 mmol) and TEA (280 μ L, 1.9 mmol) were added. The solution was stirred at room temperature for 4 h and the solvent was removed under reduced pressure. The product **B6** was purified by RP-HPLC and obtained as a white solid. It was stored under inert conditions at -20 °C [287].

Yield: 50%

RP-HPLC: $R_t = 15 \text{ min}$ (Method D/ii)

ESI MS (m/z): $[M+H-17]^+ 736.2$

^1H NMR: δ 8.07 (d, $J = 4.8 \text{ Hz}$, 3H), 7.97 (t, $J = 7.6 \text{ Hz}$, 3H), 7.78 – 7.63 (m, 6H), 7.62 (s, 2H), 7.29 (d, $J = 7.6 \text{ Hz}$, 2H), 7.25 (s, 6H), 7.18 – 7.10 (m, 4H), 4.87 (d, $J = 9.0 \text{ Hz}$, 5H), 4.78 (s, 2H), 4.01 (d, $J = 12.9 \text{ Hz}$, 3H), 3.92 (d, $J = 10.4 \text{ Hz}$, 5H), 3.85 (d, $J = 10.7 \text{ Hz}$, 2H), 3.71 (d, $J = 15.5 \text{ Hz}$, 4H), 3.62 (s, 6H), 3.60 – 3.54 (m, 4H), 3.53 (d, $J = 6.6 \text{ Hz}$, 5H), 3.38 (dt, $J = 14.8, 7.0 \text{ Hz}$, 4H), 3.29 (s, 2H), 3.20 – 3.13 (m, 3H), 2.41 (dt, $J = 13.8, 7.0 \text{ Hz}$, 10H), 2.17 (d, $J = 0.8 \text{ Hz}$, 44H), 2.10 (d, $J = 0.9 \text{ Hz}$, 22H), 2.08 (s, 10H), 2.00 (d, $J = 0.9 \text{ Hz}$, 7H), 1.34 (t, $J = 7.3 \text{ Hz}$, 4H), 1.29 – 1.21 (m, 3H)

Dimethyl-9-(3-(2-mercaptoacetamido)propoxy)-2,4-di(pyridin-2-yl)-3,7-bis(pyridin-2-ylmethyl)-3,7-diazabicyclo[3.3.1]nonane-1,5-dicarboxylate (B7)

B7 is formed in a two-step reaction. In the first reaction step, **B3** (1.3 g, 2 mmol), S-Acetylthioglycolic acid NHS ester (4.6 g, 2 mmol) and trimethylamine (50 μ L, 300 μ mol)

were dissolved in acetonitrile (10 mL). The solution was stirred at room temperature for 2.5 h. The solvent was then removed under reduced pressure. The second step of the reaction was performed *in situ*.

In the second step, the thioester formed was hydrolyzed by suspending in hydroxylamine hydrochloride, ethylenediaminetetraacetic and sodium hydrogen carbonate in a mixture of water and methanol (10 mL, 1:1 v/v). The pH was adjusted to pH 7 and the suspension was stirred at room temperature for 3 h. The product **B7** was extracted with DCM. The organic phases were combined and dried with anhydrous sodium sulfate and the solvent was removed under reduced pressure. The product was purified *via* RP-HPLC as a colorless solid. **B7** was stored under inert gas at -20 °C [287].

Yield: 0.94 g, 68%

ESI MS (*m/z*): [M+H]⁺ 725.3

Tert-butyl (4-(2-bromoacetamido)phenyl)carbamate (L2)

Bromoacetate succinimide ester (39 mg, 0.12 mmol) was dissolved in dry acetonitrile (4 mL) and a solution of p-amino-Boc-aniline (36 mg, 160 μmol) in dry acetonitrile (4 mL) added. The mixture was stirred under nitrogen atmosphere for 24 h, whereupon a solid precipitated. The solvent was removed under vacuum and the residue was dissolved in chloroform (10 mL) and washed thrice with water (10 mL each). The organic phase was separated, dried over anhydrous sodium sulfate, filtered and evaporated to yield the product **L1** as a powder [196].

Yield: 46 mg, 90%

¹H NMR: δ 8.17 (s, 1H), 7.42 (d, *J* = 6.7 Hz, 1H), 7.41 – 7.17 (m, 4H), 6.60 (s, 1H), 3.99 (s, 1H), 2.64 (s, 2H), 1.49 (s, 9H), 1.47 – 1.39 (m, 1H) (CDCl₃)

¹³C NMR: δ 115.79, 28.35, 25.31

Dimethyl-9-(3-((2-((4-((tert-butoxycarbonyl)amino)phenyl)amino)-2-oxoethyl)amino)propoxy)-2,4-di(pyridin-2-yl)-3,7-bis(pyridin-2-ylmethyl)-3,7-diazabicyclo[3.3.1]nonane-1,5-dicarboxylate (B8')

L1 (10 mg, 0.03 mmol) was dissolved in dry acetonitrile (5 mL) and a solution of **B3** (19.5 mg, 0.03 mmol) in dry acetonitrile (5 mL) was added, followed by K_2CO_3 (17 mg, 0.13 mmol) and KI (4 mg, 0.02 mmol). The mixture was stirred at room temperature for 1 h and then refluxed for 24 h. After cooling to room temperature, the solution was filtered and the solvent evaporated to yield the product **B8'** as a brown oil [196].

Yield: 12 mg, 42%

ESI MS (m/z): $[M+H]^+$ 899.3

Dimethyl-9-(3-((2-((4-aminophenyl)amino)-2-oxoethyl)amino)propoxy)-2,4-di(pyridin-2-yl)-3,7-bis(pyridin-2-ylmethyl)-3,7-diazabicyclo[3.3.1]nonane-1,5-dicarboxylate (B8**)**

B8' (10 mg, 0.01 mmol) was dissolved in a 1:1 (v/v) mixture of chloroform and TFA (2 mL). The mixture was stirred for 4 h and then the solvent was evaporated to obtain the crude product **B8** as dark brown red oil [196, 288].

Yield: ~2 mg, 19%

ESI MS (m/z): $[M+H]^+$ 799.5

Dimethyl-9-((1-(9H-fluoren-9-yl)-3,6,9,12-tetraoxo-2-oxa-4,7,10,13-tetraaza hexadecan-16-yl)oxy)-2,4-di(pyridin-2-yl)-3,7-bis(pyridin-2-ylmethyl)-3,7-diazabicyclo [3.3.1]-nonane-1,5-dicarboxylate (B9'**)**

Fmoc-(Gly)₃-OH (22.9 mg, 56 μ mol) was dissolved in 560 μ l anhydrous DMF solvent. To this solution, PyBOP (29.4 mg, 56 μ mol), HOBT (7.57 mg, 56 μ mol) and DIPEA (19 μ l, 0.11 mmol) were added to generate the active ester. The solution was stirred at room temperature for 5 min when it turns brown in color. At this point, **B3** (48.5 mg, 55 μ mol) dissolved in 560 μ l anhydrous DMF was added to the active ester solution. The mixture was stirred overnight at room temperature under inert (Argon) conditions. The solvent was then removed under low pressure and the crude product was purified by RP-HPLC [232].

Yield: 16,8 mg, 30%

RP-HPLC: $R_t = 11 \text{ min}$ (Method C)

ESI MS (m/z): $[M+H]^+ 1045.4$

¹H NMR: δ 9.00 – 8.94 (m, 1H), 8.50 (d, $J = 5.5 \text{ Hz}$, 1H), 8.13 – 8.06 (m, 1H), (CD₃CN) 7.99 (td, $J = 7.7, 1.7 \text{ Hz}$, 1H), 7.86 (dd, $J = 17.0, 7.8 \text{ Hz}$, 2H), 7.73 – 7.60 (m, 4H), 7.44 (dt, $J = 10.8, 6.4 \text{ Hz}$, 2H), 7.36 (d, $J = 7.1 \text{ Hz}$, 1H), 7.35 – 7.29 (m, 1H), 7.27 (s, 1H), 7.20 (dd, $J = 7.9, 3.9 \text{ Hz}$, 3H), 6.73 (t, $J = 5.6 \text{ Hz}$, 1H), 6.34 (s, 1H), 4.90 (s, 1H), 4.63 (s, 1H), 4.56 (s, 1H), 4.33 (d, $J = 7.0 \text{ Hz}$, 2H), 4.24 (t, $J = 7.1 \text{ Hz}$, 1H), 3.97 (d, $J = 12.6 \text{ Hz}$, 2H), 3.81 – 3.63 (m, 5H), 3.59 (d, $J = 11.1 \text{ Hz}$, 6H), 3.52 (t, $J = 6.0 \text{ Hz}$, 2H), 3.10 (d, $J = 13.1 \text{ Hz}$, 2H), 1.85 – 1.77 (m, 1H), 1.59 (p, $J = 6.4 \text{ Hz}$, 2H), 1.09 (d, $J = 6.3 \text{ Hz}$, 1H)

Dimethyl-9-(3-(2-(2-(2-aminoacetamido)acetamido)acetamido)propoxy)-2,4-di(pyridin-2-yl)-3,7-bis(pyridin-2-ylmethyl)-3,7-diazabicyclo[3.3.1]nonane-1,5-dicarboxylate (B9)

B9' (2.3 mg, 2.2 mmol) was dissolved in 500 μL 20 % Piperidine/DMF solution and stirred at room temperature for 20 min at room temperature. The crude product was then lyophilized and purified by RP-HPLC [232].

Yield: 1.2 mg, 66%

RP-HPLC: $R_t = 15 \text{ min}$ (Method C)

ESI MS (m/z): $[M+H]^+ 822.6$

^1H NMR: δ 7.80 – 7.65 (m, 4H), 3.96 – 3.81 (m, 4H), 3.75 – 3.69 (m, 4H), 3.65 (CD₃CN) (d, $J = 1.1 \text{ Hz}$, 6H), 3.59 (t, $J = 6.0 \text{ Hz}$, 2H), 3.15 (q, $J = 6.6 \text{ Hz}$, 3H), 1.99 – 1.93 (m, 72H), 1.65 (p, $J = 6.4 \text{ Hz}$, 2H)

9-oxo-2,4-di(pyridin-2-yl)-3,7-bis(pyridin-2-ylmethyl)-3,7-diazabicyclo [3.3.1]-nonane-1,5-dicarboxylic acid (B10)

A solution of **B2** (200 mg, 0.336 mmol) was prepared by dissolving it in 6 mL of anhydrous THF. Cesium hydroxide (565 mg, 3.36 mmol) was added to this solution under ice cooling followed by 34 μL water. The reaction mixture was stirred for 24 h at room temperature. A milky white emulsion was obtained towards the end of reaction. The emulsion was acidified by dilute HCl to pH 6. The solvent was then removed under low pressure and the product was purified by RP-HPLC. The product was obtained as a white crystalline solid [76].

Yield: 159 mg, 86%

RP-HPLC: $R_t = 11 \text{ min}$ (Method D/ii)

ESI MS (m/z): $[M+H]^+ 566.8$

^1H NMR: δ 8.93 (d, J = 5.1 Hz, 1H), 8.65 (d, J = 5.9 Hz, 1H), 8.19 – 8.10 (m, 1H), 8.02 (t, J = 7.9 Hz, 1H), 7.95 (d, J = 4.9 Hz, 1H), 7.72 (ddd, J = 25.7, 17.4, 7.1 Hz, 4H), 7.38 (d, J = 7.8 Hz, 1H), 7.16 (t, J = 6.4 Hz, 1H), 7.05 (d, J = 8.0 Hz, 1H), 4.93 (s, 1H), 4.87 (s, 1H), 4.60 (s, 2H), 3.75 (s, 1H), 3.70 (d, J = 12.8 Hz, 1H), 3.50 (d, J = 12.9 Hz, 1H)

^{13}C NMR (D₂O) δ 171.52, 162.87, 162.52, 149.23, 148.62, 146.35, 140.53, 127.57, 126.45, 125.89, 125.18 (d, J = 14.8 Hz), 117.64, 114.74, 71.12, 69.87, 60.58, 56.05, 52.70, 48.98

Diethyl-4-oxo-2,6-di(pyridin-2-yl)-1-(pyridin-2-ylmethyl)piperidine-3,5-dicarboxylate (B11')

$\text{C}_{27}\text{H}_{28}\text{N}_4\text{O}_5$
 $M = 488.5 \text{ g/mol}$

B11' was obtained by drop-wise addition of pyridine-2-aldehyde (9.56 mL, 100 mmol) and picolylamine (5.10 mL, 50 mmol) to an ice-cold solution of diethyl acetonedicarboxylate (9.08 mL, 50 mmol) in methanol (30 mL). After 30 min, the brown solution was extracted with chloroform/water. The organic layer was removed under low pressure and purified by ethanol crystallization [282].

Yield: 16.7 mg, 79%

ESI MS (m/z): $[\text{M}+\text{H}]^+$ 489.1

Diethyl-9-oxo-2,4-di(pyridin-2-yl)-3,7-bis(pyridin-2-ylmethyl)-3,7-diazabicyclo[3.3.1]nonane-1,5-dicarboxylate (B11)

$\text{C}_{35}\text{H}_{36}\text{N}_6\text{O}_5$
 $M = 620.7 \text{ g/mol}$

To a solution of **B11'** (16 g, 32.6 mmol) in 25 mL THF, 2(aminomethyl) pyridine (4.23 g, 39.1 mmol) and formaldehyde (37%, 6.4 mL, 78.3 mmol) were added. This solution run under reflux for 4 h till a deep brown solution was formed. The solvent was removed

under low pressure and the remaining dark brown solid was recrystallized from ethanol [282].

Yield:

ESI MS (m/z): $[M+H]^+$ 621.2

5.4.2 Synthesis of bispidines-based bioconjugates B4-EG-TATE and B5-sdAb-7C12

B4-EG-TATE-Boc

TATE Peptide (1.70 mg, 1 μ mol) and B4 (0.98 mg, 1.1 μ mol) were dissolved in 2 mL anhydrous ACN. 200 μ DIPEA was added dropwise to this solution. The reaction mixture was stirred overnight at room temperature. The product was then lyophilized and further purified by RP-HPLC. The identification of the Boc-protected B4-EG-TATE product was done by MALDI-TOF spectrometry [289].

Yield: 0.9 mg, 37%

RP-HPLC: 27 min (Method B)

MALDI-TOF (m/z): $[M+H]^+$ 2407.1

B4-EG-TATE

Boc-protected B4-EG-TATE (0.9 mg, 0.37 μ mol) was dissolved in 85 μ L TFA and 55 μ L H₂O. 5 μ L TIPS and 0.1 mg was added as scavengers to the reaction mixture. The

reaction was stirred overnight at room temperature. The product B4-EG-TATE was lyophilized and purified by RP-HPLC. The identification was performed by MALDI-TOF [185, 289, 290].

Yield: 0.25 mg, 33%
 RP-HPLC: 12.5 min (Method B)
 MALDI-TOF (m/z): $[M+H]^+$ 2038.5

B5-sdAb-7C12

Strain-promoted azide-alkyne cycloaddition

Small-scale reactions were set up in 50 μ L with variable molar ratios of azide-functionalized sdAb and **B5**. The optimal conditions were upscaled and the reaction mixture was composed of 275 nmol azide-functionalized sdAb and 1375 nmol **B5**. Reactions were incubated at 25°C for up to 4 h with gentle shaking. The unconjugated **B5** was removed by SEC using Zeba Spin Desalting Columns (7K MWCO, Thermo Scientific) with elution in MES/NaOH buffer (100 mM, 150 mM NaCl, pH 6.0). The purified conjugate was sterile filtered using Whatman Puradisc FP 30 cellulose acetate syringe filter units with a pore size of 0.2 μ m (GE Healthcare Life Sciences) and stored at 4°C.

5.4.3 Synthesis and modification of Si NPs by hydrothermal method

Synthesis of Si NPs by hydrothermal method

Amine-terminated Si NPs were prepared by slight modification of a reported protocol. Trisodium citrate (0.8211 g, mmol) was dissolved in Argon-saturated Millipore water (20 mL). Argon was bubbled into the citrate solution for 15 min. 3-Aminopropyl-trimethoxysilane (APTMS) (5 mL) was then added to the solution and stirred for another

15 min at room temperature. The reaction mixture was transferred to a Teflon vessel in an Argon environment. The Si NP precursor solution was heated up to 160 °C for 5 h in a conventional oven. Upon completion of the reaction, the sample solution was cooled down to room temperature. The Si NPs were purified by dialysis (Dialysis membrane MWCO: 1 kDa) against water. The characterization was performed by TEM, HR-TEM, FTIR, UV-Vis and TGA analysis.

Amine group quantification

TGA analysis¹⁸

The TGA analysis shows four inflexion points at 115 °C (water loss ~5%), 215 °C (organic functional group loss ~17.5%), 465 °C and 630 °C. The HR-TEM diameter of the Si NPs ~3 nm and density of amorphous silica, 2.2 g/cm³ can be used to calculate the surface amine groups.

$$\text{Organic bilayer/inorganic bilayer} = 17.5\%$$

$$\text{Radius} = 1.5 \text{ nm}$$

$$\text{Volume} = 33.5 * 10^{-18} \text{ mm}^3$$

$$\text{mass}_{\text{core}} = 888 * 10^{-18} \text{ mg}$$

$$\text{mass}_{\text{org.bilayer}} = 15.7 * 10^{-18} \text{ mg}$$

$$\text{mass org.bilayer} * NA/M.W.org.bilayer = 162 \text{ amine groups per Si NP}$$

Kaiser Test (Ninhydrin test)

Ninhydrin solution upon reaction with Si NP solution (free amine groups) forms a complex, Ruhemann's purple in weakly acidic solution. The absorbance values at 570 nm from the complex were registered for the accessible amine group quantification.

$$A = \epsilon cl$$

$$c = A / (\epsilon l)$$

$$c = (\text{Absorbance} * \text{Volume} / (\epsilon * \text{Mass Si NPs} * 1))$$

$$= 6.8 \mu\text{mol} / \text{mg Si NP}$$

Synthesis of monomodal Si NP-based imaging agents

Si NP-NOTA

Si NP-NOTA was prepared by dispersing 1 mg of amine-terminated Si NP in deionized water (150 µL) in a 1.5 mL low protein binding Eppendorf-tube. The pH of the dispersion was set between 7–7.5 by 0.01 M NaOH solution. NOTA-Bn-SCN (0.1 µmol) was

¹⁸ The experiment was performed by John Ddungu at University of Strasbourg, France.

dissolved in water (100 μ L) and added to the Si NP dispersion. The reaction was shaken on a thermomixer at 750 rpm for 16 h at 25 $^{\circ}$ C. The Si NP-NOTA was purified by dialyzing against water (MWCO: 500-1000 Da). A defined aliquot of the sample was freeze-dried to obtain the concentration of particles.

Si NP-IR800

To amine-terminated Si NPs (1 mg) dispersed in water (150 μ L) at pH 7–7.5 (0.01 M NaOH), NHS-IR 800 (0.05 μ mol) dissolved in water (100 μ L) was added. The reaction was performed in a 1.5 mL low protein binding Eppendorf-tube in dark. The reaction was shook under a thermomixer at 750 rpm at 35 $^{\circ}$ C. The green solution starts to turn blue in 15 min. Upon completion of the reaction, the product was purified by dialysis (MWCO: 5 kDa) against Millipore water. The Si NP-IR800 was analysed by fluorescence measurements *via* TLC. The sample was further purified by SEC (Sephadex G-25) with 0.9% (wt/v) NaCl as eluent, to obtain pure Si NP-IR800. A known volume of the dispersion was lyophilized to determine the concentration of the nanoconjugate.

Si NP-B4

Si NPs (1 mg) basified to pH 7–7.5 (0.01 M NaOH) was dispersed in water (150 μ L). Bispidine-Bn-SCN (**B4**) was dissolved in a mixture of Water/ACN (9:1) v/v. The reaction was stirred at 35 $^{\circ}$ C for 12 h. The dispersion turned turbid after the addition of **B4**, which clears out in 10 min. The Si NP-B4 was purified by dialysis, followed by SEC (Sephadex G-25 and G-50).

Synthesis of bimodal Si NP-based imaging agent (Si NP-IR800-NOTA)

One-pot synthesis

Si NP (1 mg) dispersed in water (500 μ L) was basified to pH 7–7.5 (0.01 M NaOH). To this solution, NOTA-Bn-SCN (0.1 μ mol) and NHS-IR 800 (0.05 μ mol) separately dissolved in 100 μ L water was added. The reaction was stirred at 35 $^{\circ}$ C for 10–12 h. The product Si NP-IR800-NOTA was purified by dialysis and SEC. The characterization studies displayed non-uniform attachment of the ligands onto the surface of Si NPs. This method was discarded and new stepwise synthetic strategy was applied.

Stepwise synthesis

Si NP (1 mg) dispersed in water (500 μ L) was basified to pH 7–7.5 (0.01 M NaOH). To this solution, NHS-IR 800 (0.05 μ mol) dissolved in 100 μ L was added and the reaction was stirred in dark at 35 $^{\circ}$ C for 8 h. NOTA-Bn-SCN (0.1 μ mol) dissolved in 100 μ L was added to Si NP-IR800 solution and the reaction was performed *in situ*. The reaction was

run for another 8 h at 35 °C. Si NP-IR800-NOTA was purified by dialyzing the solution against water (MWCO 5 kDa). SEC was also performed to yield high purity product. The concentration of the purified NP conjugate was obtained by freeze-drying a known volume of the sample.

Synthesis of targeted Si NP-based imaging agents

Si NP-IR800-PEG-5B9

To synthesize Si NP-IR800-PEG-5B9 bioconjugate, Si NP-IR800 conjugate was freshly synthesized and the next reaction was performed in situ. To this reaction solution, Mal-(PEG)₆-SH (0.15 μmol) was added and the solution was stirred in dark for 2 h at room temperature. Thiol-terminated 5B9 peptide (0.16 μmol) was dissolved in water (200 μL) and TCEP (0.16 μmol) was added to this solution. The reaction was stirred for 15 min and this mixture was then transferred to Si NP conjugate solution. The reaction was now run at room temperature for 5 h. The Si NP-IR800-PEG-5B9 was dialyzed against water and repurified by SEC (Sephadex G-25).

Si NP-IR800-NOTA-PEG-5B9

Si NP-IR800-NOTA was freshly synthesized to perform further modification of the Si NPs. Mal-(PEG)₆-SH (0.15 μmol) was added to this reaction mixture and stirred for 2 h at room temperature. Thiol-terminated 5B9 peptide (0.16 μmol) was dissolved in water (200 μL) and TCEP (0.16 μmol) was added to this solution. The reaction was stirred for 15 min then transferred to the Si NP conjugate solution. The reaction was run at room temperature for 5 h. The Si NP-IR800-NOTA-PEG-5B9 was dialyzed against water and repurified by SEC (Sephadex G-25).

Determination of dye concentration

The IR 800 dye concentration in the solution of unknown concentration after the multifunctionalization was determined by Beer-Lambert's Law. A stock solution of known concentration of IR 800 dye was prepared in Millipore water. 5 samples with increasing concentrations were prepared from the stock. Each dye sample was measured for absorbance keeping the value between 0.01–0.1 AU. A graph was plotted with this data, (absorbance vs concentration) which lead to a straight line passing through the origin. A sample with known amount of Si NP conjugates was measured for its absorbance and plotted on the same graph. The Beer-Lambert's equation was also used for the determination of the amount of dye present per mg of Si NP conjugate.

Relative Quantum Yield Determination

The photoluminescent quantum yield of the fluorophore coupled to unimodal and bimodal Si NP conjugates was determined using free IR 800 dye as a reference. The QY of free NHS-IR 800 dye according to previously published data is 14.2%. The UV-Vis absorbance spectrum of the Si NP conjugates dispersed in water, as well as for the free dye was measured. To minimize the re-absorption effects, the absorbance signals were always kept between 0.01–0.10 AU. The maximum absorbance (~615 nm) was used as the excitation wavelength for the fluorescence emission due to the blue shift in the IR 800 dye signals upon conjugation to Si NPs. A slit width of 5 mm was kept constant for all the measurements. The resulting fluorescence signal was integrated, and the quantum yield was calculated.

5.5 Radiolabeling experiments

⁶⁴Cu-radiolabeling kinetics:

Radiolabeling of (**B1–B5**), **B4**-PEG-TATE and **B5**-sdAb was performed using a standard protocol. Radio-TLC is performed to access the radiolabeling with 2 M NH₄OAc/MeOH (1:2 v/v) as solvent system. The TLC strips are air-dried and exposed to a high-resolution phosphorimaging plate (GE Healthcare). The exposed plates are scanned with an Amersham Typhoon 5 Scanner (GE Healthcare).

The chelators (**B1–B5**) were dissolved in acetonitrile/water solution (1:9 v/v) in stock solutions of 1 mg/mL. In 1.5 mL Eppendorf vials, to a 10 µL aliquot of the stock solutions 100 µL of 0.1 M MES/NaOH buffer was added. To this solution, 6–8 MBq of [⁶⁴Cu]CuCl₂ were added, and the final volume was made up to 200 µL with the buffer solution. The radiolabeling was monitored using radio-TLC using **Method I (Table 6)** at different times. The ⁶⁴Cu-complexes were purified by Radio-HPLC C-18 columns.

For **B5**-sdAb¹⁹, samples of variable volumes of the [⁶⁴Cu]CuCl₂ solution corresponding to different activities (0.5 MBq – 10 MBq) was prepared with constant amount of **B4**-sdAb (1 nmol). The reaction mixtures were incubated at 37°C for different times. The assessment was done by radio-TLC (iTLC-SG) with 5 µL of EDTA (500 mM, pH 7.4) was added to 5 µL sample solutions for challenge experiment and monitored by TLC. The [⁶⁴Cu]Cu-EDTA migrated to the solvent front ($R_f = 1$), whereas the radiolabeled proteins remained at the origin ($R_f = 0$). If necessary, the radiolabeled proteins were purified prior to further use *via* spin filtration (MWCO: 3kDa).

Si NP conjugates (Si NP-NOTA/Si NP-IR800-NOTA)

An aqueous solution of [⁶⁴Cu]CuCl₂ (10 MBq) was added to ~0.1 mg (1 mg/mL stock solution) of Si NP-NOTA and Si NP-IR800-NOTA dispersed in water. 100 mM MES buffer at pH 7.4 was used to make the final volume up to 400 µL. The reaction mixtures were shaken at RT for 30 min, which resulted in RCY ~ 80-90%. The radiolabeling was monitored using radio-TLC (**method II**). Radio-SEC was performed to purify the traces of free [⁶⁴Cu]CuCl₂. The ⁶⁴Cu-complexes were purified using short NAP-5 columns or PD-10 size exclusion columns with the 0.9% NaCl solution as a mobile phase. The radiochemical yields of ⁶⁴Cu(II)-labeled complexes are displayed in **Table** below.

¹⁹ The conjugation, radiolabeling and cell binding studies were performed by Dr. K. Zarschler, Institute of Radiopharmaceutical Cancer Research, HZDR.

⁶⁴ Cu-Bispidine complex	Radiochemical purity (RCY %)
[⁶⁴ Cu]Cu-B1	> 99.9
[⁶⁴ Cu]Cu-B2	> 99.9
[⁶⁴ Cu]Cu-B3	> 99.9
[⁶⁴ Cu]Cu-B4	> 99.1
[⁶⁴ Cu]Cu-B5	≥ 99.0
[⁶⁴ Cu]Cu-B4-PEG-TATE	≥ 99.0
[⁶⁴ Cu]Cu-B5-sdAb	≥ 99.0
[⁶⁴ Cu]Cu-Si NP-NOTA	90–95
[⁶⁴ Cu]Cu-Si NP-IR800-NOTA	80–95

Radiometric titration to determine the amount of chelator on Si NPs

Radiometric titrations were conducted using a mixture of radioactive ⁶⁴Cu(II) and non-radioactive Cu(II) for the determination of the amount of NOTA and bispidine chelator conjugated to Si NPs. To 10 µL aliquots of the Si NP-NOTA, Si NP-B4 and Si NP-IR800-NOTA (0.2 mM stock solution in 100 mM MES buffer, pH 5.5), known amounts of a mixture of [⁶⁴Cu]CuCl₂ and CuSO₄ with a final concentration of 0.4 mM (stock solution) were added. The solutions were stirred for 1 h at room temperature on a thermomixer at 750 rpm. After complete complexation (confirmed using radio-TLC), an 8-fold excess of EDTA was added to each solution vial to remove any non-specifically bound Cu(II) ions. The resulting solutions were stirred at 37 °C for 40 min at 750 rpm. The RCY was determined *via* radio-TLC and also on a radio-SEC (Sephadex G-25). The amount of chelator per mg Si NP conjugate was calculated using the percentage of Cu(II) ions bound to the conjugate.

Using the equation;

$$\text{moles Cu(II)} \cdot \left[\frac{\% \text{ } ^{64}\text{Cu(II)} (Rf < 0.2)}{\% \text{ } ^{64}\text{Cu (total)}} \right] = \text{moles of chelator conjugated}$$

The amount of chelator conjugated to Si NPs was calculated.

5.6 *In vitro* cell experiments²⁰

Cell culture

²⁰ Cell experiments of Si NPs were performed by Mrs. Utta Herzog assisted by Ms. Neena Patel, Institute of Radiopharmaceutical Cancer Research, HZDR.

All biocompatibility assays were conducted as per the protocol specified in the Iso Norm 10993–5:2009.

All cell culture reagents were purchased from Biochrom AG and Sigma-Aldrich unless otherwise stated. The human embryonic kidney cell lines HEK293 (DMSZ number ACC 305) were cultured in DMEM + 10% fetal calf serum (FCS) and incubated in humidified atmosphere of 95% air/ 5% CO₂ at 37°C. The cell number and the viability of the cells was determined using CASY cell counter (Roche Diagnostics) following the manufacturer's protocol. The cell lines were regularly tested and were confirmed to be mycoplasma negative using the LookOut mycoplasma PCR detection kit (Sigma-Aldrich). For the cell viability and LDH assays, the cells were treated with penicillin/streptomycin to avoid any bacterial growth during the experiments.

Assessment of biocompatibility (CyQuant, MTS and LDH assay)

The *in vitro* assessment was performed using MTS (3-(4,5-dimethylthiazol-2-yl)-5-(3-carboxymethoxyphenyl)-2-(4-sulfophenyl)-2H-tetrazolium) calorimetric assay and the CyQUANT® Direct Cell Proliferation Assay.

This MTS assay is based on the conversion of the tetrazolium salt to a formazan dye, presumably by NADPH and NADH produced by dehydrogenase enzymes in metabolically-active cells. The formation of the dye in the presence of the NP is monitored by optical absorbance measurement at 490 nm, and the viability is finally compared with the untreated cellular control. In addition to that, the membrane integrity was measured by an LDH assay which quantifies lactate dehydrogenase (LDH), an enzyme that is released upon cell lysis. The CyQUANT assay on the contrary is a fluorescence based proliferation and cytotoxicity assay and therefore can be used to assess cell growth, cell viability, or compound toxicity in a range of applications, from high-throughput screening to bioproduction.

For the evaluation of biocompatibility, non-cancerous cell line, HEK 293 Human epithelial kidney; was seeded in sterile 96-well microtiter plates (Greiner Bio-One) at a density of 1.8×10^4 cells per well were seeded in DMEM media and allowed to grow for 24 h prior to addition of the selected samples.

CyQUANT® is a direct fluorescence-based proliferation and cytotoxicity assay consisting of a fluorescent nucleic acid stain that permeates live cells and concentrates in its nucleus. The second component is a suppresser dye that is impermeable for live cells and suppresses the green fluorescence. 100 µL of CyQuant dye was added to the wells after 24 h of incubation. The fluorescence was measured after 60 min of incubation.

After 24 h incubation, increasing concentrations of Si NP-NOTA, Si NP-IR800 and Si NP-IR800-NOTA conjugates were added to the cells in triplicate. After exposure for 72 h, 20 μ L of MTS (Promega) was added to the wells and incubated for 40 min. Optical densities at 492 nm were measured with a microplate reader (TECAN sunrise). The viability of cells is expressed as a percentage of viable cells grown in the absence of the NPs. Additionally, to measure the membrane integration, an LDH assay was used in the same cell lines. For this, 2.1×10^4 cells per well were seeded in a 96 microtiter plate. After 24 h incubation, the stock solutions of the conjugates were added and incubated for 72 h. 10 μ L of LDH lysis buffer (cytoscan LDH from G-biosciences) was added to the cells. Optical density was measured at 492 nm. The percentage of LDH release was expressed as the percentage of LDH release of cells grown in the absence of NPs. The data were analyzed by normalizing the absorbance from each well containing the samples to the absorbance value of the cells without any samples. As a positive control, a set of a triplicate of cells were treated with the supplied lysis buffer and the percentage of LDH release was expressed relative to this control. As a negative control, bovine serum albumin in triplicate of cells was used.

5.7 Cell binding studies

A total of 60,000 A431 cells were seeded in 48-well plates in 250 μ L DMEM supplemented with 10% fetal calf serum (FCS), and incubated in a humidified atmosphere of 95% air/5% CO₂ at 37°C. After 24 h of incubation, media was replaced by serum-free DMEM and cultivation was continued. After 4 h of incubation, media was removed and 100 μ L of ice-cold PBS (total binding) or 100 μ L of 1 μ M sdAb in ice-cold PBS (unspecific binding), respectively, was added prior to incubation of the cells at 4°C. After 30 min, different concentrations of ⁶⁴Cu-labeled sdAb-**B4** in ice-cold PBS were added. The cell culture microplates were further incubated for 1 h at 4°C. Following treatment with radiolabeled sdAbs, cells were washed three times with ice-cold PBS in order to ensure removal of loosely attached proteins from the cell membrane. Finally, cell lysis was achieved by the addition of 1% SDS/0.1 M NaOH and incubation for 30 min at room temperature with vigorous shaking. The cell associated radioactivity was quantified using an automated gamma counter (PerkinElmer Life and Analytical Sciences). Total protein concentration in cell extracts was determined as described above.

5.8 Animal experiments²¹

In vivo small animal PET

All animal experiments were carried out according to the guidelines of the German Regulations for Animal Welfare and the protocols approved by the local Ethical Committee for Animal Experiments (AZ 24-9168.21-4/2004-1, AZ 24-9168.11-4/2012-1). Animals received standard food and tap water *ad libitum* and were anesthetized prior to animal experiments using 10% desflurane in 30% oxygen/air. Male Wistar-Unilever rats (HsdCpb:WU) and male and female NMRI nu/nu mice (RjOrl:NMRI-Foxn1nu/Foxn1nu) were received from Harlan Winkelmann GmbH (Borchen, Germany) and Janvier Labs (Saint-Berthevin Cedex, France), respectively. Small animal PET imaging was performed using microPET® P4 scanner (Siemens Medical Solutions, Knoxville) or NanoScanPET/CT (Mediso, Budapest). Animals were positioned on a heated bed to maintain body temperature at 37°C.

For the generation of A431 tumor xenograft model 2×10^6 human A431 epidermoid (squamous) carcinoma cells (ATCC: CRL 1555) were subcutaneously injected into the right leg of six weeks old female NMRI nu/nu mice as described elsewhere

Rats and mice received 10-30 MBq (50–100 µg) of Si NP conjugates in 0.5 mL and 0.2 mL E-153, respectively, intravenously over 0.5 min into a lateral tail vein. The activity of the injection solution was measured in a dose calibrator (Isomed 2000, Dresden, Germany) cross-calibrated to the PET scanners.

Attenuation correction of PET data was performed by means of a 10 min transmission scan using a rotating point source of ⁵⁷Co (microPET®) or by whole body CT (NanoScan®PET/CT). Data acquisition and reconstruction were performed as described elsewhere. Three-dimensional regions of interest (ROI) were determined within masks around the tumors or different organs by thresholding PET data within these masks. ROI time activity curves (TAC) were derived from subsequent data analysis. Data were calculated as standard uptake value (SUV, SUV = activity concentration in tissue [Bq/g] × body weight [g] / injected dose [Bq]) at a certain time point *p.i.* The ROI data and TAC were further analyzed using R (R is available as Free Software under the terms of the Free Software Foundation's GNU General Public License in source code form) and specially developed program packages (Jörg van den Hoff, Helmholtz-Zentrum Dresden

²¹ All animal experiments were performed under the supervision of Prof. Pietzsch. PET experiments were performed by Dr. Ralf Bergmann and Ms. Regina Herrlich at the Institute of Radiopharmaceutical Cancer Research, HZDR.

- Rossendorf, Dresden, Germany). The data were calculated in standard uptake value (SUV, in units of grams per milliliter), defined as the tracer concentration at a certain time point normalized to injected dose per unit body weight. The SUV was used for better comparison within animals of different size and weight and with other species. The graphs were calculated with GraphPad Prism version 5.00 for Windows (GraphPad Software, San Diego California USA, www.graphpad.com).

Ex-vivo whole-body autoradiography

Following PET imaging, 100 µm thick whole body coronal cryosections were prepared from representative animals using a Leica Jung Cryopolytcut (Leica Microsystems, Wetzlar, Germany) as described elsewhere.

Biodistribution studies

For biodistribution experiments, rats and A431 tumor-bearing NMRI nu/nu mice were intravenously injected with ~0.5 mL 0.5-10 MBq (10-100 µg) of ⁶⁴Cu-labeled [⁶⁴Cu]Cu-Si NP-NOTA and [⁶⁴Cu]Cu-Si NP-IR800-NOTA or ⁶⁸Ga labeled [⁶⁸Ga]Ga-Si NP-IR800-NOTA conjugates. Animals were sacrificed at 5 min, 60 min and 24 h *p.i.* Organs and tissues of interest were excised, weighed, and radioactivity was determined using Wizard™3" gamma counter. Activity in selected organs and tissues was expressed as % injected dose (%ID), %ID/g tissue, or SUV. The obtained blood volume amounted to 400-500 µL. Afterwards, urine samples (200 - 400 µL) were obtained by puncture of bladder.

Optical imaging

For optical imaging experiments, anesthetized A431 tumor-bearing and healthy NMRI nu/nu mice were intravenously injected with 20 µl IR 800-labeled Si NP-IR800 and Si NP-IR800-NOTA (~80 µg each). Fluorescence was measured using the small animal In-vivo-Imaging System FX Pro (Carestream Health, USA) in fluorescence mode (Ex 600 nm, Em 753 nm; reference image for autofluorescence: Ex 430 nm, Em 535 nm). Fluorescence pictures were merged with the X-ray acquisitions. Afterwards, animals were sacrificed, organs and tissues of interest were excised, and fluorescence was measured again.

Chapter 6 References

- [1] Bray F, F.J., Soerjomataram I, Siegel RL, Torre LA, Jemal A, Global cancer statistics 2018: GLOBOCAN estimates of incidence and mortality worldwide for 36 cancers in 185 countries. *CA Cancer J. Clin.* **6** (2018) 394-424.
- [2] Meyer, T., et al., Accumulating advantages, reducing limitations: multimodal nonlinear imaging in biomedical sciences - the synergy of multiple contrast mechanisms. *J. Biophotonics.* **6** (2013) 887-904.
- [3] Rieffel, J., U. Chitgupi, and J.F. Lovell, Recent Advances in Higher-Order, Multimodal, Biomedical Imaging Agents. *Small.* **11** (2015) 4445-4461.
- [4] James, M.L. and S.S. Gambhir, A molecular imaging primer: modalities, imaging agents, and applications. *Physiol. Rev.* **92** (2012) 897-965.
- [5] Condeelis, J. and R. Weissleder, In vivo imaging in cancer. *Cold Spring Harb. Perspect. Biol.* **2** (2010) 1-22.
- [6] Grenier, N. and P. Brader, Principles and basic concepts of molecular imaging. *Pediatr. Radiol.* **41** (2011) 144-160.
- [7] Key, J. and J.F. Leary, Nanoparticles for multimodal in vivo imaging in nanomedicine. *Int. J. Nanomedicine.* **9** (2014) 711-726.
- [8] Opacic, T., et al., Status and trends in the development of clinical diagnostic agents. *Wiley Interdiscip. Rev. Nanomed. Nanobiotechnol.* **9** (2017).
- [9] Weber, J., U. Haberkorn, and W. Mier, Cancer stratification by molecular imaging. *Int. J. Mol. Sci.* **16** (2015) 4918-4946.
- [10] Winfield, J.M., G.S. Payne, and N.M. deSouza, Functional MRI and CT biomarkers in oncology. *Eur. J. Nucl. Med. Mol. Imaging.* **42** (2015) 562-578.
- [11] Frangioni, J.V., New technologies for human cancer imaging. *J. Clin. Oncol.* **26** (2008) 4012-4021.
- [12] Wu, M. and J. Shu, Multimodal Molecular Imaging: Current Status and Future Directions. *Contrast Media Mol. Imaging.* **2018** (2018) 1-12.
- [13] Endo, K., et al., PET and PET/CT using 18F-FDG in the diagnosis and management of cancer patients. *Int. J. Clin. Oncol.* **11** (2006) 286-296.
- [14] Pandey, S.K., et al., Multimodality imaging probe for positron emission tomography and fluorescence imaging studies. *Mol. Imaging.* **13** (2014) 1-7.
- [15] Summer, D., et al., Developing Targeted Hybrid Imaging Probes by Chelator Scaffolding. *Bioconjug. Chem.* **28** (2017) 1722-1733.

-
- [16] Wadas, T.J., et al., Coordinating radiometals of copper, gallium, indium, yttrium, and zirconium for PET and SPECT imaging of disease. *Chem. Rev.* **110** (2010) 2858-2902.
- [17] Wang, L., Y. Wang, and Z. Li, Nanoparticle-based tumor theranostics with molecular imaging. *Curr. Pharm. Biotechnol.* **14** (2013) 683-692.
- [18] Pelaz, B., et al., Diverse Applications of Nanomedicine. *ACS Nano.* **11** (2017) 2313-2381.
- [19] Thakor, A.S., et al., Clinically Approved Nanoparticle Imaging Agents. *J. Nucl. Med.* **57** (2016) 1833-1837.
- [20] Anselmo, A.C. and S. Mitragotri, Nanoparticles in the clinic. *Bioeng. Transl. Med.* **1** (2016) 10-29.
- [21] Jokerst, J.V. and S.S. Gambhir, Molecular imaging with theranostic nanoparticles. *Acc. Chem. Res.* **44** (2011) 1050-1060.
- [22] Min, Y., et al., Clinical Translation of Nanomedicine. *Chem. Rev.* **115** (2015) 11147-11190.
- [23] Phillips, E., et al., Clinical translation of an ultrasmall inorganic optical-PET imaging nanoparticle probe. *Sci. Transl. Med.* **6** (2014) 260ra149.
- [24] Huang, H.C., et al., Inorganic nanoparticles for cancer imaging and therapy. *J. Control Release.* **155** (2011) 344-357.
- [25] Anselmo, A.C. and S. Mitragotri, A Review of Clinical Translation of Inorganic Nanoparticles. *AAPS J.* **17** (2015) 1041-1054.
- [26] Zarschler, K., et al., Ultrasmall inorganic nanoparticles: State-of-the-art and perspectives for biomedical applications. *Nanomedicine.* **12** (2016) 1663-1701.
- [27] Baetke, S.C., T. Lammers, and F. Kiessling, Applications of nanoparticles for diagnosis and therapy of cancer. *Br. J. Radiol.* **88** (2015) 1-12.
- [28] Ahire, J.H., et al., Synthesis of D-mannose capped silicon nanoparticles and their interactions with MCF-7 human breast cancerous cells. *ACS Appl. Mater. Interfaces.* **5** (2013) 7384-7391.
- [29] Barreto, J.A., et al., Nanomaterials: applications in cancer imaging and therapy. *Adv. Mater.* **23** (2011) H18-40.
- [30] Feliu, N., et al., In vivo degeneration and the fate of inorganic nanoparticles. *Chem. Soc. Rev.* **45** (2016) 2440-2457.
- [31] Jin, Y., et al., Tracking the Fate of Porous Silicon Nanoparticles Delivering a Peptide Payload by Intrinsic Photoluminescence Lifetime. *Adv. Mater.* **30** (2018) e1802878.
- [32] Kiessling, F., et al., Nanoparticles for imaging: top or flop? *Radiology.* **273** (2014) 10-28.

-
- [33] Kleps, I., et al., Nanostructured silicon particles for medical applications. *J. Nanosci. Nanotechnol.* **10** (2010) 2694-2700.
- [34] Kunz-Schughart, L.A., et al., Nanoparticles for radiooncology: Mission, vision, challenges. *Biomaterials.* **120** (2017) 155-184.
- [35] Kunjachan, S., et al., Noninvasive Imaging of Nanomedicines and Nanotheranostics: Principles, Progress, and Prospects. *Chem. Rev.* **115** (2015) 10907-10937.
- [36] Ma, Y.Y., et al., Molecular Imaging of Cancer with Nanoparticle-Based Theranostic Probes. *Contrast Media Mol. Imaging.* **2017** (2017) 1-11.
- [37] Sun, X., W. Cai, and X. Chen, Positron emission tomography imaging using radiolabeled inorganic nanomaterials. *Acc. Chem. Res.* **48** (2015) 286-294.
- [38] Bradbury, M.S., et al., Clinically-translated silica nanoparticles as dual-modality cancer-targeted probes for image-guided surgery and interventions. *Integr. Biol. (Camb).* **5** (2013) 74-86.
- [39] Ji, X., et al., Silicon Nanomaterials for Biosensing and Bioimaging Analysis. *Front. Chem.* **6** (2018) 1-9.
- [40] Tu, C., et al., PET Imaging and biodistribution of silicon quantum dots in mice. *ACS Med. Chem. Lett.* **2** (2011) 285-288.
- [41] Peng, F., et al., Silicon Nanomaterials Platform for Bioimaging, Biosensing, and Cancer Therapy. *Acc. Chem. Res.* **47** (2014) 612-623.
- [42] Park, J.-H., et al., Biodegradable luminescent porous silicon nanoparticles for in vivo applications. *Nat. Mater.* **8** (2009) 331-336.
- [43] Tasciotti, E., et al., Mesoporous silicon particles as a multistage delivery system for imaging and therapeutic applications. *Nat. Nanotechnol.* **3** (2008) 151-157.
- [44] Zhong, Y., et al., Microwave-Assisted Synthesis of Biofunctional and Fluorescent Silicon Nanoparticles Using Proteins as Hydrophilic Ligands. *Angew. Chem. Int. Ed. Engl.* **51** (2012) 8485-8489.
- [45] Tu, C., et al., Paramagnetic, Silicon Quantum Dots for Magnetic Resonance and Two-Photon Imaging of Macrophages. *J. Am. Chem. Soc.* **132** (2010) 2016-2023.
- [46] Blanco, E., H. Shen, and M. Ferrari, Principles of nanoparticle design for overcoming biological barriers to drug delivery. *Nat. Biotechnol.* **33** (2015) 941-951.
- [47] Choi, J.S., et al., A hybrid nanoparticle probe for dual-modality positron emission tomography and magnetic resonance imaging. *Angew. Chem. Int. Ed. Engl.* **47** (2008) 6259-6262.

-
- [48] Li, Q., et al., Surface-modified silicon nanoparticles with ultrabright photoluminescence and single-exponential decay for nanoscale fluorescence lifetime imaging of temperature. *J Am. Chem. Soc.* **135** (2013) 14924-14927.
- [49] Li, Q., et al., Silicon Nanoparticles with Surface Nitrogen: 90% Quantum Yield with Narrow Luminescence Bandwidth and the Ligand Structure Based Energy Law. *ACS Nano*. **10** (2016) 8385-8393.
- [50] Li, X., et al., Multimodality imaging in nanomedicine and nanotheranostics. *Cancer Biol. Med.* **13** (2016) 339-348.
- [51] Liu, J., et al., One-pot synthesis of silicon based nanoparticles with incorporated phthalocyanine for long-term bioimaging and photo-dynamic therapy of tumors. *Nanotechnology*. **1-8** (2017) 135601.
- [52] McBride, J.R., et al., On Ultrasmall Nanocrystals. *Chem. Phys. Lett.* **498** (2010) 1-9.
- [53] Rodio, M., et al., Direct surface modification of ligand-free silicon quantum dots prepared by femtosecond laser ablation in deionized water. *J. Colloid Interface Sci.* **465** (2016) 242-248.
- [54] Shiohara, A., et al., Chemical reactions on surface molecules attached to silicon quantum dots. *J. Am. Chem. Soc.* **132** (2010) 248-253.
- [55] Su, X., et al., Mild two-step method to construct DNA-conjugated silicon nanoparticles: scaffolds for the detection of microRNA-21. *Bioconjug. Chem.* **25** (2014) 1739-1743.
- [56] Zhang, W., et al., Fluorescent proteins as efficient tools for evaluating the surface PEGylation of silica nanoparticles. *Methods Appl. Fluoresc.* **5** (2017) 1-9.
- [57] Erogbogbo, F., et al., Biocompatible Luminescent Silicon Quantum Dots for Imaging of Cancer Cells. *ACS Nano*. **2** (2008) 873-878.
- [58] Rampazzo, E., et al., NIR-fluorescent dye doped silica nanoparticles for in vivo imaging, sensing and theranostic. *Methods Appl. Fluoresc.* **6** (2018) 1-21.
- [59] Rosenholm, J.M., et al., Prolonged Dye Release from Mesoporous Silica-Based Imaging Probes Facilitates Long-Term Optical Tracking of Cell Populations In Vivo. *Small*. **12** (2016) 1578-1592.
- [60] Tasciotti, E., et al., Near-infrared imaging method for the in vivo assessment of the biodistribution of nanoporous silicon particles. *Mol. Imaging*. **10** (2011) 56-68.
- [61] Kim, H., et al., Near-infrared light-responsive nanomaterials for cancer theranostics. *Wiley Interdiscip. Rev. Nanomed. Nanobiotechnol.* **8** (2016) 23-45.
- [62] Anderson, C.J. and R. Ferdani, Copper-64 radiopharmaceuticals for PET imaging of cancer: advances in preclinical and clinical research. *Cancer Biother. Radiopharm.* **24** (2009) 379-393.

-
- [63] Miyamoto, T., et al., [Development of new peptide-based chelating agents for site-specific radiolabeling with ^{64}Cu]. *Yakugaku Zasshi*. **134** (2014) 797-800.
- [64] Thieme, S., et al., Module-assisted preparation of ^{64}Cu with high specific activity. *Appl. Radiat. Isot.* **70** (2012) 602-608.
- [65] Banerjee, S.R. and M.G. Pomper, Clinical applications of Gallium-68. *Appl. Radiat. Isot.* **76** (2013) 2-13.
- [66] Brechbiel, M.W., Bifunctional chelates for metal nuclides. *Q. J. Nucl. Med. Mol. Imaging*. **52** (2008) 166-173.
- [67] Liu, S., Bifunctional coupling agents for radiolabeling of biomolecules and target-specific delivery of metallic radionuclides. *Adv. Drug Deliv. Rev.* **60** (2008) 1347-1370.
- [68] Pant, K., et al., Radiolabelled Polymeric Materials for Imaging and Treatment of Cancer: Quo Vadis? *Adv. Healthc. Mater.* **6** (2017) 1-31.
- [69] Sarko, D., et al., Bifunctional Chelators in the Design and Application of Radiopharmaceuticals for Oncological Diseases. *Curr. Med. Chem.* **19** (2012) 2667-2688.
- [70] Spang, P., C. Herrmann, and F. Roesch, Bifunctional Gallium-68 Chelators: Past, Present, and Future. *Semin. Nucl. Med.* **46** (2016) 373-394.
- [71] Cooper, M.S., et al., Comparison of ^{64}Cu -Complexing Bifunctional Chelators for Radioimmunoconjugation: Labeling Efficiency, Specific Activity, and in Vitro/in Vivo Stability. *Bioconjug. Chem.* **23** (2012) 1029-1039.
- [72] Price, E.W. and C. Orvig, Matching chelators to radiometals for radiopharmaceuticals. *Chem. Soc. Rev.* **43** (2014) 260-290.
- [73] Ramogida, C.F. and C. Orvig, Tumour targeting with radiometals for diagnosis and therapy. *Chem. Commun. (Camb)*. **49** (2013) 4720-4739.
- [74] Chiavarelli, S., F. Toffler, and D. Misiti, [Chemistry of bispidine and its derivatives]. *Ann. Ist. Super. Sanità*. **4** (1968) 157-185.
- [75] Tomassoli, I. and D. Gundisch, Bispidine as a Privileged Scaffold. *Curr. Top. Med. Chem.* **16** (2016) 1314-1342.
- [76] Juran, S., et al., Hexadentate bispidine derivatives as versatile bifunctional chelate agents for copper(II) radioisotopes. *Bioconjug. Chem.* **20** (2009) 347-359.
- [77] Roux, A., et al., Bifunctional bispidine derivatives for copper-64 labelling and positron emission tomography. *Org. Biomol. Chem.* **15** (2017) 1475-1483.
- [78] Stephan, H., et al., Bispidines for dual imaging. *Chemistry*. **20** (2014) 17011-17018.
- [79] Ait-Mohand, S., et al., Evaluation of ^{64}Cu -labeled bifunctional chelate-bombesin conjugates. *Bioconjug. Chem.* **22** (2011) 1729-1735.

-
- [80] Mankoff, D.A., A definition of molecular imaging. *J. Nucl. Med.* **48** (2007) 18n, 21n.
- [81] Anderson, C.J. and J.S. Lewis, Current status and future challenges for molecular imaging. *Philos. Trans. A Math. Phys. Eng. Sci.* **375** (2017).
- [82] Chen, F., et al., Dynamic Positron Emission Tomography Imaging of Renal Clearable Gold Nanoparticles. *Small*. **12** (2016) 2775-2782.
- [83] Cheng, X., et al., Versatile "Click Chemistry" Approach to Functionalizing Silicon Quantum Dots: Applications toward Fluorescent Cellular Imaging. *Langmuir*. **30** (2014) 5209-5216.
- [84] Zhou, Y., et al., The in vivo targeted molecular imaging of fluorescent silicon nanoparticles in *Caenorhabditis elegans*. *Nano Res.* **11** (2018) 2336-2346.
- [85] Weissleder, R. and U. Mahmood, Molecular imaging. *Radiology*. **219** (2001) 316-333.
- [86] Hu, Z., et al., Nanoparticle-mediated radiopharmaceutical-excited fluorescence molecular imaging allows precise image-guided tumor-removal surgery. *Nanomedicine*. **13** (2017) 1323-1331.
- [87] Hu, Z., et al., In vivo nanoparticle-mediated radiopharmaceutical-excited fluorescence molecular imaging. *Nat. Commun.* **6** (2015) 1-12.
- [88] Puranik, A.D., H.R. Kulkarni, and R.P. Baum, Companion Diagnostics and Molecular Imaging. *Cancer J.* **21** (2015) 213-217.
- [89] Saint-Aubert, L., et al., Tau PET imaging: present and future directions. *Mol. Neurodegener.* **12** (2017) 1-21.
- [90] Cutler, C.S., et al., Radiometals for combined imaging and therapy. *Chem. Rev.* **113** (2013) 858-883.
- [91] Wadas, T.J., et al., Copper chelation chemistry and its role in copper radiopharmaceuticals. *Curr Pharm Des.* **13** (2007) 3-16.
- [92] Differding, S., F.X. Hanin, and V. Gregoire, PET imaging biomarkers in head and neck cancer. *Eur. J. Nucl. Med. Mol. Imaging*. **42** (2015) 613-622.
- [93] Lane, S.R., et al., Optimization, biological evaluation and microPET imaging of copper-64-labeled bombesin agonists, [64Cu-NO2A-(X)-BBN(7-14)NH₂], in a prostate tumor xenografted mouse model. *Nucl. Med. Biol.* **37** (2010) 751-761.
- [94] Moreau, M., et al., MANOTA: a promising bifunctional chelating agent for copper-64 immunoPET. *Dalton Trans.* **46** (2017) 14659-14668.
- [95] Shokeen, M. and C.J. Anderson, Molecular Imaging of Cancer with Copper-64 Radiopharmaceuticals and Positron Emission Tomography (PET). *Acc. Chem. Res.* **42** (2009) 832-841.

- [96] Zhang, Y., et al., Positron emission tomography imaging of CD105 expression with a ^{64}Cu -labeled monoclonal antibody: NOTA is superior to DOTA. *PLoS One*. **6** (2011) 1-7.
- [97] Joshi, T., et al., Harnessing the Coordination Chemistry of 1,4,7-Triazacyclononane for Biomimicry and Radiopharmaceutical Applications. *ChemPlusChem*. **83** (2018) 554-564.
- [98] Bailey, G.A., et al., H(2)azapa: a versatile acyclic multifunctional chelator for (67)Ga, (64)Cu, (111)In, and (177)Lu. *Inorg. Chem.* **51** (2012) 12575-12589.
- [99] Li, J., et al., Cyclam-based polymeric copper chelators for gene delivery and potential PET imaging. *Biomacromolecules*. **13** (2012) 3220-3227.
- [100] Anderson, C.J., et al., ^{64}Cu -TETA-octreotide as a PET imaging agent for patients with neuroendocrine tumors. *J. Nucl. Med.* **42** (2001) 213-221.
- [101] Comba, P.K., M.; Schiek W., *Bispidine Coordination Chemistry*. Progress in Inorganic Chemistry, ed. K.D. Karlin. Vol. 55. 2008: John Wiley & Sons, Inc.
- [102] Comba, P., et al., Bispidines for radiopharmaceuticals. *Dalton Trans.* **47** (2018) 9202-9220.
- [103] Roux, A., et al., Kinetically Inert Bispidol-Based Cu(II) Chelate for Potential Application to (64/67)Cu Nuclear Medicine and Diagnosis. *Inorg. Chem.* **54** (2015) 4431-4444.
- [104] Rudin, M., Noninvasive structural, functional, and molecular imaging in drug development. *Curr. Opin. Chem. Biol.* **13** (2009) 360-371.
- [105] Ruenitz, P.C. and C.M. Mokler, Analogues of sparteine. 5. Antiarrhythmic activity of selected N,N'-disubstituted bispidines. *J. Med. Chem.* **20** (1977) 1668-1671.
- [106] Xu, F., et al., Highly selective cascade couplings for the syntheses of functionalized piperidinones and bispidines. *Org. Lett.* **9** (2007) 2669-2672.
- [107] Comba, P., Fit and misfit between ligands and metal ions. *Coord. Chem. Rev.* **238-239** (2003) 21-29.
- [108] Price, E.W. and C. Orvig, Matching chelators to radiometals for radiopharmaceuticals. *Chem Soc Rev.* **43** (2014) 260-290.
- [109] Tanaka, K. and A. R. Pradipta, Synthesis of 3,7,9- and 2,6,9-Triazabicyclo[3.3.1]Nonane Derivatives. *Heterocycles*. **87** (2013).
- [110] Comba, P., et al., Optimization of pentadentate bispidines as bifunctional chelators for ^{64}Cu positron emission tomography (PET). *Inorg. Chem.* **52** (2013) 8131-8143.
- [111] Comba, P., et al., Bispidine dioxotetraaza macrocycles: a new class of bispidines for (64)Cu PET imaging. *Inorg. Chem.* **53** (2014) 6698-6707.

- [112] Comba, P., et al., The computation of lipophilicities of (64)Cu PET systems based on a novel approach for fluctuating charges. *Dalton Trans.* **42** (2013) 11066-11073.
- [113] Comba, P., et al., Octadentate Picolinic Acid-Based Bispidine Ligand for Radiometal Ions. *Chemistry*. **23** (2017) 15945-15956.
- [114] Comba, P., M. Starke, and H. Wadepohl, Optimization of Hexadentate Bispidine Ligands as Chelators for ⁶⁴Cu^{II} PET Imaging. *ChemPlusChem*. **83** (2018) 597-604.
- [115] Cusnir, R., et al., Hydroxypyridinone Chelators: From Iron Scavenging to Radiopharmaceuticals for PET Imaging with Gallium-68. *Int. J. Mol. Sci.* **18** (2017) 1-23.
- [116] Sarko, D., et al., Bifunctional chelators in the design and application of radiopharmaceuticals for oncological diseases. *Curr Med Chem*. **19** (2012) 2667-2688.
- [117] Tornesello, A.L., M.L. Tornesello, and F.M. Buonaguro, An Overview of Bioactive Peptides for in vivo Imaging and Therapy in Human Diseases. *Mini Rev. Med. Chem.* **17** (2017) 758-770.
- [118] Dommerholt, J., F. Rutjes, and F.L. van Delft, Strain-Promoted 1,3-Dipolar Cycloaddition of Cycloalkynes and Organic Azides. *Top. Curr. Chem. (Cham)*. **374** (2016) 1-20.
- [119] Kobayashi, H., et al., Multimodal nanoprobe for radionuclide and five-color near-infrared optical lymphatic imaging. *ACS Nano*. **1** (2007) 258-264.
- [120] Mérian, J., et al., Fluorescent Nanoprobes Dedicated to in Vivo Imaging: From Preclinical Validations to Clinical Translation. *Molecules*. **17** (2012) 5564-5591.
- [121] Conner, K.P., et al., Evaluation of near infrared fluorescent labeling of monoclonal antibodies as a tool for tissue distribution. *Drug Metab. Dispos.* **42** (2014) 1906-1913.
- [122] Debie, P., et al., Effect of Dye and Conjugation Chemistry on the Biodistribution Profile of Near-Infrared-Labeled Nanobodies as Tracers for Image-Guided Surgery. *Mol. Pharm.* **14** (2017) 1145-1153.
- [123] Liu, Z., et al., Advanced nanomaterials in multimodal imaging: design, functionalization, and biomedical applications. *J. Nanomaterials*. **2010** (2010) 1-15.
- [124] Zheng, X., et al., Applications of silica-based nanoparticles for multimodal bioimaging. *Applied Spectroscopy Reviews*. **53** (2018) 377-394.
- [125] Auletta, L., et al., Advances in multimodal molecular imaging. *Q. J. Nucl. Med. Mol. Imaging*. **61** (2017) 19-32.

-
- [126] Babinec, P. and M. Babincova, Towards multimodal nanoparticle labels for molecular imaging of biological processes. *Med. Hypotheses*. **69** (2007) 703-704.
- [127] Martinez-Rodriguez, I. and I. Banzo, Advances in PET: The success of multimodal molecular imaging. *Med. Clin. (Barc)*. **148** (2017) 354-356.
- [128] Melendez-Alafort, L., P.C. Muzzio, and A. Rosato, Optical and multimodal peptide-based probes for in vivo molecular imaging. *Anticancer Agents Med. Chem.* **12** (2012) 476-499.
- [129] Spinelli, A.E., et al., Optical imaging of radioisotopes: a novel multimodal approach to molecular imaging. *Q. J. Nucl. Med. Mol. Imaging*. **56** (2012) 280-290.
- [130] Swierczewska, M., S. Lee, and X. Chen, Inorganic nanoparticles for multimodal molecular imaging. *Mol. Imaging*. **10** (2011) 3-16.
- [131] Bao, G., S. Mitragotri, and S. Tong, Multifunctional nanoparticles for drug delivery and molecular imaging. *Annu. Rev. Biomed. Eng.* **15** (2013) 253-282.
- [132] Behray, M., et al., Synthesis of Diagnostic Silicon Nanoparticles for Targeted Delivery of Thiourea to Epidermal Growth Factor Receptor-Expressing Cancer Cells. *ACS Appl. Mater. Interfaces*. **8** (2016) 8908-8917.
- [133] Joshi, B.P. and T.D. Wang, Exogenous Molecular Probes for Targeted Imaging in Cancer: Focus on Multi-modal Imaging. *Cancers (Basel)*. **2** (2010) 1251-1287.
- [134] Feldborg, L.N., R.I. Jolck, and T.L. Andresen, Quantitative evaluation of bioorthogonal chemistries for surface functionalization of nanoparticles. *Bioconjug. Chem.* **23** (2012) 2444-2450.
- [135] Lee, D.E., et al., Multifunctional nanoparticles for multimodal imaging and theragnosis. *Chem. Soc. Rev.* **41** (2012) 2656-2672.
- [136] Lee, H., et al., Silica nanoparticle-based dual imaging colloidal hybrids: cancer cell imaging and biodistribution. *Int. J. Nanomedicine*. **10 Spec Iss** (2015) 215-225.
- [137] Rizzo, L.Y., et al., Recent progress in nanomedicine: therapeutic, diagnostic and theranostic applications. *Curr. Opin. Biotechnol.* **24** (2013) 1159-1166.
- [138] Rosenholm, J.M., C. Sahlgren, and M. Linden, Multifunctional mesoporous silica nanoparticles for combined therapeutic, diagnostic and targeted action in cancer treatment. *Curr. Drug Targets*. **12** (2011) 1166-1186.
- [139] Wang, W., et al., A new optical and nuclear dual-labeled imaging agent targeting interleukin 11 receptor alpha-chain. *Bioconjug. Chem.* **18** (2007) 397-402.
- [140] Joo, J., et al., Gated Luminescence Imaging of Silicon Nanoparticles. *ACS Nano*. **9** (2015) 6233-6241.

-
- [141] Rosso-Vasic, M., et al., Amine-terminated silicon nanoparticles: synthesis, optical properties and their use in bioimaging. *J. Mater. Chem.* **19** (2009) 5926-5933.
- [142] Cheng, X., et al., Colloidal silicon quantum dots: from preparation to the modification of self-assembled monolayers (SAMs) for bio-applications. *Chem. Soc. Rev.* **43** (2014) 2680-2700.
- [143] Warner, J.H., et al., Water-soluble photoluminescent silicon quantum dots. *Angew. Chem. Int. Ed. Engl.* **44** (2005) 4550-4554.
- [144] He, Y., et al., One-Pot Microwave Synthesis of Water-Dispersible, Ultraphoto- and pH-Stable, and Highly Fluorescent Silicon Quantum Dots. *J. Am. Chem. Soc.* **133** (2011) 14192-14195.
- [145] Song, C., et al., Peptide-Conjugated Fluorescent Silicon Nanoparticles Enabling Simultaneous Tracking and Specific Destruction of Cancer Cells. *Anal. Chem.* **87** (2015) 6718-6723.
- [146] He, Y., et al., Photo and pH Stable, Highly-Luminescent Silicon Nanospheres and Their Bioconjugates for Immunofluorescent Cell Imaging. *J. Am. Chem. Soc.* **131** (2009) 4434-4438.
- [147] Zhong, Y., et al., Large-Scale Aqueous Synthesis of Fluorescent and Biocompatible Silicon Nanoparticles and Their Use as Highly Photostable Biological Probes. *J Am. Chem. Soc.* **135** (2013) 8350-8356.
- [148] Ruizendaal, L., et al., Biofunctional silicon nanoparticles by means of thiol-ene click chemistry. *Chem. Asian. J.* **6** (2011) 2776-2786.
- [149] Wang, L., V. Reipa, and J. Blasic, Silicon nanoparticles as a luminescent label to DNA. *Bioconjug. Chem.* **15** (2004) 409-412.
- [150] Intartaglia, R., et al., Bioconjugated silicon quantum dots from one-step green synthesis. *Nanoscale.* **4** (2012) 1271-1274.
- [151] Erogbogbo, F. and M.T. Swihart. *Imaging pancreatic cancer with folic acid terminated luminescent silicon nanocrystals.* in *Bonsai Project Symposium: Breakthroughs in Nanoparticles for Bio-Imaging*. 2010. ENEA Res Ctr Frascati, Rome, ITALY: American Institute Of Physics 2 Huntington Quad Melville, NY 11747-4501 USA.
- [152] Ahire, J.H., et al., Synthesis of Carbohydrate Capped Silicon Nanoparticles and their Reduced Cytotoxicity, In Vivo Toxicity, and Cellular Uptake. *Adv. Healthc. Mater.* **4** (2015) 1877-1886.
- [153] Lai, C.-H., et al., Analysis of Carbohydrate–Carbohydrate Interactions Using Sugar-Functionalized Silicon Nanoparticles for Cell Imaging. *Nano Lett.* **16** (2016) 807-811.

-
- [154] Li, Z.F. and E. Ruckenstein, Water-soluble poly(acrylic acid) grafted luminescent silicon nanoparticles and their use as fluorescent biological staining labels. *Nano Lett.* **4** (2004) 1463-1467.
- [155] Warner, J.H., et al., Water-Soluble Photoluminescent Silicon Quantum Dots. *Angewandte Chemie International Edition.* **44** (2005) 4550-4554.
- [156] Ohta, S., et al., Size- and surface chemistry-dependent intracellular localization of luminescent silicon quantum dot aggregates. *J. Mater. Chem.* **22** (2012) 10631-10638.
- [157] Ruizendaal, L., et al., Synthesis and cytotoxicity of silicon nanoparticles with covalently attached organic monolayers. *Nanotoxicology.* **3** (2009) 339-347.
- [158] Bhattacharjee, S., et al., Cytotoxicity of surface-functionalized silicon and germanium nanoparticles: the dominant role of surface charges. *Nanoscale.* **5** (2013) 4870-4883.
- [159] Bhattacharjee, S., et al., Role of surface charge and oxidative stress in cytotoxicity of organic monolayer-coated silicon nanoparticles towards macrophage NR8383 cells. *Part. Fibre Toxicol.* **7** (2010) 1-12.
- [160] Gao, F., et al., Ultrasmall [64Cu]Cu Nanoclusters for Targeting Orthotopic Lung Tumors Using Accurate Positron Emission Tomography Imaging. *ACS Nano.* **9** (2015) 4976-4986.
- [161] Zhao, Y., et al., Gold Nanoclusters Doped with 64Cu for CXCR4 Positron Emission Tomography Imaging of Breast Cancer and Metastasis. *ACS Nano.* **10** (2016) 5959-5970.
- [162] Zhao, Y., et al., Facile synthesis, pharmacokinetic and systemic clearance evaluation, and positron emission tomography cancer imaging of 64Cu–Au alloy nanoclusters. *Nanoscale.* **6** (2014) 13501-13509.
- [163] Zhou, M., et al., CuS Nanodots with Ultrahigh Efficient Renal Clearance for Positron Emission Tomography Imaging and Image-Guided Photothermal Therapy. *ACS Nano.* **9** (2015) 7085-7096.
- [164] Lewis, M.R. and R. Kannan, Development and applications of radioactive nanoparticles for imaging of biological systems. *Wiley Interdiscip. Rev. Nanomed. Nanobiotechnol.* **6** (2014) 628-640.
- [165] Erogbogbo, F., et al., *In vivo* targeted cancer imaging, sentinel lymph node mapping and multi-channel imaging with biocompatible silicon nanocrystals. *ACS Nano.* **5** (2011) 413-423.
- [166] Singh, M.P., et al., Development of Iron-Doped Silicon Nanoparticles As Bimodal Imaging Agents. *ACS Nano.* **6** (2012) 5596-5604.

-
- [167] Wu, S., et al., Biomimetic Preparation and Dual-Color Bioimaging of Fluorescent Silicon Nanoparticles. *J. Am. Chem. Soc.* **137** (2015) 14726-14732.
- [168] Qiu, M., et al., Biocompatible and biodegradable inorganic nanostructures for nanomedicine: Silicon and black phosphorus. *Nano Today*, (2019).
- [169] Dasog, M., et al., Chemical Insight into the Origin of Red and Blue Photoluminescence Arising from Freestanding Silicon Nanocrystals. *ACS Nano*. **7** (2013) 2676-2685.
- [170] Canham, L.T., Bioactive silicon structure fabrication through nanoetching techniques. *Adv. Mat.* **7** (1995) 1033-1037.
- [171] Hon, N.K., et al., Tailoring the biodegradability of porous silicon nanoparticles. *J. Biomed. Mater. Res. A*. **100A** (2012) 3416-3421.
- [172] Romero, J.J., et al., Photoluminescent 1–2 nm Sized Silicon Nanoparticles: A Surface-Dependent System. *Chem. Mater.* **25** (2013) 3488-3498.
- [173] Zhong, Y., et al., Facile, Large-Quantity Synthesis of Stable, Tunable-Color Silicon Nanoparticles and Their Application for Long-Term Cellular Imaging. *ACS Nano*. **9** (2015) 5958-5967.
- [174] Cheng, X., et al., Enhancing Quantum Dots for Bioimaging using Advanced Surface Chemistry and Advanced Optical Microscopy: Application to Silicon Quantum Dots (SiQDs). *Adv. Mat.* **27** (2015) 6144-6150.
- [175] Ji, X., et al., Highly Fluorescent, Photostable, and Ultrasmall Silicon Drug Nanocarriers for Long-Term Tumor Cell Tracking and In-Vivo Cancer Therapy. *Adv. Mat.* **27** (2015) 1029-1034.
- [176] Kharissova, O.V., et al., Ultrasmall particles and nanocomposites: state of the art. *RSC Advances*. **3** (2013) 22648-22682.
- [177] Atkins, T.M., A.Y. Louie, and S.M. Kauzlarich, An efficient microwave-assisted synthesis method for the production of water soluble amine-terminated Si nanoparticles. *Nanotechnology*. **23** (2012) 1-9.
- [178] Baldwin, R.K., et al., Room temperature solution synthesis of alkyl-capped tetrahedral shaped silicon nanocrystals. *J. Am. Chem. Soc.* **124** (2002) 1150-1151.
- [179] Cho, B., et al., Synthesis of silicon quantum dots showing high quantum efficiency. *J. Nanosci. Nanotechnol.* **14** (2014) 5868-5872.
- [180] Heath, J.R., A liquid-solution-phase synthesis of crystalline silicon. *Science*. **258** (1992) 1131-1133.
- [181] Mastronardi, M.L., et al., Preparation of monodisperse silicon nanocrystals using density gradient ultracentrifugation. *J. Am. Chem. Soc.* **133** (2011) 11928-11931.

-
- [182] Mastronardi, M.L., et al., Size-dependent absolute quantum yields for size-separated colloidally-stable silicon nanocrystals. *Nano Lett.* **12** (2012) 337-342.
- [183] Ravotto, L., et al., Bright long-lived luminescence of silicon nanocrystals sensitized by two-photon absorbing antenna. *Chem.* **2** (2017) 550-560.
- [184] Rosso-Vasic, M., et al., Alkyl-functionalized oxide-free silicon nanoparticles: synthesis and optical properties. *Small.* **4** (2008) 1835-1841.
- [185] Zhang, X., et al., A New Solution Route to Hydrogen Terminated Silicon Nanoparticles: Synthesis, Functionalization, and Water Stability. *Nanotechnology.* **18** (2007) 095601-095601.
- [186] Pujari, S.P., et al., One-Pot Gram-Scale Synthesis of Hydrogen-Terminated Silicon Nanoparticles. *Chem. Mater.* **30** (2018) 6503-6512.
- [187] Hessel, C.M., E.J. Henderson, and J.G.C. Veinot, Hydrogen Silsesquioxane: A Molecular Precursor for Nanocrystalline Si-SiO₂ Composites and Freestanding Hydride-Surface-Terminated Silicon Nanoparticles. *Chem. Mater.* **18** (2006) 6139-6146.
- [188] Henderson, E.J., J.A. Kelly, and J.G.C. Veinot, Influence of HSiO_{1.5} Sol-Gel Polymer Structure and Composition on the Size and Luminescent Properties of Silicon Nanocrystals. *Chem. Mater.* **21** (2009) 5426-5434.
- [189] Neiner, D., H.W. Chiu, and S.M. Kauzlarich, Low-Temperature Solution Route to Macroscopic Amounts of Hydrogen Terminated Silicon Nanoparticles. *J. Am. Chem. Soc.* **128** (2006) 11016-11017.
- [190] Wilcoxon, J.P., G.A. Samara, and P.N. Provencio, Optical and electronic properties of Si nanoclusters synthesized in inverse micelles. *Phys. Rev. B.* **60** (1999) 2704-2714.
- [191] Linehan, K. and H. Doyle, Size Controlled Synthesis of Silicon Nanocrystals Using Cationic Surfactant Templates. *Small.* **10** (2014) 584-590.
- [192] Yang, C.-S., et al., Synthesis of Alkyl-Terminated Silicon Nanoclusters by a Solution Route. *J. Am. Chem. Soc.* **121** (1999) 5191-5195.
- [193] Bley, R.A. and S.M. Kauzlarich, A Low-Temperature Solution Phase Route for the Synthesis of Silicon Nanoclusters. *J. Am. Chem. Soc.* **118** (1996) 12461-12462.
- [194] Toy, R., et al., Shaping cancer nanomedicine: the effect of particle shape on the in vivo journey of nanoparticles. *Nanomedicine (Lond).* **9** (2014) 121-134.
- [195] García, K.P., et al., Zwitterionic-Coated “Stealth” Nanoparticles for Biomedical Applications: Recent Advances in Countering Biomolecular Corona Formation and Uptake by the Mononuclear Phagocyte System. *Small.* **10** (2014) 2516-2529.

-
- [196] Pombo-García, K., et al., Design, synthesis, characterisation and in vitro studies of hydrophilic, colloidally stable, $^{64}\text{Cu}(\text{II})$ -labelled, ultra-small iron oxide nanoparticles in a range of human cell lines. *RSC Adv.* **3** (2013) 22443-22454.
- [197] Pant, K., et al., Surface charge and particle size determine the metabolic fate of dendritic polyglycerols. *Nanoscale.* **9** (2017) 8723-8739.
- [198] Alexis, F., et al., Factors affecting the clearance and biodistribution of polymeric nanoparticles. *Mol. Pharm.* **5** (2008) 505-515.
- [199] Iyer, A.K., et al., Exploiting the enhanced permeability and retention effect for tumor targeting. *Drug Discov. Today.* **11** (2006) 812-818.
- [200] Moghimi, S.M., A.C. Hunter, and T.L. Andresen, Factors controlling nanoparticle pharmacokinetics: an integrated analysis and perspective. *Annu. Rev. Pharmacol. Toxicol.* **52** (2012) 481-503.
- [201] Choi, H.S., et al., Renal clearance of quantum dots. *Nat. Biotechnol.* **25** (2007) 1165-1170.
- [202] Docter, D., et al., The nanoparticle biomolecule corona: lessons learned - challenge accepted? *Chem. Soc. Rev.* **44** (2015) 6094-6121.
- [203] Gunawan, C., et al., Nanoparticle-protein corona complexes govern the biological fates and functions of nanoparticles. *J. Mater. Chem. B.* **2** (2014) 2060-2083.
- [204] Lesniak, A., et al., Effects of the Presence or Absence of a Protein Corona on Silica Nanoparticle Uptake and Impact on Cells. *ACS Nano.* **6** (2012) 5845-5857.
- [205] Lundqvist, M., et al., Nanoparticle size and surface properties determine the protein corona with possible implications for biological impacts. *Proc. Natl. Acad. Sci. U.S.A.* **105** (2008) 14265-14270.
- [206] Salvati, A., et al., Experimental and theoretical comparison of intracellular import of polymeric nanoparticles and small molecules: toward models of uptake kinetics. *Nanomedicine.* **7** (2011) 818-826.
- [207] Saptarshi, S.R., A. Duschl, and A.L. Lopata, Interaction of nanoparticles with proteins: relation to bio-reactivity of the nanoparticle. *J. Nanobiotechnology.* **11** (2013) 1-12.
- [208] Tenzer, S., et al., Nanoparticle Size Is a Critical Physicochemical Determinant of the Human Blood Plasma Corona: A Comprehensive Quantitative Proteomic Analysis. *ACS Nano.* **5** (2011) 7155-7167.
- [209] Walkey, C.D., et al., Protein Corona Fingerprinting Predicts the Cellular Interaction of Gold and Silver Nanoparticles. *ACS Nano.* **8** (2014) 2439-2455.
- [210] Lee, Y.K., et al., Effect of the protein corona on nanoparticles for modulating cytotoxicity and immunotoxicity. *Int. J. Nanomedicine.* **10** (2015) 97-113.

-
- [211] Walczyk, D., et al., What the cell "sees" in bionanoscience. *J. Am. Chem. Soc.* **132** (2010) 5761-5768.
- [212] Smilowitz, H.M., et al., Biodistribution of gold nanoparticles in BBN-induced muscle-invasive bladder cancer in mice. *Int. J. Nanomedicine*. **12** (2017) 7937-7946.
- [213] Zhang, X.D., et al., Passing through the renal clearance barrier: toward ultrasmall sizes with stable ligands for potential clinical applications. *Int. J. Nanomedicine*. **9** (2014) 2069-2072.
- [214] Longmire, M., P.L. Choyke, and H. Kobayashi, Clearance Properties of Nano-sized Particles and Molecules as Imaging Agents: Considerations and Caveats. *Nanomedicine (London, England)*. **3** (2008) 703-717.
- [215] Kim, B.H., et al., Synthesis, Characterization, and Application of Ultrasmall Nanoparticles. *Chem. Mater.* **26** (2014) 59-71.
- [216] Shen, S., et al., Renal-Clearable Ultrasmall Coordination Polymer Nanodots for Chelator-Free ⁶⁴Cu-Labeling and Imaging-Guided Enhanced Radiotherapy of Cancer. *ACS Nano*. **11** (2017) 9103-9111.
- [217] Liu, J., et al., Renal clearable inorganic nanoparticles: a new frontier of bionanotechnology. *Mater. Today*. **16** (2013) 477-486.
- [218] Choi, H.S., et al., Tissue- and organ-selective biodistribution of NIR fluorescent quantum dots. *Nano. Lett.* **9** (2009) 2354-2359.
- [219] Liang, X., et al., Short- and Long-Term Tracking of Anionic Ultrasmall Nanoparticles in Kidney. *ACS Nano*. **10** (2016) 387-395.
- [220] Boselli, L., et al., Regimes of Biomolecular Ultrasmall Nanoparticle Interactions. *Angew. Chem. Int. Ed. Engl.* **56** (2017) 4215-4218.
- [221] Kang, H., et al., Renal Clearable Organic Nanocarriers for Bioimaging and Drug Delivery. *Adv Mater.* **28** (2016) 8162-8168.
- [222] Tang, S., et al., Tailoring Renal Clearance and Tumor Targeting of Ultrasmall Metal Nanoparticles with Particle Density. *Angew. Chem. Int. Ed. Engl.* **55** (2016) 16039-16043.
- [223] Truillet, C., et al., Synthesis and Characterization of (89)Zr-Labeled Ultrasmall Nanoparticles. *Mol Pharm.* **13** (2016) 2596-2601.
- [224] Sadauskas, E., et al., Kupffer cells are central in the removal of nanoparticles from the organism. *Part. Fibre Toxicol.* **4** (2007) 1-7.
- [225] Bertrand, N., et al., Cancer nanotechnology: the impact of passive and active targeting in the era of modern cancer biology. *Adv. Drug. Deliv. Rev.* **66** (2014) 2-25.

- [226] Nakamura, H., F. Jun, and H. Maeda, Development of next-generation macromolecular drugs based on the EPR effect: challenges and pitfalls. *Expert Opin. Drug Deliv.* **12** (2014) 53-64.
- [227] Viola, K.L., et al., Towards non-invasive diagnostic imaging of early-stage Alzheimer's disease. *Nat. Nanotechnol.* **10** (2015) 91-98.
- [228] Pombo Garcia, K., et al., Zwitterionic-coated "stealth" nanoparticles for biomedical applications: recent advances in countering biomolecular corona formation and uptake by the mononuclear phagocyte system. *Small.* **10** (2014) 2516-2529.
- [229] Golombek, S.K., et al., Tumor targeting via EPR: Strategies to enhance patient responses. *Adv. Drug Deliv. Rev.* **130** (2018) 17-38.
- [230] Cai, W., et al., Dual-function probe for PET and near-infrared fluorescence imaging of tumor vasculature. *J. Nucl. Med.* **48** (2007) 1862-1870.
- [231] Li, L., et al., Site-specific conjugation of monodispersed DOTA-PEGn to a thiolated diabody reveals the effect of increasing peg size on kidney clearance and tumor uptake with improved 64-copper PET imaging. *Bioconjug. Chem.* **22** (2011) 709-716.
- [232] Massa, S., et al., Sortase A-mediated site-specific labeling of camelid single-domain antibody-fragments: a versatile strategy for multiple molecular imaging modalities. *Contrast Media Mol. Imaging.* **11** (2016) 328-339.
- [233] Viehweger, K., et al., EGF receptor-targeting peptide conjugate incorporating a near-IR fluorescent dye and a novel 1,4,7-triazacyclononane-based (64)Cu(II) chelator assembled via click chemistry. *Bioconjug. Chem.* **25** (2014) 1011-1022.
- [234] Chopra, A., (111)In conjugated to benzyl-diethylenetriaminepentaacetic acid-human epidermal growth factor, in *Molecular Imaging and Contrast Agent Database (MICAD)*. 2004, National Center for Biotechnology Information (US): Bethesda (MD).
- [235] Zarschler, K., et al., High-yield production of functional soluble single-domain antibodies in the cytoplasm of Escherichia coli. *Microb. Cell. Fact.* **12** (2013) 1-13.
- [236] Anderson, C.J., et al., In vitro and in vivo evaluation of copper-64-octreotide conjugates. *J. Nucl. Med.* **36** (1995) 2315-2325.
- [237] Cihlo, J., et al., Comparison of 111In-DOTA-NOC and 111I-DOTA-TATE distribution in the target and dose-limiting tissues: conflicting results in vitro and in vivo. *Anticancer Res.* **28** (2008) 2189-2195.

- [238] Laznicek, M., et al., Octreotide and octreotate derivatives radiolabeled with yttrium: pharmacokinetics in rats. *Cancer Biother. Radiopharm.* **17** (2002) 527-533.
- [239] Lin, M., M.J. Welch, and S.E. Lapi, Effects of chelator modifications on (68)Ga-labeled [Tyr (3)]octreotide conjugates. *Mol. Imaging. Biol.* **15** (2013) 606-613.
- [240] Nedrow, J.R., et al., Positron emission tomographic imaging of copper 64- and gallium 68-labeled chelator conjugates of the somatostatin agonist tyr3-octreotate. *Mol. Imaging.* **13** (2014) 1-14.
- [241] Sprague, J.E., et al., Preparation and biological evaluation of copper-64-labeled tyr3-octreotate using a cross-bridged macrocyclic chelator. *Clin. Cancer Res.* **10** (2004) 8674-8682.
- [242] Ullrich, M., et al., Multimodal Somatostatin Receptor Theranostics Using [(64)Cu]Cu-/[(177)Lu]Lu-DOTA-(Tyr(3))octreotate and AN-238 in a Mouse Pheochromocytoma Model. *Theranostics.* **6** (2016) 650-665.
- [243] Bar-Sever, Z., et al., Guidelines on nuclear medicine imaging in neuroblastoma. *Eur. J. Nucl. Med. Mol. Imaging.* **45** (2018) 2009-2024.
- [244] Schottelius, M., M. Schwaiger, and H.-J. Wester, Rapid and high-yield solution-phase synthesis of DOTA-Tyr3-octreotide and DOTA-Tyr3-octreotate using unprotected DOTA. *Tetrahedron Lett.* **44** (2003) 2393-2396.
- [245] Comba, P., et al., Synthesis and Coordination Chemistry of Hexadentate Picolinic Acid Based Bispidine Ligands. *Inorg. Chem.* **55** (2016) 12531-12543.
- [246] Medved'ko, A.V., et al., Copper–Bispidine Complexes: Synthesis and Complex Stability Study. *ACS Omega.* **1** (2016) 854-867.
- [247] Bleiholder, C., et al., Coordination chemistry of a new rigid, hexadentate bispidine-based bis(amine)tetrakis(pyridine) ligand. *Inorg. Chem.* **44** (2005) 8145-8155.
- [248] Brox, D., et al., Cu(II)-selective bispidine-dye conjugates. *J. Inorg. Biochem.* **148** (2015) 78-83.
- [249] Gillet, R., et al., A Bispidol Chelator with a Phosphonate Pendant Arm: Synthesis, Cu(II) Complexation, and (64)Cu Labeling. *Inorg. Chem.* **56** (2017) 11738-11752.
- [250] Legdali, T., et al., Substitution-assisted stereochemical control of bispidone-based ligands. *J. Org. Chem.* **77** (2012) 11167-11176.
- [251] Bartlett, M.G. and K.L. Busch, Liquid secondary ion mass spectra of 2-iminothiolane derivatives. *Biol. Mass Spectrom.* **23** (1994) 353-356.
- [252] Lu, W., et al., Tuning the structure and function of metal–organic frameworks via linker design. *Chem. Soc. Rev.* **43** (2014) 5561-5593.

- [253] Fani, M., et al., PET of somatostatin receptor-positive tumors using ^{64}Cu - and ^{68}Ga -somatostatin antagonists: the chelate makes the difference. *J. Nucl. Med.* **52** (2011) 1110-1118.
- [254] Antunes, P., et al., Influence of different spacers on the biological profile of a DOTA-somatostatin analogue. *Bioconjug. Chem.* **18** (2007) 84-92.
- [255] Bjorkelund, H., L. Gedda, and K. Andersson, Comparing the epidermal growth factor interaction with four different cell lines: intriguing effects imply strong dependency of cellular context. *PLoS One*. **6** (2011) 1-7.
- [256] Novy, Z., et al., A comparison of in vitro methods for determining the membrane receptor expression in cell lines. *Nucl. Med. Biol.* **39** (2012) 893-896.
- [257] Miyano, M., et al., Photophysical properties of luminescent silicon nanoparticles surface-modified with organic molecules via hydrosilylation. *Photochem. Photobiol. Sci.* **15** (2016) 99-104.
- [258] Ahire, J.H., et al., Highly Luminescent and Nontoxic Amine-Capped Nanoparticles from Porous Silicon: Synthesis and Their Use in Biomedical Imaging. *ACS Appl. Mater. Interfaces*. **4** (2012) 3285-3292.
- [259] Zhong, Y., et al., Large-scale aqueous synthesis of fluorescent and biocompatible silicon nanoparticles and their use as highly photostable biological probes. *J. Am. Chem. Soc.* **135** (2013) 8350-8356.
- [260] Licciardello, N., *Design, synthesis and functionalization of ultrasmall nanoparticles for bioimaging*, in *Institut de Science et d'Ingénierie Supramoléculaires*. 2015, University of Strasbourg: Strasbourg. p. 195.
- [261] Wang, L., et al., Gram-scale synthesis of single-crystalline graphene quantum dots with superior optical properties. *Nat. Commun.* **5** (2014).
- [262] Licciardello, N., et al., Biodistribution studies of ultrasmall silicon nanoparticles and carbon dots in experimental rats and tumor mice. *Nanoscale*. **10** (2018) 9880-9891.
- [263] Dohnalova, K., et al., Surface brightens up Si quantum dots: direct bandgap-like size-tunable emission. *Light. Sci. Appl.* **2** (2013) 1-6.
- [264] Long, D.A., Infrared and Raman characteristic group frequencies. Tables and charts George Socrates John Wiley and Sons, Ltd, Chichester, Third Edition, 2001. Price £135. *J. Raman Spectrosc.* **35** (2004) 905-905.
- [265] Lillo, C.R., et al., Organic coating of 1–2-nm-size silicon nanoparticles: Effect on particle properties. *Nano Res.* **8** (2015) 2047-2062.
- [266] Frohlich, E., The role of surface charge in cellular uptake and cytotoxicity of medical nanoparticles. *Int. J. Nanomedicine*. **7** (2012) 5577-5591.

- [267] Langford, J.H., M.S. Cooper, and K.H. Orchard, Development and validation of the ^{57}Co assay for determining the ligand to antibody ratio in bifunctional chelate/antibody conjugates for use in radioimmunotherapy. *Nucl. Med. Biol.* **38** (2011) 1103-1110.
- [268] Meares, C.F., et al., Conjugation of antibodies with bifunctional chelating agents: isothiocyanate and bromoacetamide reagents, methods of analysis, and subsequent addition of metal ions. *Anal. Biochem.* **142** (1984) 68-78.
- [269] Pant, K., et al., Synthesis and biodistribution studies of (3)H- and (64)Cu-labeled dendritic polyglycerol and dendritic polyglycerol sulfate. *Bioconjug. Chem.* **26** (2015) 906-918.
- [270] Sohaebuddin, S.K., et al., Nanomaterial cytotoxicity is composition, size, and cell type dependent. *Part. Fibre Toxicol.* **7** (2010) 1-17.
- [271] Foucaud, L., et al., Measurement of reactive species production by nanoparticles prepared in biologically relevant media. *Toxicol. Lett.* **174** (2007) 1-9.
- [272] Moller, W., et al., Ultrafine particles cause cytoskeletal dysfunctions in macrophages: role of intracellular calcium. *Part. Fibre Toxicol.* **2** (2005) 7.
- [273] Jha, A., et al., Heterogeneity in Dye–TiO₂ Interactions Dictate Charge Transfer Efficiencies for Diketopyrrolopyrrole-Based Polymer Sensitized Solar Cells. *J. Phys. Chem. C.* **118** (2014) 29650-29662.
- [274] Berezin, M.Y., J. Kao, and S. Achilefu, pH-dependent optical properties of synthetic fluorescent imidazoles. *Chemistry.* **15** (2009) 3560-3566.
- [275] Szabo, A., et al., The Effect of Fluorophore Conjugation on Antibody Affinity and the Photophysical Properties of Dyes. *Biophys. J.* **114** (2018) 688-700.
- [276] Zhegalova, N.G., et al., Minimization of self-quenching fluorescence on dyes conjugated to biomolecules with multiple labeling sites via asymmetrically charged NIR fluorophores. *Contrast Media Mol. Imaging.* **9** (2014) 355-362.
- [277] Zhang, S., P. Shao, and M. Bai, In vivo type 2 cannabinoid receptor-targeted tumor optical imaging using a near infrared fluorescent probe. *Bioconjug. Chem.* **24** (2013) 1907-1916.
- [278] Hong, S., et al., Interaction of polycationic polymers with supported lipid bilayers and cells: nanoscale hole formation and enhanced membrane permeability. *Bioconjug. Chem.* **17** (2006) 728-734.
- [279] Kizlik-Masson, C., et al., 5B9, a monoclonal antiplatelet factor 4/heparin IgG with a human Fc fragment that mimics heparin-induced thrombocytopenia antibodies. *J. Thromb. Haemost.* **15** (2017) 2065-2075.

- [280] Rollin, J., et al., Characterization of 5B9, a Chimeric Monoclonal Anti-PF4/H Antibody with a Human Fc Fragment and Which Mimics the Effects of HIT Human Antibodies. *Blood*. **128** (2016) 138-138.
- [281] Arepally, G.M., et al., Characterization of a murine monoclonal antibody that mimics heparin-induced thrombocytopenia antibodies. *Blood*. **95** (2000) 1533-1540.
- [282] Borzel, H., et al., Copper-bispidine coordination chemistry: syntheses, structures, solution properties, and oxygenation reactivity. *Inorg. Chem.* **41** (2002) 5440-5452.
- [283] Huang, A.Y., et al., Concise solid-phase synthesis of inverse poly(amidoamine) dendrons using AB2 building blocks. *Chem. Commun. (Camb)*. **49** (2013) 5784-5786.
- [284] Dumas, C., R. Schibli, and P.A. Schubiger, Versatile routes to C-2- and C-6-functionalized glucose derivatives of iminodiacetic acid. *J. Org. Chem.* **68** (2003) 512-518.
- [285] Rühl, C., *Hexadentate bispidine-functionalized iron oxide nanoparticles as dual imaging agents*, in *Fakultät Mathematik und Naturwissenschaften*. 2015, Technischen Universität Dresden: Dresden. p. 129.
- [286] Polukhtin, A.C., W.; Mendoza LG., *Novel benzophenone-based chromophoric crosslinkers and reagents for incorporation of biotin or other haptens into macromolecules*, in *United States Patent Application Publication*, U.S.P.A. Publication, Editor. 2014, Polukhtin, A.
- [287] Rohrer, J., et al., Mucus permeating thiolated self-emulsifying drug delivery systems. *Eur. J. Pharm. Biopharm.* **98** (2016) 90-97.
- [288] Schlesinger, J., et al., Radiosynthesis of New [90Y]-DOTA-Based Maleimide Reagents Suitable for the Prelabeling of Thiol-Bearing I-Oligonucleotides and Peptides. *Bioconjug. Chem.* **20** (2009) 1340-1348.
- [289] Cai, Z., et al., Cu(I)-assisted click chemistry strategy for conjugation of non-protected cross-bridged macrocyclic chelators to tumour-targeting peptides. *Dalton Trans.* **44** (2015) 3945-3948.
- [290] Zhang, A.J., et al., A method for removal of N-Boc protecting groups from substrates on TFA-sensitive resins. *Tetrahedron Lett.* **39** (1998) 7439-7442.

Chapter 7 Annex

7.1 Acronyms

Ab	Antibody
A431	Epithelial cell line from epidermoid carcinoma
ADME	Absorption Distribution Metabolism Excretion
ACN	Acetonitrile
BAT	Brown adipose tissue
BFCA	Bifunctional Chelating Agent
Cyclam	1,4,8,11-Tetraazacyclotetradecane
d	Day
DCM	Dichloromethane
DIPEA	<i>N,N</i> -Diisopropylethylamine
DLS	Dynamic Light Scattering
DMEM	Dulbecco's modified Eagle's medium
DMF	<i>N,N</i> -Dimethylformamide
DMPTACN	1, 4,-Bis-(2-Pyridylmethyl)-1,4,7-triazacyclononane
DMPA	2,2-dimethoxy-2-phenylacetophenone
DMSO	Dimethyl Sulfoxide
DOTA	1,4,7,10-tetraazacyclododecane-1,4,7,10-tetraacetic acid
EDC	1-Ethyl-3-(3-dimethylaminopropyl)-carbodiimide
EDTA	Ethylenediaminetetraacetic acid
EGF	Epidermal Growth Factor
EGFR	Epidermal Growth Factor Receptor
EMA	European medicines evaluation agency
EPR	Enhanced Permeability and Retention
ESI-MS	Electrospray Ionization-Mass Spectrometry
FDA	Food and Drug Administration, USA
Fmoc	Fluorenylmethyloxycarbonyl
FTIR	Fourier transformation infrared spectroscopy
h	Hour
HCAb	Heavy chain antibodies
HD	Hydrodynamic diameter
HEK293	Human embryonic Kidney cell line
HPLC	High-Performance Liquid Chromatography
ICG	Indocyanine green
kDa	Kilo Dalton
K _d	Dissociation constant
LDH	Lactate dehydrogenase

MALDI-TOF-MS	Matrix-assisted laser desorption/ionization-Time of flight- mass spectrometry
MES	2-(N-Morpholino) ethanesulfonic acid
min	Minute (s)
MIP	Maximum Intensity Projection
MPS	Mononuclear Phagocytic System
MRI	Magnetic Resonance Imaging
MTS assay	(3-(4,5-dimethylthiazol-2-yl)-5-(3-carboxymethoxyphenyl)-2-(4-sulfophenyl)-2H-tetrazolium) assay
Mw	Weight average molecular weight
MW	Molecular Weight
MWCO	Molecular weight cut-off
NHS	N-hydroxysuccinimide
NIR	Near-infrared
NIRF	Near-infrared Fluorophores
NMR	Nuclear Magnetic Resonance
NOTA	1,4,7-Triazacyclononane-1,4,7-triacetic acid
NP	Nanoparticle
OI	Optical Imaging
PAGE	Polyacrylamide Gel electrophoresis
PAMAM	Poly(amido)amine
PBS	Phosphate buffer Saline
PET	Positron Emission Tomography
pI	Isoelectric point
<i>p.i.</i>	Post injection
PL	Photoluminescence
PLQY	Photoluminescence Quantum Yield
Py	Pyridine
QDs	Quantum Dots
QY	Quantum Yield
RCY	Radiochemical Yield
Rf	Retention Factor
RT	Room temperature
sdAb	Single domain antibody
SDS	Sodium dodecyl Sulfate
SEC	Size exclusion chromatography
SPAAC	Strain promoted alkyne-azide cycloaddition
SPECT	Single Photon Emission Computed Tomography
SST	Somatostatin
SSTR	Somatostatin Receptor
SUV	Standard Uptake Value
$t_{1/2}$	Half life
TATE	Tyr3-Octreotate peptide
TCEP	Tris(2-carboxyethyl)phosphine
TEA	Triethylamine

TEM	Transmission Electron Microscopic
TETA	1,4,8,11-Tetraazacyclotetradecane-1,4,8,11-tetraacetic acid
TFA	Trifluoroacetic Acid
TGA	Thermogravimetric Analysis
TIPS	Triisopropylsilane
TLC	Thin Layer Chromatography
R_t	Retention time
UV-Vis	Ultraviolet-Visible
WAT	White adipose tissue
λ	Wavelength
ζ	Zeta Potential
ZW	Zwitterionic

7.2 Supplementary Data

Figure 7.1: ESI-MS of B1⁺ [M+H]⁺ *m/z* 461.3 in MeOH:H₂O (1:1) v/v (left); ESI-MS of B1 [M+py+TFA]⁺ *m/z* 611.6 in ACN:H₂O (1:1) v/v (right)

Figure 7.2: ¹H NMR of B1' in CDCl₃

Figure 7.3: ¹H NMR of B1 in CDCl₃

Figure 7.4: ^1H NMR of B2 in CDCl_3 Figure 7.5: ^{13}C NMR of L1 in CDCl_3 Figure 7.6: ESI-MS of B3' $[\text{M}+\text{H}]^+$ m/z 751.8 in $\text{ACN}:\text{H}_2\text{O}$ (1:1) v/v

Figure 7.7: ESI-MS of B3 [M+H]⁺ m/z 651.4 in ACN:H₂O (1:1) v/v (left); ESI-MS of B4 [M+H]⁺ m/z 843.3 in ACN:H₂O (1:1) v/v (right)

Figure 7.8: ¹H NMR of B4 in CDCl₃

Figure 7.9: ¹H NMR of B5 in CDCl₃

Figure 7.10: ^1H NMR of B6 in CDCl_3 Figure 7.11: ^{13}C NMR of B6 in CDCl_3 Figure 7.12: ^{13}C NMR of L2 in CDCl_3

Figure 7.13: ESI-MS of B8' $[M+H]^+$ m/z 899.1 in ACN:H₂O (1:1) v/v (left); ESI-MS of B8 $[M+H]^+$ m/z 799.5 in ACN:H₂O (1:1) v/v (right)

Figure 7.14: ESI-MS of B9' $[M+H]^+$ m/z 1045.4 in ACN:H₂O (1:1) v/v

Figure 7.15: ¹H NMR of B9' in CDCl₃

Figure 7.19: ESI-MS of B11' $[M+H]^+$ m/z 489.1 in ACN:H₂O (1:1) v/v

Figure 7.20: MALDI-TOF spectra of Boc-B4-PEG-TATE with $[M+H]^+$ at m/z 2407.1.

Figure 7.21: ^1H NMR of Si NPs and trisodium citrate in D_2O .

Figure 7.22: Radiochemical purity of purified ^{64}Cu -labeled Si NP-B4 conjugate.

Radio-TLC system: iTLC SG in 2M $\text{NH}_4\text{OAc}:\text{MeOH}$ 1:1(v/v); $[^{64}\text{Cu}]\text{Cu-Si NP-B4}$ $R_f = 0$, $[^{64}\text{Cu}]\text{CuCl}_2$ $R_f = 1$.

Figure 7.23: TGA analysis of amine-terminated Si NPs.

Temperature program: Heat from 20 °C to 1200 °C @ 20 °C per minute in air)

Chapter 8 Publications, Conferences and Awards

Publications:

1. Licciardello, N., Hunoldt S; Bergmann, R.; **Singh, G.**; Mamat, M.; Faramus, A.; Ddungu, J.L.Z.; Silvestrini, S.; Maggini, M.; Cola, L. De; Stephan, H. Biodistribution studies of ultrasmall silicon nanoparticles and carbon dots in experimental rats and tumor mice. *Nanoscale* 2018, 10, 9880-9891 (Inside cover picture)

2. Joshi, T.; Kubeil, M.; Nsubuga, A.; **Singh, G.**; Gasser, G., Stephan; H. Harnessing the Coordination Chemistry of 1,4,7- Triazacyclononane for Biomimicry and Radiopharmaceutical Applications. *ChemPlusChem* 2018, 83, 554-564 (Frontpiece)

3. **Singh, G.**; Gott, M.D.; Pietzsch, H.-J.; Stephan, H. Nuclear and optical dual-labeled imaging agents. *Nuklearmedizin* 2016, 55, 41–50

Oral and Poster presentations:

1. **Singh, G.** “Cancer diagnosis by imaging agents” Science communication “FameLab Dresden” Dresden, Germany. **April 2018 (Slam/Prelims qualified)**
2. **Singh, G.** “Silicon nanoparticles as multimodal imaging agents”, Czech-HZDR Workshop: Future Collaborations in Diagnostic Cancer Imaging and Cancer Therapy, HZDR, Dresden. **April 2018 (Oral)**
3. **Singh, G.**; Ddungu, J.; Licciardello, N.; Hunoldt, S.; Bergmann, R.; Faramus, A.; Cola, L. De; Stephan, H. “Silicon nanoparticle based dual-labeled imaging agents”, Science communication Workshop: NextGen@Helmholtz, Kiel, Germany. **July 2017 (Slam and Poster)**
4. **Singh, G.**; Licciardello, N.; Hunoldt, S.; Bergmann, R.; Faramus, A.; Cola, L. De; Stephan, H. “Modified ultrasmall nanoparticles as multimodal imaging agents for biomedical applications”, International Symposium on Radiopharmaceutical Sciences (ISRS), Dresden, Germany. **May 2017 (Poster)**
5. **Singh, G.**; Licciardello, N.; Hunoldt, S.; Bergmann, R.; Faramus, A.; Cola, L. De; Stephan, H. “Surface modified ultrasmall nanoparticles as dual-labeled imaging agents”, Immuno-imaging and molecular therapy workshop, Brussels, Belgium. **April 17 (Poster)**
6. **Singh, G.** “Bispidines as novel PET imaging agents”, Annual Meeting: Nanotracking, **October 2016 (Oral)**
7. **Singh, G.** “Synthesis, characterization and functionalization of Bispidines”, Annual Meeting: NanoTracking-Monash, Dresden, Germany. **September 2016 (Oral)**
8. **Singh, G.**; Licciardello, N.; Zarschler, K.; Faramus, A.; Cola, L. De; Stephan, H. “Surface modified ultrasmall nanoparticles as dual labelled imaging agents” International Conference: VI European Association for Chemical and Molecular Sciences (EuCheMS), Seville, Spain. **September 2016 (Poster)**
9. **Singh, G.**; Licciardello, N.; Faramus, A.; Cola, L. De; Stephan, H. “Synthesis and characterization of ultrasmall nanoparticles for biomedical applications” International Conference: Analytica, Munich, Germany. **May 16 (Poster)**
10. **Singh, G.** “Synthesis and characterization of modified ultrasmall nanoparticles for biomedical imaging”, Meeting: Nanotracking meets Portfolio, Koenigstein, Germany. March 2016 **(Oral)**
11. **Singh, G.**; Hunoldt, S.; Faramus, A.; Licciardello, N.; Cola, L. De; Stephan, H. “Synthesis and characterization of ultrasmall nanoparticles for biomedical applications” International Conference “II International Symposium on

Nanomaterials and Applications”, Caparica, Portugal. January 16 (**Shotgun and Poster**)

12. **Singh, G.** “Synthesis and characterization of ultrasmall nanoparticles for biomedical applications”, Annual Meeting: NanoTracking, **October 2015 (Oral)**

Awards:

1. DAAD-RISE Fellowship 2018 to host and supervise Ms. N. Patel (Williams College, USA) **June 2018 – August 2018**
2. DAAD-RISE Fellowship 2017 to host and supervise Mr. J. Marsh (Memorial University, Canada) **May 2017 – July 2017**
3. Poster award finalist at the “ICMI Workshop: Immuno-Imaging and Molecular Therapy”, Brussels, Belgium. **April 2017**
4. Best slammer at the “Dresden concept Open Air Science Slam”, Dresden, Germany. **July 2016**
5. DAAD-RISE Fellowship 2016 to host and supervise Ms. A. Fjodor (University of Leicester, U.K.) **June 2016 – August 2016**
6. GDCh Travel award to attend the conference “Analytica”, Munich, Germany. **May 2016**
7. Best poster presentation at the International Conference “II International Symposium on Nanoparticles / Nanomaterials and Applications”, Caparica, Portugal. **January 2016**

PhD research stay:

September 2015

Institute of Supramolecular Science and Engineering (*ISIS*), University of Strasbourg, Strasbourg, France and Karlsruhe Institute of Technology (KIT), Karlsruhe, Germany.

Advisor: Prof. Luisa de Cola (*ISIS*, France / *KIT*, Germany)

Chapter 9 Acknowledgements

My PhD journey has been a period of intense learning to broaden my scientific profile and a series of new experiences. I would like to take the opportunity to thank the people who guided, supported, and helped me during this period.

I am very grateful to Prof. Jörg Steinbach for allowing me to be a part of his research institute as a PhD candidate. His constant support throughout the years, despite his busy schedule is commendable. Thank you once again for the PhD seminars and Christmas dinners.

The next most important person during my stay at HZDR was Dr. Holger Stephan. As a supervisor, he guided me throughout, making time for all the discussions and possibilities regarding the projects. Apart from the lab, he was also a great motivation for my participation at various scientific events, slams and the art exhibition. Thank you Dr. Stephan for everything.

I would like to express my gratitude to Prof. Luisa De Cola and her group members, Dr. Angelique Faramus and John Ddungu for hosting and assisting me in the labs at KIT, Karlsruhe and ISIS, Strasbourg. It was a great learning experience for me.

The long stay at HZDR could not have been any easier and fun; thanks to my awesome colleagues who became good friends over time. Kritee, thank you for everything, the scientific discussions and suggestions, the motivating conversations and the midnight parties.

Another person who I have to thank for being extremely helpful, patient and motivating is Tanmaya. Thank you for discussing the research, instant solution to all the problems and guiding me towards the end with the thesis. It really means a lot. Thank you so much. A special thanks to Dr. Martin Walther and Dr. Constantin Mamat for performing NMR and discussion over the results. I would also like to extend my thanks to Anne Nsubuga and Dr. Kristof Zarschler for keeping their cool, making the work environment stress-free with their wit. Thanks you Kristof for all the scientific help and explanations. Thanks to Dr. Nadia Licciardello, a dear friend who has been a great support system in and outside HZDR. I would like to thank Dr. Massimo Sgarzi for helping me with photochemistry and proofreading of the thesis. Thanks to Dr. Karina-Garcia for being there as a supportive friend and helping me towards the end. Big thanks to David Bauer, Sandy Weißflog, Moore Gilpin, Falco Reissig and all colleagues at HZDR who helped me in all small and big things.

I would deeply thank the technical staff at HZDR for making my PhD life easy. Thank you Mrs. Grosse for all the support in the lab, for the conversations and making me feel a little less homesick. Thank you Madlen Matterna for acquainting with me from my very

first day at HZDR, Karin Landrock for conducting the experiments for me, Bianka Kriesel for the urgent lyophilizations and FTIR measurements, Utta Herzog for the cell studies, Christian Jentschel for LC-MS analysis, Ulrike Gesche for the managing the radionuclide production and Marita Kersten for managing all my documents and planning my business trips. Big thanks to my students, Alysia, Jevon and Neena for being excellent assistance and teaching me how to supervise. I would also appreciate the DAAD-RISE projects for the successful research exchange programs and the Helmholtz Virtual Institute, 'NanoTracking' for the project funding.

I would also like to acknowledge Edith Reschke and Sara Müller from the Library for all the help and motivation for setting up my first art exhibition at HZDR.

Thanks to the friends, I made during the course of PhD who comforted and motivated me throughout. Thank you Dr. Matthias Streller for all the car rides, Thomas Brenner for solving the IT issues in no time, Dr. Matthew Gott for being a scientific support; Swati, Cem, Nilesh, Olga, Andreas, Ajay and Stefan for all the moral support, motivation and all the good times we had together. A big hug to my dear friend Vandana for always being there for me. Thank you so much. Thanks to Devadas, Jay, Karishma and Sumbul who were distant but always encouraging me for the best.

I would like to thank Prateek for being by my side all this while and motivating me towards achieving this milestone.

Last, but not least, I would thank my big family, for all the unconditional love, support and trust in me. It makes me push my limits and prove myself each time. We grew together over the years, your curiosity to know about the new culture and lifestyle and letting me adapt to it so well. Not many families are this "cool". A special thanks to my grandfather, I hold your sincerity in me and to my grandmother, who taught me to be a little more crazy and calm; to my parents, for being the best; my sister Honey, for being my everything; my cousin Rajat, you are an inspiration; Aunt and Uncle for treating me the best. Your sacrifices, your attitude towards education and your thoughtfulness, all makes me so proud. Sincere thanks.

Chapter 10 Statement of declaration / Erklärungen

Eidesstattliche Erklärung

Die vorliegende Arbeit ist am Helmholtz-Zentrum Dresden-Rossendorf im Institut für Radiopharmazeutische Krebsforschung von Juli 2015 bis Dezember 2018 unter der wissenschaftlichen Betreuung von Herrn Dr. Holger Stephan und Herrn Prof. Dr. Jörg Steinbach angefertigt worden.

Bisherige erfolglose Promotionsverfahren haben nicht stattgefunden.

Dresden, den

Versicherung

Hiermit versichere ich, dass ich die vorliegende Arbeit ohne unzulässige Hilfe Dritter und ohne Benutzung anderer als der angegebenen Hilfsmittel angefertigt habe; die aus fremden Quellen direkt oder indirekt übernommenen Gedanken sind als solche kenntlich gemacht. Die Arbeit wurde bisher weder im Inland noch im Ausland in gleicher oder ähnlicher Form einer anderen Prüfungsbehörde vorgelegt.

Dresden, den