**Dieses Dokument ist eine Zweitveröffentlichung (Verlagsversion) /
This is a self-archiving document (published version):**

S. Guloksuz, M. van Nierop, R. Lieb, R. van Winkel, H.-U. Wittchen, J. van Os

Evidence that the presence of psychosis in nonpsychotic disorder is environment-dependent and mediated by severity of non-psychotic psychopathology

Erstveröffentlichung in / First published in:

Psychological Medicine. 2015, 45(11), S. 2389-2401 [Zugriff am: 19.03.2020]. Cambridge University Press. ISSN 1469-8978.

DOI: <https://doi.org/10.1017/S0033291715000380>

Diese Version ist verfügbar / This version is available on:

<https://nbn-resolving.org/urn:nbn:de:bsz:14-qucosa2-390353>

„Dieser Beitrag ist mit Zustimmung des Rechteinhabers aufgrund einer (DFGgeförderten) Allianz- bzw. Nationallizenz frei zugänglich.“

This publication is openly accessible with the permission of the copyright owner. The permission is granted within a nationwide license, supported by the German Research Foundation (abbr. in German DFG).

www.nationallizenzen.de/

Evidence that the presence of psychosis in non-psychotic disorder is environment-dependent and mediated by severity of non-psychotic psychopathology

S. Guloksuz^{1,2}, M. van Nierop¹, R. Lieb³, R. van Winkel^{1,4}, H.-U. Wittchen^{5,6} and J. van Os^{1,7*}

¹ Department of Psychiatry and Psychology, Maastricht University Medical Centre, EUIRON, Maastricht, The Netherlands

² Department of Psychiatry, Yale University School of Medicine, New Haven, CT, USA

³ Epidemiology and Health Psychology, Institute of Psychology, University of Basel, Basel, Switzerland

⁴ Department of Psychiatry, University Psychiatric Center KU Leuven, KU Leuven, Belgium

⁵ Institute of Clinical Psychology and Psychotherapy, Technical University Dresden, Dresden, Germany

⁶ Max Planck Institute of Psychiatry, Munich, Germany

⁷ Department of Psychosis Studies, King's College London, King's Health Partners, Institute of Psychiatry, London, UK

Background. Evidence suggests that in affective, non-psychotic disorders: (i) environmental exposures increase risk of subthreshold psychotic experiences (PEs) and strengthen connectivity between domains of affective and subthreshold psychotic psychopathology; and (ii) PEs are a marker of illness severity.

Method. In 3021 adolescents from the Early Developmental Stages of Psychopathology cohort, we tested whether the association between PEs and presence of DSM-IV mood disorder (MD)/obsessive-compulsive disorder (OCD) would be moderated by risk factors for psychosis (cannabis use, childhood trauma and urbanicity), using the interaction contrast ratio (ICR) method. Furthermore, we analysed whether the interaction between environment and PEs was mediated by non-psychotic psychopathology.

Results. The association between PEs and MD/OCD was moderated by urbanicity (ICR = 2.46, $p = 0.005$), cannabis use (ICR = 3.76, $p = 0.010$) and, suggestively, trauma (ICR = 1.91, $p = 0.063$). Exposure to more than one environmental risk factor increased the likelihood of co-expression of PEs in a dose-response fashion. Moderating effects of environmental exposures were largely mediated by the severity of general non-psychotic psychopathology (percentage explained 56–68%, all $p < 0.001$). Within individuals with MD/OCD, the association between PEs and help-seeking behaviour, as an index of severity, was moderated by trauma (ICR = 1.87, $p = 0.009$) and urbanicity (ICR = 1.48, $p = 0.005$), but not by cannabis use.

Conclusions. In non-psychotic disorder, environmental factors increase the likelihood of psychosis admixture and help-seeking behaviour through an increase in general psychopathology. The findings are compatible with a relational model of psychopathology in which more severe clinical states are the result of environment-induced disturbances spreading through a psychopathology network.

Received 23 September 2014; Revised 4 February 2015; Accepted 4 February 2015; First published online 25 March 2015

Key words: Affective disorders, cannabis, childhood trauma, psychosis, risk factors, urbanicity.

Introduction

Recent evidence suggests that psychotic experiences are not confined to psychotic disorders like schizophrenia and delusional disorder, but are also common, at subthreshold level of severity, in non-psychotic disorders (Hanssen *et al.* 2003; Rossler *et al.* 2011; van Rossum *et al.* 2011; Varghese *et al.* 2011; Kelleher *et al.* 2012a, b; Wigman *et al.* 2012; DeVlylder *et al.* 2014). These findings suggest that psychotic

experiences can be conceptualized along a dimension of psychosis expression, ranging from no/low to high levels. Research on the incidence and mechanisms of psychosis thus may profit from adopting a transdiagnostic approach applied to both frequent non-psychotic and rare traditional psychotic disorders. A number of findings on the association between psychotic experiences and non-psychotic mental disorder illustrate this issue. First, the risk of co-occurring psychotic experiences in non-psychotic disorder is moderated by proxy environmental and genetic risk factors; second, co-occurring psychotic experiences are associated with greater levels of non-psychotic illness severity and poorer treatment response (Kaymaz *et al.* 2007, 2012; Craddock & Owen, 2010; Perlis *et al.*

* Address for correspondence: J. van Os, Department of Psychiatry and Neuropsychology, Maastricht University Medical Centre, EUIRON, PO Box 616, 6200 MD Maastricht, The Netherlands.
(Email: vanosj@gmail.com)

2011; Rossler et al. 2011; Wigman et al. 2011a, b; Kelleher et al. 2012a, b; Wigman et al. 2012; Cross-Disorder Group of the Psychiatric Genomics Consortium, 2013). Third, environmental impact on psychotic experiences is accompanied by increased levels of connectivity between symptoms within, e.g. delusions and hallucinations (Smeets et al. 2012, 2014), and across, e.g. affective and psychotic symptoms (Kaymaz et al. 2006; Wigman et al. 2012), the domains of psychotic and non-psychotic psychopathology, particularly between psychotic experiences on the one hand and depressive, anxiety and manic symptoms on the other (van Rossum et al. 2011; Van Nierop et al. *in press*). Fourth, in those with an at-risk mental state, non-psychotic psychopathology (e.g. mood/anxiety disorders) is common (Fusar-Poli et al. 2014) and precedes transition to psychotic disorder (Rietdijk et al. 2011).

In combination, these findings suggest that in non-psychotic disorder, environmental risk factors make psychopathology more complex, 'co-morbid', or connected, resulting in the occurrence of psychotic experiences in addition to non-psychotic symptoms, which in turn make a negative impact on treatment response and prognosis. These results echo findings in neuroscience that increased connectivity in variably defined networks is associated with increased probability of clinical transitions (Treserras et al. 2009; Baliki et al. 2012; Varotto et al. 2012; Wigman et al. 2013). There is evidence that psychopathology can be usefully represented as a network of mutually impacting symptoms (Borsboom & Cramer, 2013). In the network, causally linked sets of symptoms reciprocally make an impact on each other over time to progress toward a more distinct syndrome forged through the pattern of the dynamic network. For instance, persecutory delusions may provoke anxiety under certain circumstances (e.g. being in public). Anxiety may in turn lead to misinterpretation of perceptual experience due to increased alertness, giving rise to hallucinatory experiences. Hallucinatory experiences in turn may provoke confusion requiring a cognitive explanation – often odd and delusional –, thus creating a loop reinforcing the primary delusion.

There exists some evidence that suggests that environmental risk factors may make symptoms firmly interconnected, which in turn – depending on the degree and the pattern of connectivity between symptoms – give rise to a more complex, 'co-morbid', and distinct clinical syndrome that may further transition to a more severe mental state with a poorer prognosis as the connectedness expands (Wigman et al. 2012; van Os, 2013; Smeets et al. 2014; van Os et al. 2014). Although a formal test of this model would require an intensive time series of measures of psychopathology after environmental impact, a more general

and practical approach is to test to what degree the association between environmental exposure and the occurrence of subthreshold psychotic experiences in non-psychotic mental disorder is mediated by 'connected' non-psychotic psychopathology. Indeed, recent research suggests that the association between environmental risk and psychotic symptoms is mediated by non-psychotic psychopathology (Gracie et al. 2007; Freeman & Fowler, 2009; Reeves et al. 2014).

In the current paper, we combined the above evidence to formulate several specific hypotheses. First, we hypothesized that the association between psychotic experiences and non-psychotic disorder would be moderated by exposure to selected known environmental risk factors (i.e. urbanicity, trauma and cannabis use). Second, we hypothesized that subthreshold psychosis admixture in non-psychotic disorder associated with environmental exposure would be mediated by an increase in severity of non-psychotic psychopathology. Third, in order to examine the impact of environment-associated psychosis admixture on clinical severity, we hypothesized that, within the group of individuals diagnosed with non-psychotic disorder, the association between co-occurring psychotic experiences and help-seeking would also be moderated by exposure to known environmental risk factors. To date, these hypotheses have either not been examined before, or not been examined jointly, or not been examined in a sample of individuals with non-psychotic disorders. The aim thus was to examine, in a sample of individuals with non-psychotic disorder, to what degree environmental exposures make psychopathology more 'psychotic' – and more likely to result in contact with mental health services – and to what degree this may be driven by an underlying mechanism of increasing severity of general non-psychotic psychopathology.

Method

Data were derived from the Early Developmental Stages of Psychopathology (EDSP) Study, which collected data on the prevalence, incidence, risk factors, co-morbidity and course of mental disorders in a random representative population sample of 3021 adolescents and young adults in the general population. Following ethics committee approval, a representative population sample was randomly drawn from the 1994 German government population registers. The sample consisted of adolescents and young adults living in the Munich area aged 14–24 years at baseline. The overall design of the study was observational, longitudinal and prospective, consisting of a baseline (T0: $n=3021$, response rate 71%) and three follow-ups [T1: $n=1228$ (younger group only), response rate=88%; T2: $n=$

2548, response rate = 84%; T3: $n = 2210$, response rate = 73%], covering a time period of on average 1.6 years (T0–T1, $s.d. = 0.2$), 3.5 years (T0–T2, $s.d. = 0.3$) and 8.4 years (T0–T3, range 7.3–10.5 years, $s.d. = 0.7$), respectively. The sample at T1 only included the younger members of the cohort; assessments at T0, T2 and T3 were based on the full sample. For the current analyses, data from all waves were used (T0, T1, T2 and T3). A more detailed description of the study, fieldwork and response rates and characteristics of the respondents has been reported elsewhere (Wittchen *et al.* 1998b; Lieb *et al.* 2000; Zimmermann *et al.* 2008).

Instruments

Psychopathology

Symptoms, syndromes and disorders were assessed with the computer-assisted version of the Munich-Composite International Diagnostic Interview (DIA-X/M-CIDI) (Wittchen & Pfister, 1997), an updated and expanded version of the World Health Organization's CIDI version 1.2. The DIA-X/M-CIDI is a comprehensive, fully standardized, diagnostic interview, addressing symptoms, syndromes and diagnoses of a wide range of mental disorders in accordance with definitions and criteria of the Diagnostic and Statistical Manual of Mental Disorders (DSM), fourth edition and the International Classification of Diseases, tenth edition. The DIA-X/M-CIDI has been shown to be both reliable and valid (Reed *et al.* 1998; Wittchen *et al.* 1998a). Interviews were conducted by fully trained and experienced psychologists, who were allowed to probe with follow-up questions. This interview technique is particularly relevant for the assessment of psychotic symptoms, which are sensitive to false-positive ratings. The EDSP study covers a total observation period of up to 10 years. At T0 (baseline), the lifetime version of the DIA-X/M-CIDI was used; for subsequent waves, the respective DIA-X/M-CIDI interval versions were used.

Consistent with DSM-5 (APA, 2013), we studied psychotic experiences beyond disorders listed in the chapter 'Schizophrenia spectrum and other psychotic disorders'. Thus, according to DSM-5 (APA, 2013), a number of other disorders can be accompanied by psychotic features, namely: depression spectrum (depressive episode/dysthymia – persistent depressive disorder in DSM-5); mania spectrum (manic episode, hypomanic episode); and obsessive-compulsive disorder. Based on the relevant sections, a dichotomous variable (hereafter: 'affective spectrum disorder') for each time point was constructed representing whether an individual had been diagnosed with one or more of these disorders.

At all time points, participants also completed the Self-Report Symptom Checklist-90-R (SCL-90-R), a

comprehensive self-report symptom inventory, multi-dimensional in nature, and oriented to screen for a broad range of psychological problems and psychopathology in community respondents and respondents with somatic and psychiatric disorders. It contains 90 items, scored on a five-point severity scale, measuring nine primary symptom dimensions named 'somatization', 'obsessive-compulsive', 'interpersonal sensitivity', 'depression', 'anxiety', 'hostility', 'phobic anxiety', 'paranoid ideation' and 'psychoticism'. Reliability and validity of the SCL-90-R were established previously (Derogatis & Cleary, 1977). The timeframe is the past 2 weeks.

For the purpose of the analyses, a dichotomous psychotic experience variable was created using a cut-off point to define the group of individuals with the highest 10% scores of combined psychosis and paranoia dimensions of the SCL-90-R (hereafter 'psychosis expression'), consistent with previous analyses in this sample (Henquet *et al.* 2005; Cougnard *et al.* 2007; Dominguez *et al.* 2011) and congruent with the meta-analytic rate for the prevalence of psychotic symptoms in the general population (Linscott & van Os, 2013).

A 'non-psychotic psychopathology load' variable was created using the total score of the SCL-90-R excluding the psychotic dimensions ('paranoid ideation' and 'psychoticism'). This score thus reflects the number and the severity of SCL-90-R non-psychotic symptoms.

Trauma

Self-reported lifetime (baseline) and interval (follow-up) exposure to trauma was assessed using the N-section of the DIA-X/M-CIDI on trauma and post-traumatic stress disorder comprising nine groups of specific traumatic events (presented by a respondent list) such as 'experienced physical threat', 'experienced serious accident', or 'being sexually abused as a child'. Visual presentation of the list allowed respondents and interviewers to avoid speaking about sometimes embarrassing and stigmatizing trauma by simply indicating the number of the event. Consistent with earlier analyses (Spauwen *et al.* 2006; Wigman *et al.* 2012), positive responses to any of the events were coded as 'self-reported trauma'.

Cannabis use

Cannabis use was assessed with the L-section of the DIA-X/M-CIDI using the question 'Have you ever used cannabis five times or more?' to define cannabis exposure. Conforming to previous work (Henquet *et al.* 2005; Kuepper *et al.* 2011c), the DIA-X/M-CIDI cut-off of use of five times or more was used to

define a binary variable for cannabis exposure. Therefore, at baseline, cannabis use was defined as lifetime use of cannabis of five times or more. At each consequent time point, cannabis use was defined as interval use of cannabis of five times or more, such that period between baseline and T1 defines cannabis use variable at T1, period between T1 and T2 defines cannabis use at T2, and so on.

Urbanicity

Consistent with previous work (Spauwen *et al.* 2004; Wigman *et al.* 2012), urbanicity was defined as living, at baseline, in the urban region of the German city of Munich *versus* the surrounding areas of Munich. The urban area, hence defined, had a population density of 4061 persons per square mile; for the rural area, this was 553 persons per square mile.

Help-seeking behaviour

In line with previous analyses reported elsewhere (Wigman *et al.* 2012), help-seeking behaviour was defined as general help-seeking behaviour, which was broadly defined as having visited any mental health institution ever for any mental health problem (based on the Q-section of the DIA-X/M-CIDI), assessed at baseline and each follow-up.

Statistical analysis

All analyses were carried out with Stata 13.1 (StataCorp, 2013). Given the fact that outcome was measured at each time point, in keeping with previous work (Dominguez *et al.* 2011; Wigman *et al.* 2012), data were analysed cross-sectionally in the 'long format' (each individual contributing four observations: T0, T1, T2 and T3). Logistic regression models using the LOGIT command were applied to analyse whether the association between psychosis expression and affective spectrum disorder was greater if there was also evidence of exposure to environmental risk factors (trauma, urbanicity, cannabis use). In order to correct for the clustering of multiple observations within subjects, cluster-robust standard errors were computed, using the CLUSTER option. We tested for departure from additivity using the interaction contrast ratio (ICR) method as suggested by Knol and colleagues (Schwartz & Susser, 2006; Knol *et al.* 2007), using the NLCOM command in Stata.

A convincing case exists for additive models to provide the best representation of synergy (Rothman *et al.* 1980; Schwartz & Susser, 2006) and that they are the most useful from a public health perspective (Darroch, 1997; Kendler & Gardner, 2010). This approach allows use of odds ratios (ORs) derived from logistic models

to estimate the relative excess risk as a result of synergy for combinations of dichotomous, ordinal and continuous exposures (i.e. $ICR = OR_{\text{exposure A \& exposure B}} - OR_{\text{exposure A only}} - OR_{\text{exposure B only}} + 1$). An ICR greater than zero is defined as a positive deviation from additivity.

To test our hypotheses on synergism, we entered the four exposure states occasioned by the combination of each environmental factor (trauma, urbanicity, cannabis use) and psychosis expression as independent variables (three dummy variables with non-exposed state as the reference category), and affective spectrum disorder as the dependent variable in logistic models (Knol *et al.* 2007). Using the ORs derived from these models, ICRs (e.g. $ICR = OR_{\text{trauma \& psychosis expression}} - OR_{\text{trauma}} - OR_{\text{psychosis expression}} + 1$) for each model were calculated using the Stata NLCOM command.

In the second part of the analysis, non-psychotic psychopathology load was added as an independent variable in the logistic models to test to what degree synergism between environmental factors and psychosis expression were mediated by the severity of non-psychotic psychopathology. Additionally, using the KHB command in Stata, we formally tested whether non-psychotic psychopathology load mediated the interaction between environmental factors and psychosis expression. The KHB command provides unbiased decomposition of the total effect into direct and indirect effects (Karlson & Holm, 2011; Kohler *et al.* 2011).

Logistic regression models using the same strategy were applied to analyse whether the associations between psychosis expression and help-seeking behaviour within the group diagnosed with affective spectrum disorder was greater if there was also evidence of exposure to environmental risk factors (trauma, urbanicity, cannabis use).

Results

Characteristics of the study sample at different time points are shown in Table 1.

There was a suggestion that the association between psychosis expression and affective spectrum disorder was greater if there was also evidence of trauma exposure (Fig. 1a). The OR for those with psychosis expression and exposure to trauma was 7.78, in comparison with ORs of 1.56 for those exposed to trauma only, and 5.31 for those with psychosis expression only, yielding an ICR of 1.91 ($p = 0.063$). The ICR was mediated by non-psychotic psychopathology load: the estimated difference between the full model including non-psychotic psychopathology load and the reduced model without non-psychotic psychopathology load was large [OR difference 3.81, 95%

Table 1. Characteristics of the population at different time points

	T0	T1	T2	T3
Sex, male	1533 (50.74)	637 (51.87)	1297 (50.90)	1135 (51.36)
Mean age, years (s.d.)	18.26 (3.34)	16.72 (1.19)	21.74 (3.39)	26.62 (3.47)
Affective spectrum disorder	451 (14.93)	132 (4.37)	309 (10.23)	318 (10.53)
Major depressive episode	326 (10.79)	86 (7.01)	233 (9.21)	238 (10.77)
Dysthymia	58 (1.92)	19 (1.55)	28 (1.11)	36 (1.63)
Hypomanic episode	55 (1.82)	16 (1.30)	48 (1.88)	16 (0.72)
Manic episode	41 (1.36)	16 (1.30)	20 (0.78)	12 (0.54)
Obsessive-compulsive disorder	20 (0.66)	3 (0.24)	12 (0.47)	22 (1.00)
Psychosis expression ^a	280 (9.28)	105 (8.66)	208 (8.28)	183 (8.38)
Mean non-psychotic psychopathology load (s.d.) ^b	1.39 (0.34)	1.29 (0.30)	1.28 (0.28)	1.25 (0.28)
Urbanicity	2162 (71.57)	860 (70.03)	1796 (70.49)	1558 (70.50)
Cannabis use	390 (13.38)	167 (13.92)	519 (20.89)	574 (26.50)
Trauma	598 (19.79)	201 (16.38)	1350 (53.49)	1382 (62.56)
Help-seeking behaviour ^c	120 (26.61)	19 (14.39)	93 (30.10)	137 (43.08)

Data are given as number (percentage) unless otherwise indicated.

T0, Baseline; T1, first follow-up; T2, second follow-up; T3, third follow-up; s.d., standard deviation; SCL, Self-Report Symptom Checklist.

^a Individuals with the highest 10% of scores of the SCL-psychosis subscale.

^b The total score of the SCL-90-R excluding the psychotic dimensions ('paranoid ideation' and 'psychoticism').

^c In the subsample consisted only of individuals diagnosed with affective spectrum disorder.

confidence interval (CI) 3.03–4.79] and the percentage of total effect explained by non-psychotic psychopathology load was 62.2% (Table 2).

Similarly, the association between psychosis expression and affective spectrum disorder was greater if there was also evidence of exposure to an urban environment (Fig. 1b). The OR for those living in an urban area along with psychosis expression was 6.55, in comparison with ORs of 1.12 for those living in an urban area only, and 3.97 for those with psychosis expression only, yielding an ICR of 2.46 ($p=0.005$). The ICR was mediated by non-psychotic psychopathology load: the estimated difference between the full model including non-psychotic psychopathology load and the reduced model without non-psychotic psychopathology load was large (OR difference 3.76, 95% CI 2.99–4.73) and the percentage of total effect explained by non-psychotic psychopathology load was 68.1% (Table 2).

The association between psychosis expression and affective spectrum disorder was moderated by cannabis use (Fig. 1c). The ORs for those exposed to cannabis use along with psychosis expression was 9.47, in comparison with ORs of 1.77 for those exposed to cannabis use only, and 4.94 for those with psychosis expression, yielding an ICR of 3.76 ($p=0.010$). The ICR was mediated by non-psychotic psychopathology load: the estimated difference between the full model including non-psychotic psychopathology load and the reduced model without non-psychotic psychopathology load was large (OR difference 3.73, 95% CI 2.96–4.71) and

the percentage of total effect explained by non-psychotic psychopathology load was 56.1% (Table 2).

Analysing all environmental factors (cannabis, urbanicity, trauma), combined together as a loading variable, with psychosis expression, revealed a dose-response relationship in the level of association between psychosis expression and affective spectrum disorder as a function of the degree of environmental exposure (Fig. 2).

In the subsample of individuals with affective spectrum disorder, psychosis expression combined with trauma (ICR=1.87, 95% CI 0.47–3.27, $p=0.009$), and psychosis expression combined with urbanicity (ICR=1.48, 95% CI 0.45–2.51, $p=0.005$) in a synergistic fashion to increase the odds of help-seeking behaviour (Table 3). There was no synergy between psychosis expression and cannabis use in the model of help-seeking behaviour (ICR=−0.01, 95% CI −1.41 to 1.39, $p=0.988$).

Discussion

This study investigated to what degree expression of psychotic experiences in affective spectrum disorder (mood disorders and obsessive-compulsive disorder) is contingent on exposure to environmental factors known to be associated with psychotic disorders (trauma, urbanicity, cannabis use) (van Os *et al.* 2010), using the ICR method, and to what degree expression of psychotic experiences may be mediated by general severity of psychopathology. The principal

Fig. 1. Association between affective spectrum disorder and psychosis expression as a function of environmental exposures (cannabis, trauma and urbanicity). Unexposed, Exposed to neither environmental risk nor psychosis; ICR, Interaction contrast ratio; OR, odds ratio; CI, confidence interval.

findings were: (i) psychosis expression in affective spectrum disorder additively increased with exposure to environmental risk factors (trauma, urbanicity, cannabis), and this interaction was largely mediated by general non-psychotic psychopathology load; (ii) environmental effects appeared to make an impact on the same mechanism as evidenced by the dose-response relationship between affective spectrum disorder and psychosis expression, as a function of the extent of environmental exposure; (iii) the likelihood of help-seeking behaviour, as a proxy for

severity/suffering, as a function of psychosis expression within the group diagnosed with affective spectrum disorder was moderated by both trauma and urbanicity, but not by cannabis use.

Environment and transdiagnostic expression of psychosis

At the phenomenological level, indicators of reality distortion such as hallucinations and delusions were previously thought to be core symptoms that

Table 2. Combined effect size of psychosis expression and environmental exposure in model of affective spectrum disorder

	Unadjusted			Adjusted for non-psychotic psychopathology load			Mediation effect of non-psychotic psychopathology load			
	ICR	(95% CI)	<i>p</i>	ICR	(95% CI)	<i>p</i>	OR, difference ^a	(95% CI)	Indirect effect, % ^b	<i>p</i>
Trauma	1.91	(−0.10 to 3.93)	0.063	0.19	(−0.45 to 0.83)	0.557	3.81	(3.03 to 4.79)	62.2	<0.001
Urbanicity	2.46	(0.73 to 4.18)	0.005	0.64	(0.11 to 1.17)	0.018	3.76	(2.99 to 4.73)	68.1	<0.001
Cannabis use	3.76	(0.89 to 6.63)	0.010	0.74	(−0.18 to 1.67)	0.116	3.73	(2.96 to 4.71)	56.1	<0.001

ICR, Interaction contrast ratio; CI, confidence interval; OR, odds ratio.

^a Estimated difference between the full model including non-psychotic psychopathology load and the reduced model without non-psychotic psychopathology load.

^b Percentage of total effect explained by non-psychotic psychopathology load.

Fig. 2. Cumulative effect size of psychosis expression (PE) and environmental exposure (EE) combination in model of affective spectrum disorder. Unexposed, Exposed to neither environmental risk nor psychosis; OR, odds ratio; CI, confidence interval. * $p < 0.001$, ** $p = 0.01$.

distinguish schizophrenia from non-psychotic mental disorders *per se*. However, the findings, combined with previous work, suggest that affective syndromes and reality distortion may not only simply co-occur, but also reciprocally make an impact on each other to predict subsequent clinical outcome (Kaymaz *et al.* 2007, 2012; Rossler *et al.* 2011; Wigman *et al.* 2011a, b; Kelleher *et al.* 2012a, b; M. van Nierop, M Bak, R De Graaf, M Ten Have, S van Dorsselaer, GROUP Investigators and R van Winkel, unpublished observations), across the boundaries of traditional diagnostic constructs.

We provide evidence that the likelihood of psychosis expression in non-psychotic affective spectrum disorder was moderated by environmental exposure in a

dose–response fashion. A growing body of evidence indicates that environmental risk factors play a role in the aetiology of psychotic syndromes (van Os *et al.* 2010). Exposure to an urban environment (Vassos *et al.* 2012), early childhood trauma (Varese *et al.* 2012) and cannabis use (Minozzi *et al.* 2010) are consistently associated with psychotic outcome in rigorously designed epidemiological studies, including prospective studies, in both clinical and general populations. Confirming the shared vulnerability theory of psychopathology, environmental factors, particularly cannabis use and trauma, are also associated with affective disorders, albeit less pronounced (Bovasso, 2001; Moore *et al.* 2007; Breetvelt *et al.* 2010; Nanni *et al.* 2012; Kedzior & Laeber, 2014). Similarly, a nationwide

Table 3. Combined effect of psychosis expression and environmental exposure on help-seeking behaviour

	Odds ratio	(95% confidence interval)	<i>p</i>
Unexposed	1.00		
Trauma	1.41	(1.04–1.92)	0.027
Psychosis	1.45	(0.95–2.21)	0.085
Trauma + psychosis expression	3.73	(2.51–5.54)	<0.001
Unexposed	1.00		
Urbanicity	1.20	(0.85–1.70)	0.302
Psychosis	1.17	(0.63–2.16)	0.626
Urbanicity + psychosis expression	2.85	(1.91–4.26)	<0.001
Unexposed	1.00		
Cannabis	1.74	(1.25–2.44)	0.001
Psychosis	2.22	(1.56–3.14)	<0.001
Cannabis + psychosis expression	2.95	(1.91–4.56)	<0.001

Swedish follow-up study showed that a higher level of urbanization predicted psychosis and, to a lesser degree, depression (Sundquist *et al.* 2004).

The fact that likelihood of expression of psychosis in affective spectrum disorder was moderated by the amount of environmental exposure in a dose–response fashion may imply that environmental factors make an impact on the same underlying mechanism, reinforcing each other. A similar additive effect across multiple environmental exposures was previously shown on the likelihood of persistence of psychotic experiences in the general population over time (Cougnard *et al.* 2007). Additionally, we demonstrated that psychosis expression and environmental exposure (urbanicity, trauma, but not cannabis use) synergistically increased odds of help-seeking behaviour, reflecting onset of treatment needs, within the group of individuals diagnosed with affective spectrum disorder. The findings suggest that the cumulative environmental risk load moderated the level of psychosis admixture in non-psychotic disorder, increasing the likelihood of help-seeking behaviour. To date, several studies, with one exception (Kuepper *et al.* 2011a), have demonstrated that environmental risk factors may add to each other's effects in the causation of psychotic outcomes (Houston *et al.* 2008; Harley *et al.* 2010; Kuepper *et al.* 2011b; Konings *et al.* 2012; Morgan *et al.* 2014). The cumulative effect of environmental factors predicting worse outcome appears to be evident across diagnostic categories; for example additive effects between childhood trauma and cannabis use on earlier

age at onset, increased rapid cycling, and suicide attempts were also demonstrated in bipolar disorder (Aas *et al.* 2014).

Environmental impact as increased connectivity spreading through the psychopathology network

It has been suggested that symptoms of psychopathology do not vary in isolation, but rather make an impact on each other over time. For example, there is evidence that affective dysregulation underlies psychosis (Freeman *et al.* 2013a, b; Hartley *et al.* 2013), that motivational impairments may predict psychosis (Dominguez *et al.* 2010) and that abnormal perceptions may make an impact on risk of delusional ideation (Smeets *et al.* 2012). In non-psychotic affective disorder, expression of psychosis is associated with severity of anxiety and depression (Armando *et al.* 2013), and there is evidence that the association between environmental risk factors such as cannabis and trauma on the one hand, and psychosis on the other, is mediated by non-psychotic symptoms such as anxiety and depression (Gracie *et al.* 2007; Freeman & Fowler, 2009; Reeves *et al.* 2014). This literature is compatible with a relational model of psychopathology, with environmental factors making an impact on symptoms, and symptoms making an impact on each other in causal networks (Borsboom & Cramer, 2013; van Os, 2013). The current findings are in agreement with the literature, in that environmental risk factors increased the probability of psychosis admixture in non-psychotic disorder, and that non-psychotic psychopathology mediated this relationship. In combination, the data suggest that a relational model of psychopathology in which the environment makes an impact on connectivity and 'tips' psychopathology towards increased levels of psychotic experiences may be useful in clinical practice, informing on severity and onset of help-seeking behaviour. However, prospective studies are required in order to examine the temporal relationship between environment and symptoms, as well as between different symptoms connecting with and making an impact on each other.

Methodological issues

One of the major strengths of this study was the standardized assessment of a large, representative population through clinical interviews conducted by psychologists who were allowed to probe with clinical questioning. The sample was followed up over 8 years with three post-baseline assessments, thus facilitating the interpretation of interplay between dimensions of psychopathology and risk factors. Nevertheless, more frequent clinical assessments could model the dynamic trajectory of psychopathology over time. Furthermore, our cross-sectional analyses, combining observations

from all four time points in the 'long format', positively make an impact on reliability but provide little evidence for causality, with the exception of dose-response relationships. However, to our knowledge, there has been no prospective study with sufficiently intensive repeated assessments of psychopathology and environmental exposures over a long period of time, spanning the period of transition from adolescence to adulthood – the critical period for emerging psychopathology. At this stage, notwithstanding its limitations, a more practical approach – such as the cross-sectional analysis performed in the present study – may be used to test to what degree psychotic experiences and help-seeking behaviour in non-psychotic disorder are contingent on exposure to environmental risks and its impact on the severity of general psychopathology. Individuals with more severe psychopathology may have a tendency to report more exposure to risk factors, such as childhood trauma (van Winkel *et al.* 2013). There are also several other determinants that play critically important roles in mediating the impact of environmental exposure on shaping the subsequent psychopathology, such as the extent and the timing of environmental exposure. Evidence indicates that the more severe the environmental exposure, the more severe the psychopathology. For instance, psychosis may be associated with childhood abuse stronger than neglect (Read *et al.* 2005; Heins *et al.* 2011; van Dam *et al.* 2014); likewise, the risk for psychosis cumulatively increases by the amount and the duration of cannabis exposure (Moore *et al.* 2007; Manrique-Garcia *et al.* 2012). Moreover, any dose-response effect is evident across diagnostic categories, such that both childhood trauma (Leverich *et al.* 2002; Hammersley *et al.* 2003; Hovens *et al.* 2010, 2012; van Dam *et al.* 2014) and cannabis use (Aas *et al.* 2014) are associated with more psychotic episodes, poor outcome, and treatment resistance in mood and anxiety disorders. Further, environmental exposures may have the greatest impact on psychopathology during the critically sensitive period from childhood to early adolescence, during which neurodevelopment still continues. The timing of the exposure is an influential factor within this sensitive period (Fisher *et al.* 2010; Bentall *et al.* 2012; van Winkel & Kuepper, 2014). In the present study, we only analysed environmental exposures coded binary as either present or not present; therefore our analysis did not allow us for further interpretation of distinctive impacts of the extent (e.g. sexual abuse *v.* physical abuse), the timing (e.g. childhood *v.* adolescence) and the amount of exposure (e.g. heavy cannabis use *v.* occasional cannabis use) on psychopathology.

We tested models of interaction where the variables making up the interaction (psychosis expression and

environmental exposure) were not independent (Linscott & van Os, 2013), making it difficult to distinguish between mediation (environmental risk causing psychosis expression in affective spectrum disorder) and moderation (environmental risk moderating the strength of the association between psychosis expression and affective spectrum disorder). However, both models, particularly the test of mediation by underlying severity of psychopathology, are conceptually converging (psychotic forms of non-psychotic disorder are more likely in the presence of environmental risks) and of similar clinical interest.

Our findings further reinforce the notion that psychopathology may represent a network of interacting symptom dimensions, which are affected by environmental risk factors across traditional diagnoses, potentially adding to diagnosis using a system of environmental stratification.

Acknowledgements

This work is part of the EDSP Study and is funded by the German Federal Ministry of Education and Research (BMBF) project no. 01EB9405/6, 01EB9910/6, EB10106200, 01EB0140 and 01EB0440. Part of the fieldwork and analyses were also additionally supported by grants of the Deutsche Forschungsgemeinschaft (DFG) LA1148/1-1, WI2246/7-1 and WI709/8-1. Principal investigators of the EDSP Study are H.-U. Wittchen and R. Lieb. Core staff members of the EDSP group are: K. von Sydow, G. Lachner, A. Perkonig, P. Schuster, M. Höfler, H. Sonntag, T. Brückl, E. Garczynski, B. Isensee, A. Nocon, C. Nelson, H. Pfister, V. Reed, B. Spiegel, A. Schreier, U. Wunderlich, P. Zimmermann, D. Beesdo and A. Bitner. Scientific advisors are J. Angst (Zurich), J. Margraf (Basel), G. Esser (Potsdam), K. Merikangas (National Institute of Mental Health, Bethesda), R. Kessler (Harvard, Boston) and J. van Os (Maastricht). The research leading to these results has received funding from the European Community's Seventh Framework Program under grant agreement no. HEALTH-F2-2009-241909 (project EU-GEI).

Declaration of Interest

None.

References

- Aas M, Etain B, Bellivier F, Henry C, Lagerberg T, Ringen A, Agartz I, Gard S, Kahn JP, Leboyer M, Andreassen OA, Melle I (2014). Additive effects of childhood abuse and cannabis abuse on clinical expressions of bipolar disorders. *Psychological Medicine* 44, 1653–1662.

- APA (2013). *Diagnostic and Statistical Manual of Mental Disorders*, 5th edn. American Psychiatric Publishing: Arlington, VA.
- Armando M, Lin A, Girardi P, Righetti V, Dario C, Saba R, Decrescenzo F, Mazzone L, Vicari S, Birchwood M, Fiori Nastro P (2013). Prevalence of psychotic-like experiences in young adults with social anxiety disorder and correlation with affective dysregulation. *Journal of Nervous and Mental Disease* **201**, 1053–1059.
- Baliki MN, Petre B, Torbey S, Herrmann KM, Huang L, Schnitzer TJ, Fields HL, Apkarian AV (2012). Corticostriatal functional connectivity predicts transition to chronic back pain. *Nature Neuroscience* **15**, 1117–1119.
- Bentall RP, Wickham S, Shevlin M, Varese F (2012). Do specific early-life adversities lead to specific symptoms of psychosis? A study from the 2007 Adult Psychiatric Morbidity Survey. *Schizophrenia Bulletin* **38**, 734–740.
- Borsboom D, Cramer AO (2013). Network analysis: an integrative approach to the structure of psychopathology. *Annual Review of Clinical Psychology* **9**, 91–121.
- Bovasso GB (2001). Cannabis abuse as a risk factor for depressive symptoms. *American Journal of Psychiatry* **158**, 2033–2037.
- Breetvelt EJ, Boks MP, Numans ME, Selten JP, Sommer IE, Grobbee DE, Kahn RS, Geerlings MI (2010). Schizophrenia risk factors constitute general risk factors for psychiatric symptoms in the population. *Schizophrenia Research* **120**, 184–190.
- Cognard A, Marcelis M, Myin-Germeys I, De Graaf R, Vollebergh W, Krabbendam L, Lieb R, Wittchen HU, Henquet C, Spauwen J, Van Os J (2007). Does normal developmental expression of psychosis combine with environmental risk to cause persistence of psychosis? A psychosis proneness–persistence model. *Psychological Medicine* **37**, 513–527.
- Craddock N, Owen MJ (2010). The Kraepelinian dichotomy – going, going ... but still not gone. *British Journal of Psychiatry* **196**, 92–95.
- Cross-Disorder Group of the Psychiatric Genomics Consortium (2013). Identification of risk loci with shared effects on five major psychiatric disorders: a genome-wide analysis. *Lancet* **381**, 1371–1379.
- Darroch J (1997). Biologic synergism and parallelism. *American Journal of Epidemiology* **145**, 661–668.
- Derogatis LR, Cleary PA (1977). Confirmation of the dimensional structure of the SCL-90: a study in construct validation. *Journal of Clinical Psychology* **33**, 981–989.
- DeVylder JE, Burnette D, Yang LH (2014). Co-occurrence of psychotic experiences and common mental health conditions across four racially and ethnically diverse population samples. *Psychological Medicine* **44**, 3503–3513.
- Dominguez MD, Saka MC, Lieb R, Wittchen HU, van Os J (2010). Early expression of negative/disorganized symptoms predicting psychotic experiences and subsequent clinical psychosis: a 10-year study. *American Journal of Psychiatry* **167**, 1075–1082.
- Dominguez MD, Wichers M, Lieb R, Wittchen HU, van Os J (2011). Evidence that onset of clinical psychosis is an outcome of progressively more persistent subclinical psychotic experiences: an 8-year cohort study. *Schizophrenia Bulletin* **37**, 84–93.
- Fisher HL, Jones PB, Fearon P, Craig TK, Dazzan P, Morgan K, Hutchinson G, Doody GA, McGuffin P, Leff J, Murray RM, Morgan C (2010). The varying impact of type, timing and frequency of exposure to childhood adversity on its association with adult psychotic disorder. *Psychological Medicine* **40**, 1967–1978.
- Freeman D, Dunn G, Fowler D, Bebbington P, Kuipers E, Emsley R, Jolley S, Garety P (2013a). Current paranoid thinking in patients with delusions: the presence of cognitive–affective biases. *Schizophrenia Bulletin* **39**, 1281–1287.
- Freeman D, Fowler D (2009). Routes to psychotic symptoms: trauma, anxiety and psychosis-like experiences. *Psychiatry Research* **169**, 107–112.
- Freeman D, Startup H, Dunn G, Cernis E, Wingham G, Pugh K, Cordwell J, Kingdon D (2013b). The interaction of affective with psychotic processes: a test of the effects of worrying on working memory, jumping to conclusions, and anomalies of experience in patients with persecutory delusions. *Journal of Psychiatric Research* **47**, 1837–1842.
- Fusar-Poli P, Nelson B, Valmaggia L, Yung AR, McGuire PK (2014). Comorbid depressive and anxiety disorders in 509 individuals with an at-risk mental state: impact on psychopathology and transition to psychosis. *Schizophrenia Bulletin* **40**, 120–131.
- Gracie A, Freeman D, Green S, Garety PA, Kuipers E, Hardy A, Ray K, Dunn G, Bebbington P, Fowler D (2007). The association between traumatic experience, paranoia and hallucinations: a test of the predictions of psychological models. *Acta Psychiatrica Scandinavica* **116**, 280–289.
- Hammersley P, Dias A, Todd G, Bowen-Jones K, Reilly B, Bentall RP (2003). Childhood trauma and hallucinations in bipolar affective disorder: preliminary investigation. *British Journal of Psychiatry* **182**, 543–547.
- Hanssen M, Peeters F, Krabbendam L, Radstake S, Verdoux H, van Os J (2003). How psychotic are individuals with non-psychotic disorders? *Social Psychiatry and Psychiatric Epidemiology* **38**, 149–154.
- Harley M, Kelleher I, Clarke M, Lynch F, Arseneault L, Connor D, Fitzpatrick C, Cannon M (2010). Cannabis use and childhood trauma interact additively to increase the risk of psychotic symptoms in adolescence. *Psychological Medicine* **40**, 1627–1634.
- Hartley S, Barrowclough C, Haddock G (2013). Anxiety and depression in psychosis: a systematic review of associations with positive psychotic symptoms. *Acta Psychiatrica Scandinavica* **128**, 327–346.
- Heins M, Simons C, Lataster T, Pfeifer S, Versmissen D, Lardinois M, Marcelis M, Delespaul P, Krabbendam L, van Os J, Myin-Germeys I (2011). Childhood trauma and psychosis: a case–control and case–sibling comparison across different levels of genetic liability, psychopathology, and type of trauma. *American Journal of Psychiatry* **168**, 1286–1294.
- Henquet C, Krabbendam L, Spauwen J, Kaplan C, Lieb R, Wittchen HU, van Os J (2005). Prospective cohort study of

- cannabis use, predisposition for psychosis, and psychotic symptoms in young people. *BMJ* 330, 11.
- Houston JE, Murphy J, Adamson G, Stringer M, Shevlin M (2008). Childhood sexual abuse, early cannabis use, and psychosis: testing an interaction model based on the National Comorbidity Survey. *Schizophrenia Bulletin* 34, 580–585.
- Hovens JG, Giltay EJ, Wiersma JE, Spinhoven P, Penninx BW, Zitman FG (2012). Impact of childhood life events and trauma on the course of depressive and anxiety disorders. *Acta Psychiatrica Scandinavica* 126, 198–207.
- Hovens JG, Wiersma JE, Giltay EJ, van Oppen P, Spinhoven P, Penninx BW, Zitman FG (2010). Childhood life events and childhood trauma in adult patients with depressive, anxiety and comorbid disorders *vs.* controls. *Acta Psychiatrica Scandinavica* 122, 66–74.
- Karlson KB, Holm A (2011). Decomposing primary and secondary effects: a new decomposition method. *Research in Social Stratification and Mobility* 29, 221–237.
- Kaymaz N, Drukker M, Lieb R, Wittchen HU, Werbeloff N, Weiser M, Lataster T, van Os J (2012). Do subthreshold psychotic experiences predict clinical outcomes in unselected non-help-seeking population-based samples? A systematic review and meta-analysis, enriched with new results. *Psychological Medicine* 42, 2239–2253.
- Kaymaz N, Krabbendam L, de Graaf R, Nolen W, Ten Have M, van Os J (2006). Evidence that the urban environment specifically impacts on the psychotic but not the affective dimension of bipolar disorder. *Social Psychiatry and Psychiatric Epidemiology* 41, 679–685.
- Kaymaz N, van Os J, de Graaf R, Ten Have M, Nolen W, Krabbendam L (2007). The impact of subclinical psychosis on the transition from subclinical mania to bipolar disorder. *Journal of Affective Disorders* 98, 55–64.
- Kedzior KK, Laeber LT (2014). A positive association between anxiety disorders and cannabis use or cannabis use disorders in the general population – a meta-analysis of 31 studies. *BMC Psychiatry* 14, 136.
- Kelleher I, Keeley H, Corcoran P, Lynch F, Fitzpatrick C, Devlin N, Molloy C, Roddy S, Clarke MC, Harley M, Arseneault L, Wasserman C, Carli V, Sarchiapone M, Hoven C, Wasserman D, Cannon M (2012a). Clinicopathological significance of psychotic experiences in non-psychotic young people: evidence from four population-based studies. *British Journal of Psychiatry* 201, 26–32.
- Kelleher I, Murtagh A, Molloy C, Roddy S, Clarke MC, Harley M, Cannon M (2012b). Identification and characterization of prodromal risk syndromes in young adolescents in the community: a population-based clinical interview study. *Schizophrenia Bulletin* 38, 239–246.
- Kendler KS, Gardner CO (2010). Interpretation of interactions: guide for the perplexed. *British Journal of Psychiatry* 197, 170–171.
- Knol MJ, van der Tweel I, Grobbee DE, Numans ME, Geerlings MI (2007). Estimating interaction on an additive scale between continuous determinants in a logistic regression model. *International Journal of Epidemiology* 36, 1111–1118.
- Kohler U, Karlson KB, Holm A (2011). Comparing coefficients of nested nonlinear probability models. *Stata Journal* 11, 420–438.
- Konings M, Stefanis N, Kuepper R, de Graaf R, ten Have M, van Os J, Bakoula C, Henquet C (2012). Replication in two independent population-based samples that childhood maltreatment and cannabis use synergistically impact on psychosis risk. *Psychological Medicine* 42, 149–159.
- Kuepper R, Henquet C, Lieb R, Wittchen HU, van Os J (2011a). Non-replication of interaction between cannabis use and trauma in predicting psychosis. *Schizophrenia Research* 131, 262–263.
- Kuepper R, van Os J, Lieb R, Wittchen HU, Henquet C (2011b). Do cannabis and urbanicity co-participate in causing psychosis? Evidence from a 10-year follow-up cohort study. *Psychological Medicine* 41, 2121–2129.
- Kuepper R, van Os J, Lieb R, Wittchen HU, Hoffer M, Henquet C (2011c). Continued cannabis use and risk of incidence and persistence of psychotic symptoms: 10 year follow-up cohort study. *BMJ* 342, d738.
- Leverich GS, McElroy SL, Suppes T, Keck PE Jr, Denicoff KD, Nolen WA, Altshuler LL, Rush AJ, Kupka R, Frye MA, Autio KA, Post RM (2002). Early physical and sexual abuse associated with an adverse course of bipolar illness. *Biological Psychiatry* 51, 288–297.
- Lieb R, Isensee B, von Sydow K, Wittchen HU (2000). The Early Developmental Stages of Psychopathology Study (EDSP): a methodological update. *European Addiction Research* 6, 170–182.
- Linscott RJ, van Os J (2013). An updated and conservative systematic review and meta-analysis of epidemiological evidence on psychotic experiences in children and adults: on the pathway from proneness to persistence to dimensional expression across mental disorders. *Psychological Medicine* 43, 1133–1149.
- Manrique-Garcia E, Zammit S, Dalman C, Hemmingsson T, Andreasson S, Allebeck P (2012). Cannabis, schizophrenia and other non-affective psychoses: 35 years of follow-up of a population-based cohort. *Psychological Medicine* 42, 1321–1328.
- Minozzi S, Davoli M, Bargagli AM, Amato L, Vecchi S, Perucci CA (2010). An overview of systematic reviews on cannabis and psychosis: discussing apparently conflicting results. *Drug Alcohol Review* 29, 304–317.
- Moore THM, Zammit S, Lingford-Hughes A, Barnes TRE, Jones PB, Burke M, Lewis G (2007). Cannabis use and risk of psychotic or affective mental health outcomes: a systematic review. *Lancet* 370, 319–328.
- Morgan C, Reininghaus U, Reichenberg A, Frissa S, SELCoH study team, Hotopf M, Hatch SL (2014). Adversity, cannabis use and psychotic experiences: evidence of cumulative and synergistic effects. *British Journal of Psychiatry* 204, 346–353.
- Nanni V, Uher R, Danese A (2012). Childhood maltreatment predicts unfavorable course of illness and treatment outcome in depression: a meta-analysis. *American Journal of Psychiatry* 169, 141–151.
- Perlis RH, Uher R, Ostacher M, Goldberg JF, Trivedi MH, Rush AJ, Fava M (2011). Association between bipolar

- spectrum features and treatment outcomes in outpatients with major depressive disorder. *Archives of General Psychiatry* 68, 351–360.
- Read J, van Os J, Morrison AP, Ross CA (2005). Childhood trauma, psychosis and schizophrenia: a literature review with theoretical and clinical implications. *Acta Psychiatrica Scandinavica* 112, 330–350.
- Reed V, Gander F, Pfister H, Steiger A, Sonntag H, Trenkwalder C, Hundt W, Wittchen H-U (1998). To what degree does the Composite International Diagnostic Interview (CIDI) correctly identify DSM-IV disorders? Testing validity issues in a clinical sample. *International Journal of Methods in Psychiatric Research* 7, 142–155.
- Reeves LE, Anglin DM, Heimberg RG, Gibson LE, Fineberg AM, Maxwell SD, Kerns CM, Ellman LM (2014). Anxiety mediates the association between cannabis use and attenuated positive psychotic symptoms. *Psychiatry Research* 218, 180–186.
- Rietdijk J, Hogerzeil SJ, van Hemert AM, Cuijpers P, Linszen DH, van der Gaag M (2011). Pathways to psychosis: help-seeking behavior in the prodromal phase. *Schizophrenia Research* 132, 213–219.
- Rosler W, Hengartner MP, Ajdacic-Gross V, Haker H, Gamma A, Angst J (2011). Sub-clinical psychosis symptoms in young adults are risk factors for subsequent common mental disorders. *Schizophrenia Research* 131, 18–23.
- Rothman KJ, Greenland S, Walker AM (1980). Concepts of interaction. *American Journal of Epidemiology* 112, 467–470.
- Schwartz S, Susser E (2006). Relationships among causes. In *Psychiatric Epidemiology: Searching for the Causes of Mental Disorders* (ed. E. Susser, S. Schwartz, A. Morabia and E. Bromet). Oxford and New York: Oxford University Press.
- Smeets F, Lataster T, Dominguez MD, Hommes J, Lieb R, Wittchen HU, van Os J (2012). Evidence that onset of psychosis in the population reflects early hallucinatory experiences that through environmental risks and affective dysregulation become complicated by delusions. *Schizophrenia Bulletin* 38, 531–542.
- Smeets F, Lataster T, Viechtbauer W, Delespaul P, G.R.O.U.P. (2014). Evidence that environmental and genetic risks for psychotic disorder may operate by impacting on connections between core symptoms of perceptual alteration and delusional ideation. *Schizophrenia Bulletin*. Published online 12 September 2014. doi:10.1093/schbul/sbu122.
- Spauwen J, Krabbendam L, Lieb R, Wittchen HU, van Os J (2004). Does urbanicity shift the population expression of psychosis? *Journal of Psychiatric Research* 38, 613–618.
- Spauwen J, Krabbendam L, Lieb R, Wittchen HU, van Os J (2006). Impact of psychological trauma on the development of psychotic symptoms: relationship with psychosis proneness. *British Journal of Psychiatry* 188, 527–533.
- StataCorp (2013). *Stata/SE Statistical Software, Release 13.1*. StataCorp LP: College Station, TX.
- Sundquist K, Frank G, Sundquist J (2004). Urbanisation and incidence of psychosis and depression: follow-up study of 4.4 million women and men in Sweden. *British Journal of Psychiatry* 184, 293–298.
- Treserras S, Boulanouar K, Conchou F, Simonetta-Moreau M, Berry I, Celsis P, Chollet F, Loubinoux I (2009). Transition from rest to movement: brain correlates revealed by functional connectivity. *NeuroImage* 48, 207–216.
- van Dam DS, van Nierop M, Viechtbauer W, Velthorst E, van Winkel R, Genetic Risk and Outcome of Psychosis (GROUP) investigators, Bruggeman R, Cahn W, de Haan L, Kahn RS, Meijer CJ, Myin-Germeys I, van Os J, Wiersma D (2014). Childhood abuse and neglect in relation to the presence and persistence of psychotic and depressive symptomatology. *Psychological Medicine*. Published online 17 July 2014. doi:10.1017/S0033291714001561.
- Van Nierop M, Viechtbauer W, Gunther N, van Zelst C, de Graaf R, ten Have M, van Dorsselaer S, Bak M, G.R.O.U.P. Investigators, van Winkel R (in press). Childhood trauma is associated with a specific admixture of affective, anxiety, and psychotic symptoms cutting across traditional diagnostic boundaries. *Psychological Medicine*.
- van Os J (2013). The dynamics of subthreshold psychopathology: implications for diagnosis and treatment. *American Journal of Psychiatry* 170, 695–698.
- van Os J, Kenis G, Rutten BP (2010). The environment and schizophrenia. *Nature* 468, 203–212.
- van Os J, Lataster T, Delespaul P, Wichers M, Myin-Germeys I (2014). Evidence that a psychopathology interactome has diagnostic value, predicting clinical needs: an experience sampling study. *PLOS ONE* 9, e86652.
- van Rossum I, Dominguez MD, Lieb R, Wittchen HU, van Os J (2011). Affective dysregulation and reality distortion: a 10-year prospective study of their association and clinical relevance. *Schizophrenia Bulletin* 37, 561–571.
- van Winkel R, Kuepper R (2014). Epidemiological, neurobiological, and genetic clues to the mechanisms linking cannabis use to risk for nonaffective psychosis. *Annual Review of Clinical Psychology* 10, 767–791.
- van Winkel R, van Nierop M, Myin-Germeys I, van Os J (2013). Childhood trauma as a cause of psychosis: linking genes, psychology, and biology. *Canadian Journal of Psychiatry. Revue Canadienne de Psychiatrie* 58, 44–51.
- Varese F, Smeets F, Drukker M, Lieveer R, Lataster T, Viechtbauer W, Read J, van Os J, Bentall RP (2012). Childhood adversities increase the risk of psychosis: a meta-analysis of patient-control, prospective- and cross-sectional cohort studies. *Schizophrenia Bulletin* 38, 661–671.
- Varghese D, Scott J, Welham J, Bor W, Najman J, O'Callaghan M, Williams G, McGrath J (2011). Psychotic-like experiences in major depression and anxiety disorders: a population-based survey in young adults. *Schizophrenia Bulletin* 37, 389–393.
- Varotto G, Visani E, Canafoglia L, Franceschetti S, Avanzini G, Panzica F (2012). Enhanced frontocentral EEG connectivity in photosensitive generalized epilepsies: a partial directed coherence study. *Epilepsia* 53, 359–367.
- Vassos E, Pedersen CB, Murray RM, Collier DA, Lewis CM (2012). Meta-analysis of the association of urbanicity with schizophrenia. *Schizophrenia Bulletin* 38, 1118–1123.
- Wigman JT, van Nierop M, Vollebergh WA, Lieb R, Beesdo-Baum K, Wittchen HU, van Os J (2012). Evidence that psychotic symptoms are prevalent in disorders of anxiety and depression, impacting on illness onset, risk,

- and severity – implications for diagnosis and ultra-high risk research. *Schizophrenia Bulletin* **38**, 247–257.
- Wigman JT, van Os J, Thiery E, Derom C, Collip D, Jacobs N, Wichers M** (2013). Psychiatric diagnosis revisited: towards a system of staging and profiling combining nomothetic and idiographic parameters of momentary mental States. *PLOS ONE* **8**, e59559.
- Wigman JT, van Winkel R, Raaijmakers QA, Ormel J, Verhulst FC, Reijneveld SA, van Os J, Vollebergh WA** (2011a). Evidence for a persistent, environment-dependent and deteriorating subtype of subclinical psychotic experiences: a 6-year longitudinal general population study. *Psychological Medicine* **41**, 2317–2329.
- Wigman JT, Vollebergh WA, Raaijmakers QA, Iedema J, van Dorsselaer S, Ormel J, Verhulst FC, van Os J** (2011b). The structure of the extended psychosis phenotype in early adolescence – a cross-sample replication. *Schizophrenia Bulletin* **37**, 850–860.
- Wittchen HU, Lachner G, Wunderlich U, Pfister H** (1998a). Test–retest reliability of the computerized DSM-IV version of the Munich-Composite International Diagnostic Interview (M-CIDI). *Social Psychiatry and Psychiatric Epidemiology* **33**, 568–578.
- Wittchen HU, Perkonig A, Lachner G, Nelson CB** (1998b). Early Developmental Stages of Psychopathology Study (EDSP): objectives and design. *European Addiction Research* **4**, 18–27.
- Wittchen HU, Pfister H** (1997). *DIA-X-Interviews: Manual für Screening-Verfahren und Interview; Interviewheft Langsschnittuntersuchung (DIA-X-Lifetime); Ergänzungsheft (DIA-X-Lifetime); Interviewheft Querschnittuntersuchung (DIA-X-Monatsversion); Ergänzungsheft (DIA-X-12 Monatsversion); PC-Programm zur Durchführung der Interviews (Langsund Querschnittuntersuchung). Auswertungsprogramm*. Swets & Zeitlinger: Frankfurt, Germany.
- Zimmermann P, Bruckl T, Lieb R, Nocon A, Ising M, Beesdo K, Wittchen HU** (2008). The interplay of familial depression liability and adverse events in predicting the first onset of depression during a 10-year follow-up. *Biological Psychiatry* **63**, 406–414.