Dieses Dokument ist eine Zweitveröffentlichung (Verlagsversion) /
This is a self-archiving document (published version):

W. X. Chmielewski, N. Wolff, M. Mückschel, V. Roessner, C. Beste

**Effects of multisensory integration processes on response inhibition
in adolescent autism spectrum disorder**

Erstveröffentlichung in / First published in:

Psychological Medicine. 2016, 46 (13), S. 2705– 2716 [Zugriff am: 15.04.2020]. Cambridge University Press. ISSN 1469-8978.

DOI: <https://doi.org/10.1017/S0033291716001008>

Diese Version ist verfügbar / This version is available on:

<https://nbn-resolving.org/urn:nbn:de:bsz:14-qucosa2-706813>

„Dieser Beitrag ist mit Zustimmung des Rechteinhabers aufgrund einer (DFGgeförderten) Allianz- bzw. Nationallizenz frei zugänglich.“

This publication is openly accessible with the permission of the copyright owner. The permission is granted within a nationwide license, supported by the German Research Foundation (abbr. in German DFG).
www.nationallizenzen.de/

Effects of multisensory integration processes on response inhibition in adolescent autism spectrum disorder

W. X. Chmielewski¹, N. Wolff¹, M. Mückschel¹, V. Roessner¹ and C. Beste^{1,2*}

¹Cognitive Neurophysiology, Department of Child and Adolescent Psychiatry, Faculty of Medicine of the TU Dresden, Germany

²Experimental Neurobiology, National Institute of Mental Health, Klecany, Czech Republic

Background. In everyday life it is often required to integrate multisensory input to successfully conduct response inhibition (RI) and thus major executive control processes. Both RI and multisensory processes have been suggested to be altered in autism spectrum disorder (ASD). It is, however, unclear which neurophysiological processes relate to changes in RI in ASD and in how far these processes are affected by possible multisensory integration deficits in ASD.

Method. Combining high-density EEG recordings with source localization analyses, we examined a group of adolescent ASD patients ($n = 20$) and healthy controls ($n = 20$) using a novel RI task.

Results. Compared to controls, RI processes are generally compromised in adolescent ASD. This aggravation of RI processes is modulated by the content of multisensory information. The neurophysiological data suggest that deficits in ASD emerge in attentional selection and resource allocation processes related to occipito-parietal and middle frontal regions. Most importantly, conflict monitoring subprocesses during RI were specifically modulated by content of multisensory information in the superior frontal gyrus.

Conclusions. RI processes are overstrained in adolescent ASD, especially when conflicting multisensory information has to be integrated to perform RI. It seems that the content of multisensory input is important to consider in ASD and its effects on cognitive control processes.

Received 2 December 2015; Revised 31 March 2016; Accepted 14 April 2016; First published online 18 July 2016

Key words: Autism spectrum disorder, EEG, executive functions, multisensory processing, response inhibition.

Introduction

Autism spectrum disorder (ASD) is a major neurodevelopmental disorder (APA, 2013). While social communication and interaction deficits, as well as stereotypic behaviour and interests, represent the core problem of ASD, alterations in behavioural and executive control are also evident (Chmielewski & Beste, 2015). An important executive control function is the ability to inhibit responses (Simson *et al.* 1977; Pfefferbaum *et al.* 1985; Kok, 1986; Jodo & Kayama, 1992; Falkenstein *et al.* 1999; Bokura *et al.* 2001; Menon *et al.* 2001; Bari & Robbins, 2013). In ASD, findings on response inhibition (RI) are rather ambiguous (for review see Chmielewski & Beste, 2015) with some findings accounting for deficits (Ozonoff *et al.* 1994; Raymaekers *et al.* 2004; Bishop & Norbury, 2005; Kana *et al.* 2007), while others account for non-

compromised RI functions (Hughes, 1996; Geurts *et al.* 2004; Schmitz *et al.* 2006; Duerden *et al.* 2013; Huster *et al.* 2013). In children with ASD a lack of behavioural deficits in RI is reflected by unchanged connectivity patterns between right inferior frontal and medial frontal areas (Lee *et al.* 2009). Other studies accounted for a decreased number of recruited brain regions during RI in ASD (Duerden *et al.* 2013), even though behavioural data do not show differences. This suggests that it is important for measures of neuronal activity to deepen the understanding of RI processes in ASD. Apart from functional imaging this can be achieved using neurophysiological (EEG) measures with the advantage that subprocesses during RI can be examined. Using event-related potentials (ERPs), two RI subprocesses can be distinguished: One process refers to conflict monitoring and/or pre-motor inhibition processes (NoGo-N2) (Falkenstein *et al.* 1999; Nieuwenhuis *et al.* 2003, 2004; Beste *et al.* 2010; Chmielewski *et al.* 2014, 2015c). The other process, reflected by NoGo-P3, is the motor inhibition process *per se* and/or evaluation processes of the successful outcome of an inhibition (Bruin *et al.* 2001; Friedman *et al.* 2001; Roche *et al.* 2004; Schmajuk *et al.* 2006; Beste &

* Address for correspondence: Dr C. Beste, Cognitive Neurophysiology, Department of Child and Adolescent Psychiatry, Faculty of Medicine of the TU Dresden, Schubertstrasse 42, D-01309 Dresden, Germany.
(Email: christian.beste@uniklinikum-dresden.de)

Saft, 2015; Huster *et al.* 2013; Quetscher *et al.* 2015; Chmielewski *et al.* 2015b; Mückschel *et al.* 2015; Wessel & Aron, 2015). In the current study we explore how far these subprocesses are specifically modulated in ASD.

So far, in healthy subjects it has been shown that RI processes are modulated by the complexity of sensory integration processes (Chmielewski *et al.* 2015a): when there is a conflict between information from different sensory modalities, RI processes are complicated (Chmielewski *et al.* 2015a). These cross-modal conflicts during RI processes are of particular relevance for ASD, since deficits in multisensory functions, i.e. in the processing of information across the senses, are increasingly recognized in ASD (Baum *et al.* 2015). For example, even though single senses seem not to be explicitly affected in ASD (Baum *et al.* 2015), it has been shown that ASD patients have difficulties detecting asynchronies between auditory and visual stimuli, as well as in audio-visual integration processes in general (Kwakye *et al.* 2011; Stevenson *et al.* 2014). Moreover, it has been shown that ASD patients have difficulties processing complex stimuli (Bertone *et al.* 2005; Minshew & Hobson, 2008) and do not seem to benefit from redundant information in the same modality (Fuxe *et al.* 2015). Thus, the current study has two questions: (i) Which neurophysiological subprocesses are modulated in ASD during RI?; (ii) How far are RI processes in ASD differentially affected by multisensory integration processes? To answer these questions we use a system neurophysiological approach using high-density EEG recordings and source localization.

Recent findings suggest that RI becomes compromised when conflicting audio-visual information has to be integrated (Chmielewski *et al.* 2015a). Considering deficits in multisensory integration processes in ASD (Baum *et al.* 2015), one hypothesis is that ASD patients may show a superior and impaired RI performance compared to controls, depending on the content of multi-modal stimuli: In case of conflicting multi-modal stimuli, ASD patients might benefit from impairments in multisensory integration processes, since less integrated conflicting information should entail a less compromised RI process. By contrast, in situations where multisensory information usually enhances RI, performance should be decreased in ASD patients compared to controls. However, if multisensory information is properly processed in ASD, then ASD patients may show increased RI deficits under conflicting multisensory information, as this may add up with possible RI deficits reported in ASD.

Regarding neurophysiological processes, we hypothesize the following: Given that multisensory integration processes may be deficient in ASD, it is conceivable that neurophysiological correlates reflecting perceptual or attentional gating and integration processes (i.e. P1 and N1; Herrmann & Knight, 2001)

are affected. Moreover, as mechanisms of resource allocation, reflected by the P2 ERP (Geisler & Murphy, 2000; Campbell & Sharma, 2013; Sugimoto & Katayama, 2013), are modulated by multisensory information during RI (Chmielewski *et al.* 2015a), it is possible that changes (i.e. a decreased P2) in ASD already occur at the early response selection level. The manipulation of multisensory information should, however, specifically affect conflict monitoring processes and thus the NoGo-N2 amplitude in ASD. That is, depending on the extent of conflict, the strength of the N2 amplitude should systematically vary between ASD patients and controls in conditions with concurrent auditory information. Moreover, the P3 amplitude, i.e. the evaluation of, or the RI process *per se* (Bruin *et al.* 2001; Friedman *et al.* 2001; Roche *et al.* 2004; Schmajuk *et al.* 2006; Beste & Saft, 2015; Huster *et al.* 2013; Quetscher *et al.* 2015; Chmielewski *et al.* 2015b; Mückschel *et al.* 2015; Wessel & Aron, 2015), should be generally decreased in ASD, given that RI deficits are evident in ASD.

Materials and method

Participants

A group consisting of adolescents with ASD ($n = 20$, 17 males; mean age 14.8, $S.D. = 1.2$, age range 13.5–16 years) and a control group ($n = 20$, 16 males; mean age 15.0, $S.D. = 1.1$, age range 13.5–16 years) was recruited. All ASD patients and controls were right-handed. Both groups did not differ in IQ scores (ASD: 111.31 ± 3.91 ; controls: 114.06 ± 3.55) ($p > 0.5$), as obtained using the German version of the Wechsler Intelligence Scale for Children (WISC-IV; Petermann & Petermann, 2011). Potential co-morbidities in the ASD group were assessed with the parent questionnaire Child Behavior Checklist (CBCL/4–18; Aschenbach, 1991) and the MINI-KID (Sheehan *et al.* 2010). Two ASD patients were diagnosed with co-morbid attention deficit disorder (ADD) and another two with co-morbid ADD. These co-morbidities were either treated with atomoxetine or with Medikinet. All controls reported no history of neurological or psychiatric disorders and were within normal limits in the MINI-KID and were free of medication. Controls and ASD patients had normal or corrected vision and hearing. Participants in the ASD group fulfilled the diagnostic criteria of childhood autism (F84.0, $n = 8$), Atypical autism (F84.1, $n = 3$), or Asperger syndrome (F84.5, $n = 9$) according to ICD-10-GM (Dilling *et al.* 2015). To ensure participants in the ASD group fulfilled the diagnostic criteria of ASD the Autism Diagnostic Observation Schedule (ADOS, module 3; Lord & Risi, 1998), an observational instrument based on a series of structured and semi-

structured tasks, involving social interaction between the examiner and the subject, was conducted with each participant. ASD participants had an overall score of 14.00 ± 0.99 points (cut-off: 6) in the ADOS. On average 4.26 ± 0.49 points were scored in the communication domain, 7.47 ± 0.49 in the social reciprocity domain, 1.47 ± 0.17 in play skills and imaginative use of objects and 0.89 ± 0.26 in restricted and repetitive behaviours. Moreover on ADOS calibrated severity score (Gotham *et al.* 2007, 2009; Shumway *et al.* 2012) participants scored on average 7.68 ± 1.70 , thus confirming the ASD diagnosis. To complement this, the Autism Diagnostic Interview – Revised (ADI-R; Lord *et al.* 1994) was conducted with the parents, or caregivers. In this test, an average of 14.47 ± 1.38 points were scored in the reciprocal social interaction domain (cut-off: 10), 11.11 ± 0.99 in the communication and language domain (cut-off: 8) and 3.16 ± 0.48 points in the restricted and repetitive behaviours domain (cut-off: 3). Written informed consent was obtained from all participants (and parents).

Task

We used a task introduced by Chmielewski *et al.* (2015a), i.e. a Go/NoGo task, with 70% (672) Go trials, requiring a right-hand response and 30% (288) NoGo trials, requiring withholding of the response. This distribution was chosen to induce a tendency of prepotent response tendencies (Beste *et al.* 2009, 2011). As stimuli, the German words for ‘stop’ (i.e. STOPP) and ‘press’ (DRÜCK) were used. Go trials were always presented visually without any second stimulus occurring, to intensify the effects of concurrent stimuli on NoGo trials (Chmielewski *et al.* 2015a). Out of all NoGo trials, 33% were facilitated (NoGo_{compatible}), and 33% were aggravated (NoGo_{incompatible}) by means of simultaneously presenting either the compatible (same word) or incompatible/conflicting (opposite word) auditory version of the NoGo stimuli verbalized by a female voice. The remaining 33% of NoGo trials were not accompanied by concurrent auditory information (NoGo_{without}). Auditory stimuli were created using ‘Google translate’ to ensure emotional neutrality. To ensure that no effects of exposure time would occur, the presentation onset and offset of visual and auditory stimuli were equated to 400 ms. Participants were explicitly instructed to only respond to visual stimuli, while ignoring auditory stimuli. Trials were separated by inter-trial intervals jittered between 1700 and 2100 ms. Go trials were coded as misses, when no response was obtained within 1000 ms, while NoGo trials were coded as false alarms (FAs), when a response was given in the same time window. To familiarize subjects with the task, a standardized instruction was given and

an exercise with 60 trials was conducted, before the experiment was started.

EEG recording and analysis

EEG data was recorded from 60 Ag/AgCl electrodes mounted in elastic caps (EasyCap Inc., USA) arranged in equidistant positions using BrainAmp amplifiers (Brain Products Inc., USA). The ground electrode was placed at coordinates $\theta = 58$, $\phi = 78$ and the reference electrode at $\theta = 90$, $\phi = 90$, respectively. A sampling rate of 500 Hz was employed and electrode impedances were kept $< 5 \text{ k}\Omega$. Processing of the data was performed using the BrainVision Analyzer 2 software package (Brain Products). Offline, data was downsampled to 256 Hz. A band-pass filter from 0.5 to 20 Hz (with a slope of 48 dB/oct each) and a notch filter at 50 Hz were applied. After a manual inspection of the data, an automatic independent component analysis (ICA; infomax algorithm; Makeig *et al.* 1996) was run to remove recurring artifacts. Only ICA components revealing pulse artifacts, blinks and vertical or horizontal eye movements were discarded. Afterwards, the EEG data was segmented for Go trials, NoGo trials without concurrent information (NoGo_{without}), compatible NoGo trials (NoGo_{compatible}), and incompatible NoGo (NoGo_{incompatible}) trials. Go trials were only included when the correct response was given in a time window until 1000 ms after target onset. Likewise, NoGo trials were only segmented when no response was given in the same time window. The segments were locked to the onset of the target stimulus (Go or NoGo stimulus). After epoching the data, an automated artifact rejection procedure was run for all segments. A difference exceeding $200 \mu\text{V}$ in a 100 ms interval and an activity $< 0.5 \mu\text{V}$ in a 200 ms period, were used as rejection criteria. In order to eliminate reference potential from the data, a current source density (CSD) transformation (Nunez & Pilgreen, 1991) re-referencing the data was then applied. The resulting CSD values are given in $\mu\text{V}/\text{m}^2$. An additional advantage of the CSD transformation is that it serves as a spatial filter (Nunez & Pilgreen, 1991), which makes it possible to identify electrodes that best reflect activity related to cognitive processes (Cohen, 2014; Mückschel *et al.* 2014). Afterwards, a baseline correction was applied in the time interval from -500 to -300 ms prior to target onset. For each condition, individual averages were calculated for every participant. The ERP components at the early processing stage were quantified by detecting peaks at electrodes P7 and P8 (P1: 70–150 ms; N1: 150–220 ms). The ERP components at the response selection stage were quantified at electrode Cz (P2: 160–240 ms; N2: 210–350; P3: 340–540 ms). The electrode sites were identified with the help of the scalp

topographies. This choice of electrodes was validated using statistical methods (cf. Mückschel et al. 2014). Peak quantification was conducted semi-automatically. For all ERP components, peak-to-baseline amplitudes were computed.

Standardized low-resolution brain electromagnetic tomography (sLORETA) analyses

Source localization was conducted using sLORETA (Pascual-Marqui, 2002). sLORETA gives a single linear solution to the inverse problem, based on extra-cranial measurements without a localization bias (Sekihara et al. 2005). sLORETA reveals high convergence with fMRI data and it has been mathematically proven that sLORETA provides reliable results without localization bias (Sekihara et al. 2005). There is also evidence of EEG/fMRI and EEG/TMS studies underlining the validity of the sources estimated using sLORETA (e.g. Sekihara et al. 2005; Dippel & Beste, 2015). For sLORETA, the intracerebral volume is partitioned into 6239 voxels at 5 mm spatial resolution. The standardized current density at each voxel is calculated in a realistic head model (Fuchs et al. 2002) using the MNI152 template (Mazziotta et al. 2001). In this study, the voxel-based sLORETA images were compared across conditions using the sLORETA built-in voxel-wise randomization tests with 3000 permutations, based on statistical non-parametric mapping. Voxels with significant differences ($p < 0.01$, corrected for multiple comparisons) between contrasted conditions were located in the MNI brain (www.unizh.ch/keyinst/NewLORETA/sLORETA/sLORETA.htm).

Statistics

For all ANOVAs (i.e. for the behavioural and neurophysiological data), Greenhouse–Geisser correction was applied and the conducted *post-hoc* tests were Bonferroni-corrected, whenever necessary. All variables included in the analyses were normally distributed, as indicated by Kolmogorov–Smirnov tests (all $z < 0.9$, $p > 0.3$). The mean and standard error of the mean (S.E.M.) are given. Statistic analyses were conducted using PASW 20.

Ethical standards

The authors assert that all procedures contributing to this work comply with the ethical standards of the relevant national and institutional committees on human experimentation and with the Helsinki Declaration of 1975, as revised in 2008. The study was approved by the ethics committee of TU Dresden.

Results

Behavioural data

Concerning Go trial reaction times (RTs) and Go trials misses no significant differences could be found between ASD patients (RTs: 474 ± 22 ms; misses: $3.13 \pm 0.85\%$) and the control group (RTs: 511 ± 20 ms; misses: $2.93 \pm 0.149\%$) (all $t < 1.2$, $p > 0.4$). However, the rate of FAs is the most important behavioural parameter in Go/NoGo tasks. For FAs, repeated-measures ANOVA revealed a main effect of ‘condition’ ($F_{2,76} = 50.46$, $p < 0.001$, $\eta^2 = 0.570$), showing FA rates increasing from the NoGo_{compatible} (14.0 ± 2.0) to the NoGo_{without} (20.1 ± 2.2) to the NoGo_{incompatible} (29.1 ± 2.7) conditions. Moreover, a main effect of ‘group’ was detected ($F_{1,38} = 97.15$, $p < 0.001$, $\eta^2 = 0.719$), showing the ASD group (27.8 ± 3.0) committed more FAs than controls (14.4 ± 3.0). Most importantly, a ‘condition $\times$ group’ interaction was observed ($F_{2,76} = 3.95$, $p = 0.023$, $\eta^2 = 0.570$), which is shown in Fig. 1. *Post-hoc* paired t tests revealed significant differences between the two groups in the NoGo_{without} NoGo_{compatible} and NoGo_{incompatible} conditions (all $t \geq 2.39$, $p \leq 0.022$). We calculated the difference between groups in the FA rate for each condition separately. *Post-hoc* paired t tests with these group differences (NoGo_{without}: 10.4 ± 4.4 ; NoGo_{compatible}: 11.5 ± 4.4 ; NoGo_{incompatible}: 18.2 ± 5.2) revealed that in the NoGo_{incompatible} condition, group differences were larger than in the NoGo_{without} and the NoGo_{compatible} conditions (all $t \geq 2.44$, $p \leq 0.025$). The latter conditions did not differ from each other ($t_{19} = -0.40$, $p = 0.695$).

Regarding the rate of FAs, there was no difference between the first and the second half of trials in each condition in ASD patients and controls (all $p > 0.4$), suggesting that the length of the experiment did not bias the results obtained for patients relative to the controls.

Neurophysiological data

For all ERP components, there were no latency effects (all $F < 1.24$, $p > 0.189$).

Early processing stage

The neurophysiological data for P1 and N1 is shown in Fig. 2. (This data, using no CSD transformation, is shown in Supplementary Fig. S1.)

Concerning the P1 amplitude, a main effect of ‘group’ was found ($F_{1,38} = 186.31$, $p < 0.001$, $\eta^2 = 0.831$), showing ASD patients ($60.5 \pm 5.7 \mu\text{V}/\text{m}^2$) exhibiting larger (i.e. more positive) P1 amplitudes than controls ($50.4 \pm 5.7 \mu\text{V}/\text{m}^2$). sLORETA analysis revealed that this group difference was related to a decreased activation in the cuneus [Brodmann Area (BA) 18] in ASD (Fig. 2). Moreover, a main effect of ‘electrode’ was

Fig. 1. False alarm (FA) rates (with corresponding S.E.M.s) for both groups in the NoGo_{without}, NoGo_{compatible} and NoGo_{incompatible} conditions. ■ Controls; ■ individuals with autism spectrum disorder (ASD). The bars depict the FA group differences in each NoGo condition.

observed ($F_{1,38} = 13.84$, $p = 0.001$, $\eta^2 = 0.267$), showing the P1 amplitude to be smaller (i.e. less positive) in P7 ($47.2 \pm 4.5 \mu\text{V}/\text{m}^2$) than in P8 ($63.7 \pm 7.3 \mu\text{V}/\text{m}^2$). No main effect of 'condition' was observed ($F_{1,38} = 0.56$, $p = 0.640$, $\eta^2 = 0.015$). An 'condition $\times$ group' interaction was observed ($F_{3,114} = 3.24$, $p = 0.025$, $\eta^2 = 0.079$), but *post-hoc* tests did not withstand Bonferroni-corrected *post-hoc* testing (all $t < 1.65$, $p > 0.2$). No further significant effects were found (all $F < 2.03$, $p > 0.265$).

For the N1 amplitudes the repeated-measures ANOVA revealed a main effect of 'condition' ($F_{3,114} = 11.79$, $p < 0.001$, $\eta^2 = 0.237$), showing the N1 amplitudes becoming larger (i.e. more negative) from the Go ($45.8 \pm 4.6 \mu\text{V}/\text{m}^2$) to the NoGo_{without} ($47.5 \pm 4.4 \mu\text{V}/\text{m}^2$) to the NoGo_{compatible} ($55.6 \pm 4.8 \mu\text{V}/\text{m}^2$) to the NoGo_{incompatible} ($56.1 \pm 5.2 \mu\text{V}/\text{m}^2$) conditions. However, *post-hoc* paired *t* tests revealed that all N1 amplitudes differed significantly from each other (all $t \geq 3.72$, $p \leq 0.001$), except for the Go from NoGo_{without} and the NoGo_{compatible} from NoGo_{incompatible} (all $t \leq 1.08$, $p \geq 0.2$) trials. Moreover, a main effect of 'group' was detected ($F_{1,38} = 126.57$, $p < 0.001$, $\eta^2 = 0.769$), showing ASD patients ($43.7 \pm 7.5 \mu\text{V}/\text{m}^2$) with smaller (i.e. less negative) N1 amplitudes than controls ($58.8 \pm 5.1 \mu\text{V}/\text{m}^2$). sLORETA analysis revealed that this group difference was related to a decreased activation in the inferior parietal lobe (IPL; BA 40) in ASD (Fig. 2). No further significant effects were found (all $F < 1.05$, $p > 0.374$). The results of the sLORETA analysis are given in Table 1.

Response selection stage

The P2, N2 and P3 ERPs are shown in Fig. 3. (This data, using no CSD transformation, is shown in Supplementary Fig. S2).

For the P2 amplitude a main effect of 'condition' was observed ($F_{3,114} = 39.85$, $p < 0.001$, $\eta^2 = 0.512$), showing that P2 amplitudes increased from NoGo_{without} ($18.6 \pm 2.7 \mu\text{V}/\text{m}^2$) to Go ($20.2 \pm 2.6 \mu\text{V}/\text{m}^2$) to NoGo_{incompatible} ($35.7 \pm 3.8 \mu\text{V}/\text{m}^2$) to NoGo_{compatible} ($42.3 \pm 4.6 \mu\text{V}/\text{m}^2$) trials. *Post-hoc* paired *t* tests revealed all conditions to differ significantly from each other (all $t \geq 2.27$, $p \leq 0.029$), except for the Go from NoGo_{without} trials ($t_{39} = 1.38$, $p = 0.175$). Moreover, a main effect of 'group' was detected ($F_{1,38} = 85.44$, $p < 0.001$, $\eta^2 = 0.692$), showing ASD patients ($17.8 \pm 4.7 \mu\text{V}/\text{m}^2$) exhibited a smaller (i.e. less positive) P2 than controls ($40.6 \pm 4.5 \mu\text{V}/\text{m}^2$). sLORETA analysis revealed that this group difference was related to a decreased activation in the middle frontal gyrus (MFG; BA 9) in ASD patients (Fig. 3, Table 1). A 'condition $\times$ group' interaction was not found ($F_{3,114} = 1.35$, $p > 0.15$, $\eta^2 = 0.058$).

For the N2 amplitude (see Fig. 3), a main effect of 'condition' ($F_{3,114} = 17.80$, $p < 0.001$, $\eta^2 = 0.319$) was found: N2 amplitudes increased (in negativity) from NoGo_{compatible} ($-22.5 \pm 3.0 \mu\text{V}/\text{m}^2$) to Go ($-33.9 \pm 2.8 \mu\text{V}/\text{m}^2$) to NoGo_{without} ($-38.2 \pm 2.9 \mu\text{V}/\text{m}^2$) to NoGo_{incompatible} ($-42.1 \pm 3.9 \mu\text{V}/\text{m}^2$) trials. All conditions differed significantly from each other (all $t \geq 2.44$, $p \leq 0.019$), except for the NoGo_{without} from NoGo_{incompatible} ($t_{39} = 1.33$, $p = 0.193$) trials. Moreover, a main effect of 'group' was detected ($F_{1,38} = 165.23$, $p < 0.001$, $\eta^2 = 0.813$), showing ASD patients ($-35.9 \pm 3.8 \mu\text{V}/\text{m}^2$) exhibited a larger (i.e. more negative) N2 than controls ($-32.4 \pm 3.8 \mu\text{V}/\text{m}^2$). sLORETA analysis revealed that this group difference was related to an increased activation in the superior frontal gyrus (SFG; BA 8) in ASD (Fig. 3). Most importantly, a 'condition $\times$ group' interaction was observed ($F_{3,114} = 3.47$, $p = 0.019$, $\eta^2 = 0.084$). *Post-hoc* tests

Fig. 2. Event-related potentials (ERPs) on Go and NoGo trials averaged across electrodes P7 and P8. Time-point zero denotes the time-point of Go and NoGo stimulus presentation. The different lines show the NoGo_{without} condition (blue lines), NoGo_{compatible} condition (orange lines) and the NoGo_{incompatible} condition (red lines). Go trials are coloured in green. The scalp topography plots show the distribution of the scalp electrical potential for the P1 (upper row), and N1 (lower row) on Go and NoGo trials. Panel (a) shows the ERPs and scalp topographies for controls. Additionally sLORETA sources for the (main effect of) group differences between controls and autism spectrum disorder (ASD) patients are displayed for P1 and N1. Panel (b) shows the ERPs and scalp topographies for ASD patients.

revealed that this interaction was due to N2 amplitude differences between both groups in NoGo_{compatible} trials ($t_{38} = 2.10$, $p = 0.042$) (ASD: $-28.8 \pm 4.6 \mu\text{V}/\text{m}^2$ *v.* controls: $-16.2 \pm 3.8 \mu\text{V}/\text{m}^2$). sLORETA analysis revealed that this group difference between compatible trials was related to increased activation in the SFG (BA 8) in ASD patients (Fig. 3 and Table 1). The groups did not differ in the other conditions (all $t \leq 0.72$, $p \geq 0.4$).

For the P3 amplitude, a main effect of 'condition' ($F_{3,114} = 29.13$, $p < 0.001$, $\eta^2 = 0.434$) was detected. The P3 amplitude increased from Go ($12.3 \pm 2.5 \mu\text{V}/\text{m}^2$) to NoGo_{without} ($28.9 \pm 3.0 \mu\text{V}/\text{m}^2$) to NoGo_{incompatible} ($32.0 \pm 2.7 \mu\text{V}/\text{m}^2$) to NoGo_{compatible} ($35.8 \pm 4.2 \mu\text{V}/\text{m}^2$)

trials. *Post-hoc* paired *t* tests revealed that all conditions differed significantly from each other (all $t \geq 2.83$, $p \leq 0.007$), except for NoGo_{incompatible} trials, which did not significantly differ from NoGo_{without} and NoGo_{compatible} trials (all $t \leq 1.40$, $p \geq 0.16$). Moreover, a main effect of 'group' was detected ($F_{1,38} = 106.36$, $p < 0.001$, $\eta^2 = 0.737$), showing ASD patients ($22.2 \pm 3.7 \mu\text{V}/\text{m}^2$) exhibited a smaller (i.e. less positive) P3 amplitude, than controls ($32.2 \pm 3.7 \mu\text{V}/\text{m}^2$). sLORETA analysis revealed that this group difference was related to a decreased activation in the MFG (BA 6, BA 10, BA 11) and cingulate gyrus (BA 32) in ASD (Fig. 3, Table 1). A 'condition $\times$ group' interaction was not found ($F_{3,114} = 1.75$, $p = 0.161$, $\eta^2 = 0.044$).

Table 1. Results from sLORETA analysis. The Table provides coordinates of obtained activation differences together with the neuroanatomical structure and Brodmann Areas (BAs) for each of the calculated contrasts

Comparison	Coordinates (x, y, z)	Structure	BA
P1 _{ASD} > N1 _{control}	0, -80, -1	Lingual gyrus	18
	0, -90, -9		
	0, -92, 3	Lingual gyrus	17
	0, -91, -2		
	0, -82, 8	Cuneus	18
	0, -88, 12		
N1 _{control} > N1 _{ASD}	0, -40, 31	Cingulate gyrus	31
	-52, -51, 48	Inferior parietal lobule	40
	-52, -52, 41		
	-39, -44, 59		
P3 _{control} > P3 _{ASD}	-50, -55, 38		
	-50, -47, 42		
	-9, -1, 71	Superior frontal gyrus	6
	-9, 1, 67	Medial frontal gyrus	6
	-9, 0, 63		
	-9, -5, 56		
	-9, -2, 57		
	-9, 64, 13	Medial frontal gyrus	10
	-9, 56, 4		
	-9, 50, -4		
N2 _{ASD} > N2 _{control}	-9, 53, -8		11
	-9, 55, -14		
	-9, 20, 53	Superior frontal gyrus	8
	-9, 15, 53		8
	-9, 14, 65		6
Nogo _{compatible} N2 _{ASD} > N2 _{control}	-9, 16, 60		6
	-9, 14, 54		6
	-8, 39, 44	Medial frontal gyrus	8
	-8, 27, 43		8
	-8, 30, 47		8
	-8, 34, 45		8
	-8, 25, 42	Medial frontal gyrus	6
	-8, 30, 42		6
	-8, 22, 56	Superior frontal gyrus	8

Discussion

In this study we examined the role of conflict in multisensory integration processes during RI in adolescent ASD. The results show that RI performance was

compromised in adolescent ASD in all three RI conditions tested (i.e. without concurrent information, with conflicting information and with redundant information), compared to healthy controls. This shows that there is a general RI deficit in adolescent ASD. Interestingly, there was also a 'group × condition' interaction effect, showing that multisensory information differentially affected RI processes in ASD and controls. Redundant sensory input did not improve RI performance in ASD more than in controls. By contrast, group differences were evident when the multisensory information was conflicting; i.e. RI processes were more dysfunctional in ASD patients than in controls, when conflicting multisensory information was presented during trials requiring the inhibition of responses. This suggests that auditory information is taken into account along with visual information. Importantly, what seems to be central is the content of the auditory information being processed along with the visual information: conflicting multisensory content leads to differences between ASD patients and controls, while non-conflict multisensory content does not lead to differences between ASD patients and controls when RI function as an instance of executive control functions is concerned. With respect to the field of multisensory processing in ASD (cf. Baum *et al.* 2015) this suggests that it is important to consider the content of multisensory information, when cognitive control processes in adolescent ASD are concerned, especially since potential deficits in conflict monitoring processes in ASD have been shown to be related to the involvement of other factors (Chmielewski & Beste, 2015).

The neurophysiological data show which mechanisms in the cognitive processing cascade are modulated between controls and ASD patients, as indexed by increased P1 and decreased N1 amplitudes in ASD deficits already occurring at the perceptual gating and attentional selection level (Herrmann & Knight, 2001). These group differences were due to activation differences in the cuneus (BA 18) and IPL (BA 40), which have frequently been shown to be associated with perceptual gating and attentional selection processes (e.g. Salazar *et al.* 2004; Schintu *et al.* 2014). It seems that perceptual gating processes (i.e. P1 ERP) are paradoxically enhanced, while attentional selection processes (i.e. N1 ERP) are decreased, which is in line with the assumption that multisensory integration processes might be altered in ASD (Baum *et al.* 2015). However, since P1 and N1 are not differentially modulated across NoGo conditions, these modulations cannot explain the observed behavioural effects. Apart from perceptual gating and attentional selection processes, the P2, indexing resource allocation processes (e.g. Geisler & Murphy, 2000; Campbell & Sharma,

Fig. 3. Event-related potentials (ERPs) on Go and NoGo trials at the Cz electrode. Time-point zero denotes the time-point of the Go and NoGo stimulus presentation. The different lines show the NoGo_{without} condition (blue lines), NoGo_{compatible} condition (orange lines) and the NoGo_{incompatible} condition (red lines). Go trials are coloured in green. The scalp topography plots show the distribution of the scalp electrical potential for P2 (upper row), N2 (middle row) and P3 (lower row) on Go and NoGo trials. Panel (a) shows the ERPs and scalp topographies for controls. Additionally sLORETA sources for the P2 main effect of group are displayed in the proximity of the corresponding scalp topographies. Panel (b) shows the ERPs and scalp topographies for autism spectrum disorder (ASD) patients. For N2, the sLORETA source of the group differences in the NoGo_{compatible} condition is displayed.

2013; Sugimoto & Katayama, 2013), was smaller in each of the experimental conditions in ASD patients compared to controls. This suggests that ASD patients are not able to allocate similar large processing resources to the task at hand, which might contribute to the observed behavioural RI deficits. The source localization analyses suggest that modulations seen for P2 are due to functional neuro-anatomical structures in MFG (BA 9). These have frequently been shown to be involved in mechanisms of resource allocation (Peelle *et al.* 2010). However, since resource allocation mechanisms were not differentially modulated between groups across experimental conditions, this suggests that this process only contributes to RI deficits in ASD non-specifically.

Regarding subsequent neurophysiological correlates of subprocesses during RI, the data show that NoGo-N2 and NoGo-P3 were smaller in ASD patients than in controls, which parallels the finding of a generally increased FA rate in the behavioural data. However, only NoGo-N2, but not NoGo-P3 revealed differential effects across conditions. This suggests that conflict and/or pre-motor inhibition processes (reflected by NoGo-N2) (Falkenstein *et al.* 1999; Nieuwenhuis *et al.* 2003, 2004; Beste *et al.* 2010; Chmielewski *et al.* 2014, 2015c), but not subprocesses related to motor inhibition process *per se* and/or evaluation processes of the successful outcome of an inhibition (Schmajuk *et al.* 2006; Beste & Saft, 2015; Huster *et al.* 2013; Quetscher *et al.* 2015; Mückschel *et al.* 2015) play a role in the modulation of RI by multisensory information in ASD. Such a dissociation of affected RI subprocesses in ASD has until now not been shown, which is due to the methods applied in studying RI in ASD (Chmielewski & Beste, 2015). In the NoGo condition without concurrent information no differences in N2 amplitude could be observed. Regarding the condition with conflicting concurrent information, the NoGo-N2 amplitudes were not different between ASD patients and controls. However, NoGo-N2 was larger in ASD patients in the compatible condition. As NoGo-N2 is increased on compatible trials, but not affected in conditions without concurrent information, this suggests that the processing of multisensory information comes with increased demands in patients in ASD. As this increase in N2 amplitude is not evident in the conflicting condition, this result pattern suggests that conflict monitoring processes cannot sufficiently be triggered in ASD patients when conflicting multisensory information has to be controlled. This may in turn lead to the increased FA rates in the conflicting NoGo condition. The source localization analysis suggests that networks in the SFG (BA 8) are differentially modulated between controls and ASD patients in the time range of NoGo-N2. These areas have frequently been shown

to be modulated by RI processes (Chmielewski *et al.* 2015a). Moreover, they are known to show alterations in ASD (for review see Chmielewski & Beste, 2015), suggesting that the functional changes observed are in line with structural changes found in ASD patients. It may, therefore, be speculated that the observed functional changes are due to structural changes in this brain region. The precise relation, however, remains to be tested in future studies. It can also not be ruled out that the generally deficient resource allocation processes (cf. P2 effects), just prior to the conflict monitoring or pre-motor inhibition subprocess, play a role. That is, due to the decreased allocation of processing resources conflict monitoring processes may not unfold to an extent necessary to resolve the conflict imposed by multisensory information.

Regarding clinical relevance it has been suggested that the assessment of cognitive control functions may serve as useful cognitive biomarkers for pharmacological and behavioural treatment studies in ASD (Chmielewski & Beste, 2015). The experimental paradigm used in this study yields strong effects between ASD patients and controls. Therefore, processes examined in this study may be considered for inclusion in clinical studies and in the assessment of cognitive control abilities in ASD patients in routine clinical care.

A limitation of this study is that sample size of the ASD group is relatively low and that the ASD group included childhood autism, atypical autism and Asperger syndrome. It is therefore unclear whether there are differences in the examined cognitive processes between different forms of ASD. This may be a subject for future studies. However, the effect sizes were strong, suggesting that the effects obtained are reliable. Future studies may also extend the scope to ASD in adults.

In summary, the study shows that RI performance is compromised in adolescent ASD and that multisensory processes modulate RI. The neurophysiological processes related to the changes affect the entire processing cascade from perceptual gating to RI subprocesses. However, conflict monitoring and pre-motor RI processes were specifically modulated between ASD patients and controls depending on multisensory information. The data suggest that conflict monitoring processes are overstrained in adolescent ASD upon conflicting multisensory information, which leads to further declines in RI processes. Multisensory integration processes have very specific effects on cognitive control processes in ASD and constitute an important factor in future research to disentangle cognitive control deficits in ASD.

Supplementary material

For supplementary material accompanying this paper visit <http://dx.doi.org/10.1017/S0033291716001008>.

Acknowledgements

We thank all participants for their participation. This work was supported by a grant from the Deutsche Forschungsgemeinschaft (DFG) BE4045/10-2.

Declaration of Interest

V.R. has received payment for consulting and writing activities from Lilly, Novartis, and Shire Pharmaceuticals, lecture honoraria from Lilly, Novartis, Shire Pharmaceuticals, and Medice Pharma, and support for research from Shire and Novartis. He has carried out (and is currently carrying out) clinical trials in cooperation with the Novartis, Shire, and Otsuka companies. C.B. has received payment for consulting from GlaxoSmithKline, and Teva. W.X.C., N.W. and M.M. declare no competing or potential conflicts of interest.

References

- APA (2013). *Diagnostic and Statistical Manual of Mental Disorders*, 5th edn. American Psychiatric Publishing: Arlington, VA.
- Aschenbach TM (1991). *Integrative Guide for the 1991CBCL/4-18, YSR, and TRF Profiles*. Department of Psychiatry: University of Vermont.
- Bari A, Robbins TW (2013). Inhibition and impulsivity: behavioral and neural basis of response control. *Progress in Neurobiology* **108**, 44–79.
- Baum SH, Stevenson RA, Wallace MT (2015). Behavioral, perceptual, and neural alterations in sensory and multisensory function in autism spectrum disorder. *Progress in Neurobiology*, 140–160.
- Bertone A, Motttron L, Jelenic P, Faubert J (2005). Enhanced and diminished visuo-spatial information processing in autism depends on stimulus complexity. *Brain* **128**, 2430–2441.
- Beste C, Dziobek I, Hielscher H, Willemssen R, Falkenstein M (2009). Effects of stimulus-response compatibility on inhibitory processes in Parkinson's disease. *European Journal of Neuroscience* **29**, 855–860.
- Beste C, Ness V, Falkenstein M, Saft C (2011). On the role of fronto-striatal neural synchronization processes for response inhibition—evidence from ERP phase-synchronization analyses in pre-manifest Huntington's disease gene mutation carriers. *Neuropsychologia* **49**, 3484–3493.
- Beste C, Saft C (2015). Action selection in a possible model of striatal medium spiny neuron dysfunction: behavioral and EEG data in a patient with benign hereditary chorea. *Brain Structure and Function* **220**, 221–228.
- Beste C, Willemssen R, Saft C, Falkenstein M (2010). Response inhibition subprocesses and dopaminergic pathways: basal ganglia disease effects. *Neuropsychologia* **48**, 366–373.
- Bishop DVM, Norbury CF (2005). Executive functions in children with communication impairments, in relation to autistic symptomatology 2: response inhibition. *Autism* **9**, 29–43.
- Bokura H, Yamaguchi S, Kobayashi S (2001). Electrophysiological correlates for response inhibition in a Go/NoGo task. *Clinical Neurophysiology* **112**, 2224–2232.
- Bruin KJ, Wijers AA, van Staveren AS (2001). Response priming in a go/nogo task: do we have to explain the go/nogo N2 effect in terms of response activation instead of inhibition? *Clinical Neurophysiology* **112**, 1660–1671.
- Campbell J, Sharma A (2013). Compensatory changes in cortical resource allocation in adults with hearing loss. *Frontiers in Systems Neuroscience* **7**, 71.
- Chmielewski WX, Beste C (2015). Action control processes in autism spectrum disorder – insights from a neurobiological and neuroanatomical perspective. *Progress in Neurobiology* **124**, 49–83.
- Chmielewski WX, Mückschel M, Dippel G, Beste C (2015a). Concurrent information affects response inhibition processes via the modulation of theta oscillations in cognitive control networks. *Brain Structure and Function*. doi:10.1007/s00429-015-1137-1.
- Chmielewski WX, Mückschel M, Roessner V, Beste C (2014). Expectancy effects during response selection modulate attentional selection and inhibitory control networks. *Behavioural Brain Research* **274C**, 53–61.
- Chmielewski WX, Mückschel M, Stock A-K, Beste C (2015b). The impact of mental workload on inhibitory control subprocesses. *NeuroImage* **112**, 96–104.
- Chmielewski WX, Roessner V, Beste C (2015c). Predictability and context determine differences in conflict monitoring between adolescence and adulthood. *Behavioural Brain Research* **292**, 10–18.
- Cohen MX (2014). *Analyzing Neural Time Series Data. Theory and Practice*. MIT Press.
- Dilling H, Mombour W, Schmidt MH, Schulte-Markwort E, Remschmidt H (eds) (2015). *Internationale Klassifikation psychischer Störungen ICD-10 Kapitel V (F). Klinisch-diagnostische Leitlinien*. Huber: Bern.
- Dippel G, Beste C (2015). A causal role of the right inferior frontal cortex in implementing strategies for multi-component behaviour. *Nature Communications* **6**, 6587.
- Duerden EG, Taylor MJ, Soorya LV, Wang T, Fan J, Anagnostou E (2013). Neural correlates of inhibition of socially relevant stimuli in adults with autism spectrum disorder. *Brain Research* **1533**, 80–90.
- Falkenstein M, Hoormann J, Hohnsbein J (1999). ERP components in Go/Nogo tasks and their relation to inhibition. *Acta Psychologica* **101**, 267–291.
- Foxe JJ, Molholm S, Del Bene VA, Frey H-P, Russo NN, Blanco D, Saint-Amour D, Ross LA (2015). Severe multisensory speech integration deficits in high-functioning school-aged children with Autism Spectrum Disorder (ASD) and their resolution during early adolescence. *Cerebral Cortex (New York)* **25**, 298–312.
- Friedman D, Cycowicz YM, Gaeta H (2001). The novelty P3: an event-related brain potential (ERP) sign of the brain's evaluation of novelty. *Neuroscience and Biobehavioral Reviews* **25**, 355–373.
- Fuchs M, Kastner J, Wagner M, Hawes S, Ebersole JS (2002). A standardized boundary element method volume conductor model. *Clinical Neurophysiology* **113**, 702–712.

- Geisler MW, Murphy C (2000). Event-related brain potentials to attended and ignored olfactory and trigeminal stimuli. *International Journal of Psychophysiology* **37**, 309–315.
- Geurts HM, Verté S, Oosterlaan J, Roeyers H, Sergeant JA (2004). How specific are executive functioning deficits in attention deficit hyperactivity disorder and autism? *Journal of Child Psychology and Psychiatry* **45**, 836–854.
- Gotham K, Pickles A, Lord C (2009). Standardizing ADOS scores for a measure of severity in autism spectrum disorders. *Journal of Autism and Developmental Disorders* **39**, 693–705.
- Gotham K, Risi S, Pickles A, Lord C (2007). The autism diagnostic observation schedule: revised algorithms for improved diagnostic validity. *Journal of Autism and Developmental Disorders* **37**, 613–627.
- Herrmann CS, Knight RT (2001). Mechanisms of human attention: event-related potentials and oscillations. *Neuroscience and Biobehavioral Reviews* **25**, 465–476.
- Hughes C (1996). Control of action and thought: normal development and dysfunction in autism: a research note. *Journal of Child Psychology and Psychiatry* **37**, 229–236.
- Huster RJ, Enriquez-Geppert S, Lavalée CF, Falkenstein M, Herrmann CS (2013). Electroencephalography of response inhibition tasks: functional networks and cognitive contributions. *International Journal of Psychophysiology* **87**, 217–233.
- Jodo E, Kayama Y (1992). Relation of a negative ERP component to response inhibition in a Go/No-go task. *Electroencephalography and Clinical Neurophysiology* **82**, 477–482.
- Kana RK, Keller TA, Minshew NJ, Just MA (2007). Inhibitory control in high-functioning autism: decreased activation and underconnectivity in inhibition networks. *Biological Psychiatry* **62**, 198–206.
- Kok A (1986). Effects of degradation of visual stimuli on components of the event-related potential (ERP) in go/nogo reaction tasks. *Biological Psychology* **23**, 21–38.
- Kwakye LD, Foss-Feig JH, Cascio CJ, Stone WL, Wallace MT (2011). Altered auditory and multisensory temporal processing in autism spectrum disorders. *Frontiers in Integrative Neuroscience* **4**, 129.
- Lee PS, Yerys BE, Rosa AD, Foss-Feig J, Barnes KA, James JD, VanMeter J, Vaidya CJ, Gaillard WD, Kenworthy LE (2009). Functional connectivity of the inferior frontal cortex changes with age in children with autism spectrum disorders: a fMRI study of response inhibition. *Cerebral Cortex* **19**, 1787–1794.
- Lord C, Risi S (1998). Frameworks and methods in diagnosing autism spectrum disorders. *Mental Retardation and Developmental Disabilities Research Reviews* **4**, 90–96.
- Lord C, Rutter M, Couteur AL (1994). Autism diagnostic interview-revised: a revised version of a diagnostic interview for caregivers of individuals with possible pervasive developmental disorders. *Journal of Autism and Developmental Disorders* **24**, 659–685.
- Makeig S, Bell AJ, Jung T-P, Sejnowski TJ (1996). Independent component analysis of electroencephalographic data. In *Advances in Neural Information Processing Systems*, Vol. 8 (ed. D.S. Touretzky), pp. 141–151. MIT Press: Cambridge, Massachusetts, USA.
- Mazziotta J, Toga A, Evans A, Fox P, Lancaster J, Zilles K, Woods R, Paus T, Simpson G, Pike B, Holmes C, Collins L, Thompson P, MacDonald D, Iacoboni M, Schormann T, Amunts K, Palomero-Gallagher N, Geyer S, Parsons L, Narr K, Kabani N, Le Goualher G, Boomsma D, Cannon T, Kawashima R, Mazoyer B (2001). A probabilistic atlas and reference system for the human brain: International Consortium for Brain Mapping (ICBM). *Philosophical Transactions of the Royal Society of London. Series B* **356**, 1293–1322.
- Menon V, Adelman NE, White CD, Glover GH, Reiss AL (2001). Error-related brain activation during a Go/NoGo response inhibition task. *Human Brain Mapping* **12**, 131–143.
- Minshew NJ, Hobson JA (2008). Sensory sensitivities and performance on sensory perceptual tasks in high-functioning individuals with autism. *Journal of Autism and Developmental Disorders* **38**, 1485–1498.
- Mückschel M, Smitka M, Hermann A, von der Hagen M, Beste C (2015). Deep brain stimulation in the globus pallidus compensates response inhibition deficits: evidence from pantothenate kinase-associated neurodegeneration. *Brain Structure and Function* **221**, 2251–2257.
- Mückschel M, Stock A-K, Beste C (2014). Psychophysiological mechanisms of interindividual differences in goal activation modes during action cascading. *Cerebral Cortex* **24**, 2120–2129.
- Nieuwenhuis S, Yeung N, Cohen JD (2004). Stimulus modality, perceptual overlap, and the go/no-go N2. *Psychophysiology* **41**, 157–160.
- Nieuwenhuis S, Yeung N, van den Wildenberg W, Ridderinkhof KR (2003). Electrophysiological correlates of anterior cingulate function in a go/no-go task: effects of response conflict and trial type frequency. *Cognitive, Affective, and Behavioral Neuroscience* **3**, 17–26.
- Nunez PL, Pilgreen KL (1991). The spline-Laplacian in clinical neurophysiology: a method to improve EEG spatial resolution. *Journal of Clinical Neurophysiology* **8**, 397–413.
- Ozonoff S, Strayer DL, McMahon WM, Filloux F (1994). Executive function abilities in autism and tourette syndrome: an information processing approach. *Journal of Child Psychology and Psychiatry* **35**, 1015–1032.
- Pascual-Marqui RD (2002). Standardized low-resolution brain electromagnetic tomography (sLORETA): technical details. *Methods and Findings in Experimental and Clinical Pharmacology* **24**(Suppl. D), 5–12.
- Peelle JE, Troiani V, Wingfield A, Grossman M (2010). Neural processing during older adults' comprehension of spoken sentences: age differences in resource allocation and connectivity. *Cerebral Cortex (New York, N.Y.: 1991)* **20**, 773–782.
- Petermann F, Petermann U (2011). *Wechsler Intelligence Scale for Children*, 4th edn. Pearson Assessment: Frankfurt/Main.
- Pfefferbaum A, Ford JM, Weller BJ, Kopell BS (1985). ERPs to response production and inhibition. *Electroencephalography and Clinical Neurophysiology* **60**, 423–434.
- Quetscher C, Yildiz A, Dharmadhikari S, Glaubit B, Schmidt-Wilcke T, Dydak U, Beste C (2015). Striatal GABA-MRS predicts response inhibition performance and

- its cortical electrophysiological correlates. *Brain Structure and Function* **220**, 3555–3564.
- Raymaekers R, van der Meere J, Roeyers H** (2004). Event-rate manipulation and its effect on arousal modulation and response inhibition in adults with high functioning autism. *Journal of Clinical and Experimental Neuropsychology* **26**, 74–82.
- Roche RAP, Garavan H, Foxe JJ, O'Mara SM** (2004). Individual differences discriminate event-related potentials but not performance during response inhibition. *Experimental Brain Research* **160**, 60–70.
- Salazar RF, Kayser C, König P** (2004). Effects of training on neuronal activity and interactions in primary and higher visual cortices in the alert cat. *Journal of Neuroscience* **24**, 1627–1636.
- Schintu S, Hadj-Bouziane F, Dal Monte O, Knutson KM, Pardini M, Wassermann EM, Grafman J, Krueger F** (2014). Object and space perception – is it a matter of hemisphere? *Cortex* **57**, 244–253.
- Schmajuk M, Liotti M, Busse L, Woldorff MG** (2006). Electrophysiological activity underlying inhibitory control processes in normal adults. *Neuropsychologia* **44**, 384–395.
- Schmitz N, Rubia K, Daly E, Smith A, Williams S, Murphy DGM** (2006). Neural correlates of executive function in autistic spectrum disorders. *Biological Psychiatry* **59**, 7–16.
- Sekihara K, Sahani M, Nagarajan SS** (2005). Localization bias and spatial resolution of adaptive and non-adaptive spatial filters for MEG source reconstruction. *NeuroImage* **25**, 1056–1067.
- Sheehan DV, Sheehan KH, Shytle RD, Janavs J, Bannon Y, Rogers JE, Milo KM, Stock SL, Wilkinson B** (2010). Reliability and validity of the mini international neuropsychiatric interview for children and adolescents (MINI-KID). *Journal of Clinical Psychiatry* **71**, 313–326.
- Shumway S, Farmer C, Thurm A, Joseph L, Black D, Golden C** (2012). The ADOS calibrated severity score: relationship to phenotypic variables and stability over time. *Autism Research* **5**, 267–276.
- Simson R, Vaughan Jr. HG, Ritter W** (1977). The scalp topography of potentials in auditory and visual discrimination tasks. *Electroencephalography and Clinical Neurophysiology* **42**, 528–535.
- Stevenson RA, Siemann JK, Woynaroski TG, Schneider BC, Eberly HE, Camarata SM, Wallace MT** (2014). Evidence for diminished multisensory integration in autism spectrum disorders. *Journal of Autism and Developmental Disorders* **44**, 3161–3167.
- Sugimoto F, Katayama J** (2013). Somatosensory P2 reflects resource allocation in a game task: assessment with an irrelevant probe technique using electrical probe stimuli to shoulders. *International Journal of Psychophysiology* **87**, 200–204.
- Wessel JR, Aron AR** (2015). It's not too late: the onset of the frontocentral P3 indexes successful response inhibition in the stop-signal paradigm. *Psychophysiology* **52**, 472–480.