**Dieses Dokument ist eine Zweitveröffentlichung (Verlagsversion) /
This is a self-archiving document (published version):**

Gabriel Fuhrmann

Non-smooth saddle-node bifurcations I: existence of an SNA

Erstveröffentlichung in / First published in:

*Ergodic theory and dynamical systems. 2017, 36(4), S. 1130– 1155 [Zugriff am: 15.04.2020].
Cambridge University Press. ISSN 1469-4417.*

DOI: <https://doi.org/10.1017/etds.2014.92>

Diese Version ist verfügbar / This version is available on:

<https://nbn-resolving.org/urn:nbn:de:bsz:14-qucosa2-707074>

„Dieser Beitrag ist mit Zustimmung des Rechteinhabers aufgrund einer (DFGgeförderten) Allianz- bzw. Nationallizenz frei zugänglich.“

This publication is openly accessible with the permission of the copyright owner. The permission is granted within a nationwide license, supported by the German Research Foundation (abbr. in German DFG).
www.nationallizenzen.de/

Non-smooth saddle-node bifurcations I: existence of an SNA

GABRIEL FUHRMANN

*Emmy Noether Group: Low-dimensional and Nonautonomous Dynamics,
TU-Dresden, Dresden, Germany
(e-mail: Gabriel.Fuhrmann@mailbox.tu-dresden.de)*

(Received 6 August 2013 and accepted in revised form 18 August 2014)

Abstract. We study one-parameter families of quasi-periodically forced monotone interval maps and provide sufficient conditions for the existence of a parameter at which the respective system possesses a non-uniformly hyperbolic attractor. This is equivalent to the existence of a sink-source orbit, that is, an orbit with positive Lyapunov exponent both forwards and backwards in time. The attractor itself is a non-continuous invariant graph with negative Lyapunov exponent, often referred to as ‘SNA’. In contrast to former results in this direction, our conditions are $\mathcal{C}^2$ -open in the fibre maps. By applying a general result about saddle-node bifurcations in skew-products, we obtain a conclusion on the occurrence of non-smooth bifurcations in the respective families. Explicit examples show the applicability of the derived statements.

1. Introduction

Bifurcation theory investigates qualitative changes in the long-term behaviour of a dynamical system along a continuous variation of the system. For the simple case of a monotonously increasing interval map, the dynamics are qualitatively understood if the fixed points of the respective system are known. Hence in this case bifurcation theory investigates the bifurcation of fixed points. A well-known example is the saddle-node bifurcation of a one-parameter family of concave functions: if the considered parameter is ‘small’, we have one attracting and one repelling fixed point, which approach each other upon the growth of the parameter until some threshold is reached. Above this threshold, these two fixed points have vanished. At the threshold itself, the two points merge together to one neutral fixed point.

In general, non-autonomous systems ask for objects other than fixed points (which might not even exist) to describe the qualitative dynamics [22]. In the context of quasi-periodically forced monotone maps, a natural choice are *invariant graphs*. Like fixed points of monotone interval maps, these are barriers which can’t be crossed by an orbit.

Furthermore, there is a one-to-one correspondence between the ergodic measures of a quasi-periodically forced monotone map and its invariant graphs (cf. [2, Theorem 1.8.4], [1, Theorem 2.2]).

We consider bifurcations of the invariant graphs of a class of quasi-periodically driven (also: forced) monotone interval maps

$$f : \mathbb{T}^d \times X \rightarrow \mathbb{T}^d \times X, \quad (\theta, x) \mapsto (\theta + \omega, \tilde{f}(\theta, x)), \quad (1.1)$$

where X denotes an interval, ω is irrational, $\tilde{f}(\theta, \cdot)$ is strictly increasing, and $\tilde{f}$ is $\mathcal{C}^2$. The most studied bifurcation patterns related to such maps are the pitchfork bifurcation (cf. [7, 12, 20, 21]) and the saddle-node bifurcation (cf. [1, 3, 12, 20]). In this article, we deal with the latter phenomenon: we consider a one-parameter family of driven interval maps of the form (1.1) and assume that for small parameters there are two invariant graphs (an attracting and a repelling one) which approach each other along the growth of the parameter until some threshold is reached. Above this threshold, these two invariant graphs have vanished.

In contrast to the autonomous situation, the forced case allows a dichotomy at the threshold: either there is just one neutral invariant graph, or there are two invariant graphs; an attracting and a repelling one. Further, these graphs are *pinched*, that is, they coincide in a point, and hence on a residual set [24], while they almost surely differ from each other (cf. Figure 1). This is called a *non-smooth saddle-node bifurcation*. The attracting graph is referred to as a *strange non-chaotic attractor* (SNA); the repelling one as a *strange non-chaotic repeller* (SNR).

Quasi-periodic forcing and SNAs play an important role in a large class of models for real life systems: the Harper map is a mathematically well-understood dynamical system related to a certain kind of quasi-periodic Schrödinger equations (see below); there is numerical evidence for the existence of SNAs in the physiologically relevant Izhikevich Neuron Model [17]; [23] shows that strange non-chaotic attractors have to be considered not only to get a complete description of the tides – as the result of the gravitational interaction between the Earth, the Moon, and the Sun – but also even to predict interdecadal atmospheric variations. Further, [5] investigates the succession of ice ages and numerically encounters bifurcation phenomena creating SNAs.

It is thus desirable to understand the underlying principles of the creation of SNAs. First results in this direction were obtained by Millionščikov [18], Vinograd [26] and Herman [11] who considered quasi-periodic $SL(2, \mathbb{R})$ -cocycles. In this context, the phenomenon is also known under the name of *non-uniform hyperbolicity*. In 1984, Grebogi *et al* found numerical evidence for SNAs in so-called *pinched skew-products* [8]; a rigorous proof in this setting is due to Keller [16]. Still, these findings lack some flexibility. By implementing parameter exclusion techniques, this has been overcome in more recent results by Young [27], Bjerklov [3] and Jäger [15].

Inspired by these works, it is the goal of this article to derive general conditions for the existence of SNAs, and thus for the occurrence of non-smooth saddle-node bifurcations for maps of the form (1.1). So far, related results have dealt only with special families of skew-products [3, 12]. While [3] considers the particular case of the Harper map, [12] yields a result for additive forcing by imposing a non-differentiability assumption on the forcing

FIGURE 1. The non-smooth saddle-node bifurcation of the family $(\theta, x) \mapsto (\theta + \omega, \arctan(\alpha x) - \beta(1 + \cos 2\pi\theta))$, where ω is the golden mean and $\alpha = 100$. We see how the invariant graphs approach each other on a measure zero set as we increase the parameter β from the left to the right. The red graph is attracting; the blue one repelling. (a) $\beta = 0.7769$; (b) $\beta = 0.7805$; (c) $\beta = 0.7805931$.

term and therefore excludes the application to smooth examples. The main achievement of this article is the merging of these two technically demanding approaches in order to get a natural and flexible statement on the non-smoothness of saddle-node bifurcations. The following assertion is a direct consequence of our results.

THEOREM. *Let X be an interval, suppose $\omega \in \mathbb{T}^d$ is Diophantine and consider the space of one-parameter families*

$$\mathcal{F}_\omega = \{(f_\beta)_{\beta \in [0,1]} : f_\beta \text{ is of the form (1.1) with rotation number } \omega \text{ and phase space } X; \\ f_\beta \text{ is } C^1 \text{ in } \beta\}$$

equipped with the metric

$$d((f_\beta)_{\beta \in [0,1]}, (g_\beta)_{\beta \in [0,1]}) = \sup_{\beta \in [0,1]} (\|f_\beta - g_\beta\|_2 + \|\partial_\beta f_\beta - \partial_\beta g_\beta\|_0).$$

There exists an open set $\mathcal{U} \subseteq \mathcal{F}_\omega$ such that each $(f_\beta)_{\beta \in [0,1]} \in \mathcal{U}$ undergoes a non-smooth saddle-node bifurcation.

The precise description of the set $\mathcal{U}$ by means of conditions on the maps $\tilde{f}_\beta$ is given in §3. Though a bit technical, these conditions are explicit and easy to check in examples. In order to demonstrate the applicability, we further show that for each Diophantine $\omega \in \mathbb{T}^1$ the family $(f_\beta)_{\beta \in [0,1]}$

$$f_\beta : \mathbb{T}^1 \times \mathbb{R} \rightarrow \mathbb{T}^1 \times \mathbb{R}, \quad (\theta, x) \mapsto (\theta + \omega, \arctan(\alpha x) - \beta(1 + \cos 2\pi\theta))$$

undergoes a non-smooth saddle-node bifurcation if α is big enough, by showing that $(f_\beta)_{\beta \in [0,1]}$ lies in the open set $\mathcal{U} \subseteq \mathcal{F}_\omega$ (cf. Figure 1).

With the existence of SNAs there naturally arise several questions: at the bifurcation point, the set which is bounded by the pinched graphs is invariant. Is it even minimal? For the Harper map there is a positive answer to this problem for particular parameter regions [3]; the case of *pinched* skew-products has been studied in [13]. Further, and closely related: what is the Hausdorff dimension of the pinched graphs [9]? These problems are dealt with in a follow-up article [6]. Finally, we don't address the problem of the speed with which the two initially continuous invariant graphs converge to the SNA and SNR, respectively [4, 10]. Note that this information is crucial when it comes to the

prediction of a bifurcation in real life problems as in [25], where the collapse of a bacteria population exposed to an increasing light pressure is studied.

The present work provides the basis for these further studies and the developed techniques should allow us to answer the above questions and approach further problems of similar type.

2. Preliminaries

2.1. *Basic setting and notation.* Throughout this article, we consider families of $\mathcal{C}^2$ -skew-products[†] $(f_\beta)_{\beta \in [0,1]}$

$$\begin{aligned} f_\beta : \mathbb{T}^d \times X &\rightarrow \mathbb{T}^d \times X, \\ (\theta, x) &\mapsto (\theta + \omega, \tilde{f}_\beta(\theta, x)), \end{aligned} \quad (2.1)$$

where X is an interval (possibly the real line, half-open, $\dots$), $\tilde{f}_\beta(\theta, \cdot)$ is strictly increasing and $\omega \in \mathbb{T}^d$ is *Diophantine* in the following sense.

Definition 2.1. Let $\mathcal{C}, \eta > 0$. We say $\omega \in \mathbb{T}^d$ is *Diophantine* of type $(\mathcal{C}, \eta)$ if $d(k\omega, 0) \geq \mathcal{C}|k|^{-\eta}$ for all $k \in \mathbb{Z} \setminus \{0\}$, where $|\cdot|$ denotes the absolute value.

It is obvious that the above definition includes all those $\omega \in \mathbb{T}^d$ which are Diophantine in the usual sense. We refer to a skew-product of the form (2.1) as a *quasi-periodically forced (qpf) monotone interval map*. $\mathbb{T}^d$ is called the *base* and $\{\theta\} \times X$ a *fibre* of (2.1) for $\theta \in \mathbb{T}^d$.

By writing $f_\beta^{-l}(\theta, x)$ for $l \in \mathbb{N}$, we implicitly assume $(\theta, x) \in f_\beta^l(\mathbb{T}^d, X)$ such that the respective expression is well-defined due to the injectivity of f_β . For $l \in \mathbb{Z}$ and $(\theta_0, x_0) \in \mathbb{T}^d \times X$, we set $(\theta_l, x_l) := f^l(\theta_0, x_0)$. Given β and θ , we call

$$f_{\beta, \theta} : x \mapsto \tilde{f}_\beta(\theta, x)$$

a *fibre map* of the skew-product (2.1). The fibre map of f_β^n at θ is denoted by $f_{\beta, \theta}^n$, where $n \in \mathbb{Z}$. Hence, for $n \in \mathbb{N}$ we have

$$f_{\beta, \theta}^n(x) = \pi_2 \circ f_\beta^n(\theta, x) = f_{\beta, \theta + (n-1)\omega} \circ \dots \circ f_{\beta, \theta}(x),$$

where π_2 is the canonical projection to the second coordinate. Further, $f_{\beta, \theta}^{-1}(x) = (f_{\beta, \theta - \omega})^{-1}(x)$.

We denote the derivative of $\pi_2 \circ f_\beta^n(\theta, x)$ with respect to β by $\partial_\beta f_{\beta, \theta}^n(x)$; the directional derivative of $\pi_2 \circ f_\beta^n(\theta, x)$ with respect to a direction $\vartheta \in \mathbb{T}^d \setminus \{0\}$ is denoted by $\partial_\vartheta f_{\beta, \theta}(x)$. Higher derivatives are denoted in an analogous way. Typically, we will consider ϑ to be a unit vector and write $\vartheta \in \mathbb{S}^{d-1}$. The derivatives of the fibre maps are denoted by $\partial_x f_{\beta, \theta}(x)$.

Given $\beta \in [0, 1]$, we are interested in studying the *invariant graphs* of f_β . These are measurable functions $\phi : \mathbb{T}^d \rightarrow X$ satisfying

$$f_\beta(\phi(\theta)) = \phi(\theta + \omega),$$

[†] As the hypothesis of our main result are of a local form, we need the systems to be $\mathcal{C}^2$ only in a section of the phase space, while on the complement of this section it suffices to assume they are just continuous and continuously differentiable with respect to x .

such that the corresponding graph $\Phi := \{(\theta, \phi(\theta)) \in \mathbb{T}^d \times X : \theta \in \mathbb{T}^d\}$ is an invariant set under f_β in the sense that $f_\beta(\Phi) = \Phi$. Like fixed points in monotone, unforced (that is, autonomous) interval maps, invariant graphs are barriers that can not be crossed by orbits of qpf monotone interval maps.

Similar to the unforced situation, the stability of an invariant graph ϕ is closely related to its associated Lyapunov exponent [14, Proposition 3.3], which is given by

$$\lambda(\phi) := \int_{\mathbb{T}^d} \log |\partial_x f_\theta(\phi(\theta))| d\theta.$$

We say an invariant graph ϕ is an *attractor* if $\lambda(\phi) < 0$; we call it *repeller* if $\lambda(\phi) > 0$; and we call it *neutral* if $\lambda(\phi) = 0$.

As a matter of fact, many statements for fixed points of unforced maps remain true (with slight changes) when going over to invariant graphs of qpf interval maps.

THEOREM 2.2. (Cf. [1, Theorem 2.1]) *Consider a qpf monotone interval map f of the form (2.1). Assume that for each $\theta \in \mathbb{T}^d$ there exist measurable functions $\gamma^- \leq \gamma^+ : \mathbb{T}^d \rightarrow X$ such that for all $\theta \in \mathbb{T}^d$ the fibre maps are strictly concave on $\Gamma(\theta) := [\gamma^-(\theta), \gamma^+(\theta)]$. Further, assume that $h(\theta) := \inf_{x \in \Gamma(\theta)} \log \partial_x f_\theta(x)$ has an integrable minorant.*

Then there exist at most two distinct invariant graphs in $\Gamma := \{(\theta, x) \in \mathbb{T}^d \times X : x \in \Gamma(\theta)\}^\dagger$. Moreover, if there exist two distinct invariant graphs $\phi^- \leq \phi^+$ in Γ , then ϕ^- is a repeller and ϕ^+ is an attractor.

We call two invariant graphs ϕ, ψ *pinched* if $\phi(\theta) = \psi(\theta)$ for some θ . Note that due to the minimality of the *base map* $\theta \mapsto \theta + \omega$, this implies that the two graphs coincide on a residual set in $\mathbb{T}^d$ [24]. It is obvious that in this case at least one of the two graphs is non-continuous. A non-continuous invariant graph ϕ is called a *strange non-chaotic attractor* (SNA) if $\lambda(\phi) < 0$; it is called a *strange non-chaotic repeller* (SNR) if $\lambda(\phi) > 0$.

In 1984, Grebogi *et al* found numerical evidence for the existence of an SNA in the case of *pinched systems* [8]. These are qpf interval maps f with monotone fibre maps which leave the zero line invariant (that is, $f_\theta(0) = 0$) and possess a *pinched point*, that is, there exists a point $\theta_0 \in \mathbb{T}^d$ with $f_{\theta_0}(x) = 0$. Under additional assumptions, the zero-line turns out to be repelling. Thus, by proving the existence of an attracting graph (which necessarily has to share a residual set with the zero-line), Keller gave a rigorous argument for the existence of an SNA for pinched systems [16]. In the injective setting (where there are no pinched points), we cannot argue in such a comparably direct way. Instead, the following concept proves helpful.

Definition 2.3. A *sink-source orbit* is an orbit whose backward and forward vertical Lyapunov exponent is positive, that is,

$$\limsup_{n \rightarrow \infty} \frac{1}{n} \ln |\partial_x f_{\beta, \theta}^n(x)| > 0 \quad \text{and} \quad \limsup_{n \rightarrow \infty} \frac{1}{n} \ln |\partial_x f_{\beta, \theta}^{-n}(x)| > 0.$$

[†] A graph ϕ is said to be contained in Γ if $\phi(\theta) \in \Gamma(\theta)$ for all $\theta \in \mathbb{T}^d$. Note that we identify invariant graphs which coincide almost surely.

THEOREM 2.4. [12, Theorem 2.4] Suppose $f_\beta : \mathbb{T}^d \times X \rightarrow \mathbb{T}^d \times X$ is a continuous qpf monotone interval map and $(\theta, x) \mapsto \partial_x f_\beta(x) > 0$ is continuous. Further, assume the existence of a sink-source-orbit contained entirely in $\mathbb{T}^d \times I$, where $I \subseteq X$ is some closed interval. Then there exist both an SNA and an SNR within $\mathbb{T}^d \times I$.

Remark 2.5. The respective statement in [12] is slightly weaker. However, by restricting the map f_β to the closure of the sink-source-orbit, the proof in [12] applies literally in the above setting.

2.2. Saddle-node bifurcations in quasi-periodically forced systems. Non-autonomous bifurcation theory in the setting of one-parameter families of qpf monotone interval maps studies the bifurcation of invariant graphs along the change of the parameter. An often considered situation is that of a saddle-node bifurcation [1, 12, 19, 20]: there exists a critical parameter β_c such that for $\beta < \beta_c$ there are two continuous invariant graphs, while there is no invariant graph for $\beta > \beta_c$. At $\beta = \beta_c$ there exists (in contrast to the unforced case) a dichotomy: there occurs either a *smooth* or a *non-smooth saddle-node* bifurcation.

THEOREM 2.6. (Cf. [1, Theorem 6.1]) Let $\omega \in \mathbb{T}^d$ and suppose $(f_\beta)_{\beta \in [0,1]}$ is a family of qpf monotone C^2 interval maps. Further, assume that there exist continuous functions $\gamma^-, \gamma^+ : \mathbb{T}^d \rightarrow X$ with $\gamma^- < \gamma^+$ such that the following holds (for all $\beta \in [0, 1]$ and $\theta \in \mathbb{T}^d$ where applicable):

- (i) there exist two distinct continuous f_0 -invariant graphs and no f_1 -invariant graph in Γ ;
- (ii) $f_{\beta,\theta}(\gamma^\pm(\theta)) \leq \gamma^\pm(\theta + \omega)$;
- (iii) the maps $(\beta, \theta, x) \mapsto \partial_x^i f_\beta(\theta, x)$ with $i = 0, 1, 2$ and $(\beta, \theta, x) \mapsto \partial_\beta f_\beta(\theta, x)$ are continuous;
- (iv) $\partial_x f_{\beta,\theta}(x) > 0$ for all $x \in \Gamma(\theta)$;
- (v) $\partial_x^2 f_{\beta,\theta}(x) < 0$ ($x \in \mathring{\Gamma}(\theta)$);
- (vi) $\partial_\beta f_{\beta,\theta}(x) < 0$ ($x \in \Gamma(\theta)$).

Then there exists a unique critical parameter $\beta_c \in (0, 1)$ such that the following hold.

- If $\beta < \beta_c$, then there exist two continuous f_β -invariant graphs $\phi_\beta^- < \phi_\beta^+$ in Γ with $\lambda(\phi_\beta^-) > 0$ and $\lambda(\phi_\beta^+) < 0$.
- If $\beta = \beta_c$, then either there exists exactly one f_β -invariant graph ϕ_β in Γ , or there exist two semi-continuous and pinched f_β -invariant graphs $\phi_\beta^- < \phi_\beta^+$ almost surely in Γ , with ϕ_β^- lower and ϕ_β^+ upper semi-continuous. If there is only one invariant graph ϕ_β , then $\lambda(\phi_\beta) = 0$. If there are two graphs, then $\lambda(\phi_\beta^-) > 0$ and $\lambda(\phi_\beta^+) < 0$.
- If $\beta > \beta_c$, then no f_β -invariant graph exists in Γ .

Remark 2.7. If there exist two invariant graphs at the critical parameter β_c , we speak of a *non-smooth saddle-node bifurcation*. The other case is referred to as a *smooth* bifurcation.

The main goal of this article is to provide natural conditions under which the occurrence of a non-smooth saddle-node bifurcation is guaranteed.

3. Statement of the main result and applications

We first collect a number of assumptions on the considered skew-products needed to formulate our main result. In order to both make the reader familiar with these assumptions and to demonstrate how they apply to some standard skew-product families, we explicitly show that they are satisfied by

$$f_\beta : \mathbb{T}^1 \times \mathbb{R} \rightarrow \mathbb{T}^1 \times \mathbb{R} \quad (*)$$

$$(\theta, x) \mapsto \left(\theta + \omega, \arctan(\alpha x) - \beta \cdot \frac{\pi}{4} (1 + \cos 2\pi\theta) \right),$$

for Diophantine $\omega \in \mathbb{T}^1$ and large enough α .

To guarantee the existence of a sink source orbit, we need to ensure that the respective orbit spends most of the positive times in regions of (vertical) expansion and most of the negative times in (vertically) contracting regions. For that reason, we assume the existence of both an interval of expansion $E = [e^-, e^+]$ and contraction $C = [c^-, c^+]$ with $e^+ < c^-$ and such that:

$$(A1) \quad \partial_x f_{\beta, \theta}(x) < \alpha_c \text{ for } (\theta, x) \in \mathbb{T}^d \times C;$$

$$(A2) \quad \partial_x f_{\beta, \theta}(x) > \alpha_e \text{ for } (\theta, x) \in \mathbb{T}^d \times E;$$

where $0 < \alpha_l < \alpha_c < 1 < \alpha_e < \alpha_u$ and

$$(A3) \quad \alpha_l < \partial_x f_{\beta, \theta}(x) < \alpha_u \text{ for all } (\theta, x) \in \mathbb{T}^d \times [e^-, c^+].$$

Instead of considering all of the phase space $\mathbb{T}^d \times X$, we restrict our analysis to the section $\mathbb{T}^d \times [e^-, c^+]$. Thus, e^- and c^+ play the roles of $\gamma^\mp$ in Theorem 2.6.

$$(A4) \quad f_{\beta, \theta}(c^+) \leq c^+ \text{ and } f_{\beta, \theta}(e^-) \leq e^-.$$

In Theorem 2.6, there exist two invariant graphs between γ^- and γ^+ for $\beta = 0$ and no invariant graphs for $\beta = 1$. In addition to (A4), we also assume (A5) to ensure this.

$$(A5) \quad f_{0, \theta}(c^-) \geq c^- \text{ for all } \theta \in \mathbb{T}^d \text{ and } f_{1, \theta}(c^+) \leq e^- \text{ for some } \theta \in \mathbb{T}^d.$$

Before formulating further assumptions, let us define the introduced quantities for (*) and see how (A1)–(A5) are verified in this particular case. Set $e^- := 0$, $e^+ := r/\alpha$ for some $r > 1$, $c^- := 1/r$ and $c^+ := \pi/2$. Note that $e^- < e^+ < c^- < c^+$ for large enough α . As

$$\partial_x f_{\beta, \theta}(x) = \frac{\alpha}{1 + (\alpha x)^2},$$

we get (A1)–(A3) with $\alpha_e, \alpha_c^{-1} = \alpha^{2/p}$ and $\alpha_u, \alpha_l^{-1} = \alpha^p$ for some fixed $p > 2$ if α is large enough. Further, (A4) is evident and (A5) holds trivially under the assumption of large enough α .

As in Theorem 2.6, we naturally assume monotone dependence on β .

$$(A6) \quad f_{(\cdot)}(\theta, x) \text{ is strictly decreasing for fixed } (\theta, x) \in \mathbb{T}^d \times [e^-, c^+].$$

Furthermore, we need the dependence on β to be smooth enough, that is, we suppose (A7) below.

$$(A7) \quad (\beta, \theta, x) \mapsto f_\beta(\theta, x) \text{ and } (\beta, \theta, x) \mapsto \partial_x f_\beta(\theta, x) \text{ as well as } (\beta, \theta, x) \mapsto \partial_\theta f_\beta(\theta, x) \text{ are continuous.}$$

Both assumptions are trivially fulfilled by (*). As we want to restrict ourselves to $\mathbb{T}^d \times [e^-, c^+]$, we exclude parameters β for which obviously every orbit leaves $\mathbb{T}^d \times [e^-, c^+]$. In other words, we only consider parameters not bigger than $\beta_+(0) := \min\{\beta \in [0, 1] \mid \exists \theta \in \mathbb{T}^d : f_{\beta, \theta}(c^+) = e^-\}$. On the other hand, as we want the sink-source orbit to basically

stay in the contracting region $\mathbb{T}^d \times C$ for negative times while we want it in the expanding region $\mathbb{T}^d \times E$ for positive times, we need to ensure that there is a connection between the two regions. Therefore, we only consider parameters β not too small in order to make it possible to jump from one region to the other. That is, we deal with parameters not smaller than $\beta_-(0) := \max\{\beta \in [0, \beta_+(0)] \mid \forall \theta \in \mathbb{T}^d : f_{\beta,\theta}(c^-) \geq e^+\}$. We don't need to compute $\beta_{\pm}(0)$ for (*) explicitly. Instead, it suffices to know that $\beta_{\pm}(0) \in (0, 1)$, which is true for obvious reasons.

Setting $\mathcal{B}(0) := [\beta_-(0), \beta_+(0)]$, we hence only consider $\beta \in \mathcal{B}(0)$ from now on. For each such β there is a so-called (first) *critical region*, $\mathcal{I}_{0,\beta} \subseteq \mathbb{T}^d$ such that outside of $\mathcal{I}_{0,\beta}$, orbits in the contracting region stay in the contracting region.

(A8) $f_{\beta,\theta}(x) \in C$ for all $x \in [e^+, c^+]$, $\theta \notin \mathcal{I}_{0,\beta}$.

By means of the monotonicity in (A3) and by (A4), this is equivalent to (A8) below.

(A8') $f_{\beta,\theta}^{-1}(x) \in E$ for all $x \in [e^-, c^-]$, $\theta \notin \mathcal{I}_{0,\beta} + \omega$.

Notice that a priori we did not assume invertibility of f_{β} . However, due to the inverse function theorem and (A3), we have that for a small open neighbourhood U of $\mathbb{T}^d \times [e^-, c^+]$ the map $(f_{\beta}|_U)^{-1}$ is well-defined and $\mathcal{C}^2$. We will refer to it simply as f_{β}^{-1} . Observe that it also verifies (A7).

In general, a natural choice for the critical region is given by

$$\mathcal{I}_{0,\beta} := \{\theta \in \mathbb{T}^d : f_{\beta,\theta}(e^+) \leq c^-\},$$

which verifies (A8) by definition. In the particular case of (*), this choice reads

$$\mathcal{I}_{0,\beta} := \left\{ \theta \in \mathbb{T}^1 : \cos 2\pi\theta \geq \frac{4}{\pi} \cdot (\arctan r - 1/r)/\beta - 1 \right\}. \quad (3.1)$$

The critical region $\mathcal{I}_{0,\beta}$ allows jumps from the contracting to the expanding region and vice versa. On the other hand, we also want the sink-source orbit to spend long times in the respective regions without jumping out too often, that is, we don't want $\mathcal{I}_{0,\beta}$ to be too big.

In (3.1), we see that by choosing large r , we can make $\mathcal{I}_{0,\beta}$ arbitrarily small for large enough α . This results from the fact that the second derivative $\partial_{\theta}^2 f_{\beta,\theta}(x) = \beta\pi^3 \cdot \cos 2\pi\theta$ is bounded away from 0 on the interval $\mathcal{I}_{0,\beta}$. In general, we thus assume there exists $s > 0$ such that

(A9) $\partial_{\vartheta}^2 f_{\beta,\theta}(x) > s$ for each $\vartheta \in \mathbb{S}^{d-1}$ and $\theta \in \mathcal{I}_{0,\beta}$, $x \in C$, $\beta \in \mathcal{B}(0)$,

(A10) $\mathcal{I}_{0,\beta}$ is closed and *convex* and $\mathcal{I}_{0,\beta} \subseteq \mathcal{I}_{0,\beta'}$ for $\beta \leq \beta'$.

To motivate further assumptions, we need to provide a rough sketch of how to prove the existence of a sink-source orbit. Assuming that $\mathcal{I}_{0,\beta}$ is small, there is a positive number M_0 such that the first M_0 forward and backward iterates of $\mathcal{I}_{0,\beta} + \omega$ under the base transformation (that is, under the rigid rotation with rotation vector ω) don't intersect, that is,

$$\mathcal{I}_{0,\beta} + \omega \cap \bigcup_{k=\pm 1, \dots, \pm M_0} (\mathcal{I}_{0,\beta} + (k+1)\omega) = \emptyset.$$

If this is true, $f_{\beta}^l(\theta, x)$ never leaves the contracting region for $\theta \in \mathcal{I}_{0,\beta} - (M_0 - 1)\omega$, $x \in C$ and $l = 0, \dots, M_0 - 1$, while $f_{\beta}^{-l}(\theta, x)$ never leaves the expanding region for $\theta \in \mathcal{I}_{0,\beta} + (M_0 + 1)\omega$, $x \in E$ and $l = 0, \dots, M_0$, due to (A8) and (A8'). However, $f_{\beta}^{M_0-1}(\theta, x)$ might jump into the expanding region under the action of f_{β} or even fall into

the set $f_\beta^{-M_0}(\mathcal{I}_{0,\beta} + (M_0 + 1)\omega, E)$. In the latter case, $f_\beta^{M_0-1}(\theta, x)$ is a first candidate for a sink-source orbit as it stays in the expanding region $M_0 + 1$ times while its backward iterates stay in the contracting region $M_0 - 1$ times.

The projection of the set of all such sink-source orbit candidates to the base $\mathbb{T}^d$ is denoted by $\mathcal{I}_{1,\beta}$. As in the case of $\mathcal{I}_{0,\beta}$, we need that $\mathcal{I}_{1,\beta}$ is small enough to guarantee that it visits itself with an even smaller frequency than $\mathcal{I}_{0,\beta}$. To that end, we need that the second derivatives of $\phi^\pm(\theta) := f_{\beta,\theta-M_0\omega}^{M_0}(c^\pm)$ and $\psi^\pm(\theta) := f_{\beta,\theta+M_0\omega}^{-M_0}(e^\pm)$ (for $\theta \in \mathcal{I}_{0,\beta} + \omega$) with respect to θ are basically bounded from below by the lower bound s in (A9)). This amounts to keeping all the other derivatives of f_β and its inverse small. Let $S > 0$ be such that:

$$(A11) \quad |\partial_\vartheta f_{\beta,\theta}(x)| < S \text{ for all } (\theta, x) \in \mathbb{T}^d \times [e^-, c^+] \text{ and } \vartheta \in \mathbb{S}^{d-1};$$

$$(A12) \quad |\partial_\vartheta^2 f_{\beta,\theta}(x)| < S^2 \text{ for all } (\theta, x) \in \mathbb{T}^d \times [e^-, c^+] \text{ and } \vartheta \in \mathbb{S}^{d-1};$$

$$(A13) \quad |\partial_\vartheta \partial_x f_{\beta,\theta}(x)| < \begin{cases} S\alpha_c & \text{for } (\theta, x) \in \mathbb{T}^d \times C, \\ S\alpha_u^2 & \text{for } (\theta, x) \in \mathbb{T}^d \times [e^-, c^-), \end{cases} \text{ for each } \vartheta \in \mathbb{S}^{d-1}.$$

Further, suppose

$$(A14) \quad |\partial_x^2 f_{\beta,\theta}(x)| < \begin{cases} \alpha_c & \text{for } (\theta, x) \in \mathbb{T}^d \times C, \\ \alpha_u^2 & \text{for } (\theta, x) \in \mathbb{T}^d \times [e^-, c^-). \end{cases}$$

For the derivatives of the inverse, we get some of the above estimates by means of the inverse function theorem. However, we additionally need

$$(A15) \quad |\partial_x^2 f_{\beta,\theta}^{-1}(x)| < \alpha_e^{-1} \text{ for each } \theta \notin \mathcal{I}_{0,\beta} + \omega \text{ and } x \in E,$$

$$(A16) \quad |\partial_\vartheta \partial_x f_{\beta,\theta}^{-1}(x)| < S\alpha_e^{-1} \text{ for each } \theta \notin \mathcal{I}_{0,\beta} + \omega, x \in E \text{ and } \vartheta \in \mathbb{S}^{d-1}.$$

Coming back to (*), we get (A11) and (A12) by setting $S := \max_{\beta,\theta,x} \partial_\theta f_{\beta,\theta}(x) = \pi^2/2$.

(A13) is trivial, as mixed derivatives vanish. With

$$\partial_x^2 f_{\beta,\theta}(x) = \frac{-2\alpha^3 x}{(1 + (\alpha x)^2)^2},$$

we get $|\partial_x^2 f_{\beta,\theta}(x)| < \alpha^{-2/p}$ for big enough α and $x \in C$. Further, basic calculus yields $|\partial_x^2 f_{\beta,\theta}(x)| \leq \partial_x^2 f_{\beta,\theta}(\sqrt{1/(3\alpha^2)}) = \mathcal{O}(\alpha^2)$ as $\alpha \rightarrow \infty$. This shows (A14) for big enough α . If $x \in E$ and $\theta \notin \mathcal{I}_{0,\beta} + \omega$, we moreover have

$$\begin{aligned} \partial_x^2 f_{\beta,\theta}^{-1}(x) &= 2/\alpha \cdot \frac{\sin(x + \beta \cdot (\pi/4)(\cos 2\pi(\theta - \omega) + 1))}{\cos^3(x + \beta \cdot (\pi/4)(\cos 2\pi(\theta - \omega) + 1))}, \\ \partial_\theta \partial_x f_{\beta,\theta}^{-1}(x) &= -\beta\pi^2/\alpha \cdot \frac{\sin 2\pi(\theta - \omega) \cdot \sin(x + \beta \cdot (\pi/4)(\cos 2\pi(\theta - \omega) + 1))}{\cos^3(x + \beta \cdot (\pi/4)(\cos 2\pi(\theta - \omega) + 1))}. \end{aligned}$$

As $\theta \notin \mathcal{I}_{0,\beta} + \omega$, (3.1) yields $\beta \cdot (\pi/4)(\cos 2\pi(\theta - \omega) + 1) < \arctan r - c^-$ which proves (A15) and (A16) for large enough α , since $0 \leq x \leq r/\alpha$.

We are now in a position to state the main theorem of this article.

THEOREM 3.1. *Suppose ω is Diophantine of type $(\mathcal{C}, \eta)$ and $(f_\beta)_{\beta \in [0,1]}$ satisfies (A1)–(A16). Let there be $p \geq \sqrt{2}$, $\alpha > 1$ with*

$$\alpha_c^{-1} = \alpha_e = \alpha^{2/p}, \quad \alpha_l^{-1} = \alpha_u = \alpha^p.$$

Then there exist strictly positive constants

$$\varepsilon_0 = \varepsilon_0(p, \mathcal{C}, \eta) \quad \text{and} \quad \alpha_0 = \alpha_0(s, S, p, |C|, |E|, \mathcal{C}, \eta)$$

such that if $|\mathcal{I}_{0,\beta_+(0)}| < \varepsilon_0$ and $\alpha > \alpha_0$, there is $\beta_c \in [0, 1]$ such that f_{β_c} has a sink-source orbit, and hence an SNA and an SNR in $\mathbb{T}^d \times [e^-, c^+]$.

Note that together with the previous discussion, this theorem proves the occurrence of a non-smooth bifurcation for (*).

Remark 3.2.

- (i) α_0 can be chosen to be monotonously increasing in $|C|$ and $|E|$.
- (ii) The conjugacy $(\theta, x) \mapsto (\theta, -x)$ and the parametrisation $\beta \mapsto 1 - \beta$ yield a symmetric version of the theorem if the contracting region is below the expanding one, that is, if $c^+ < e^-$.

Theorem 3.1 is proved in §4 by showing the existence of a critical parameter $\beta_c \in \mathcal{B}(0)$ for which f_{β_c} has a sink-source orbit in $\mathbb{T}^d \times [e^-, c^+]$. By means of Theorem 2.4, this implies the statement. It is important to note that the assumptions of Theorem 3.1 are stable under $\mathcal{C}^2$ -small perturbations of the fibre maps $f_{\beta,\theta}$ which respect (A6) and (A7). This is the main advantage over previous results in this direction, which establish the existence of an SNA only under comparably strong technical constraints: in [12, Theorem 2.7] it is necessary to assume the existence of a ‘sharp peak’ for the maps $\phi^\pm$, which implies non-differentiability of f_β with respect to the base coordinates[†].

An important step towards the understanding of the creation of SNAs was the verification of a non-smooth saddle-node bifurcation for the Harper map

$$(\theta, x) \mapsto \left(\theta + \omega, \arctan\left(\frac{-1}{\tan(x) - E + \lambda V(\theta)}\right) \right),$$

which is closely related to the discrete quasi-periodic Schrödinger equation. In [3] it is shown that if the *potential* V is $\mathcal{C}^2$ and if it assumes its unique global maximum at a point with non-vanishing second derivative, then we observe a non-smooth saddle node-bifurcation upon a decrease of E if λ is large enough.

The geometric idea of our proof is inspired by the proof in [3] as can be readily seen from the pictures in Figure 2. It is thus not surprising that we can recover Bjerklov’s result with the same regularity assumptions[‡]. However, as we don’t restrict ourselves to fibre maps of a particular shape, more work is needed in order to get control over the sink-source orbit.

Despite the fact that (A1)–(A16) seem rather technical, they just capture the main qualitative properties of some standard examples which possess an SNA and turn out to be flexible enough to treat different skew-product families at the same time. We have seen that (*) verifies the assumptions of Theorem 3.1. As a generalisation of the arctan-family (*), for each $q > 1$ we can apply Theorem 3.1 to

$$(\theta, x) \mapsto \left(\theta + \omega, h_q(\alpha x) - \beta \cdot \frac{h_q(\infty)}{2}(1 + \cos 2\pi\theta) \right),$$

[†] Note that the positive lower bound for $\partial_\theta^2 f_{\beta,\theta}(x)$ in (A9) can be understood as a replacement for this sharp peak assumption in the respective statement in [12].

[‡] The application of Theorem 3.1 to the Harper map works by means of a similar argument to that in [12, §2.4.2]. However, it is necessary to control the dependence of α_0 on s and S in this particular case. To that end, we provide a slightly different formulation of the above theorem (cf. Theorem 4.18), which specifies the relationship between α and s as well as S in an appropriate way.

where $h_q(x) := \operatorname{sgn}(x) \cdot \tilde{h}_q(|x|)$ with $\tilde{h}_q(x) := \int_0^x (1 + \zeta^q)^{-1} d\zeta$, which can be seen similarly as for (*). Analogously, we obtain a non-smooth saddle-node bifurcation for the family

$$(\theta, x) \mapsto h_q(\alpha x) - 2\beta - \frac{1 + \sin 2\pi\theta}{2},$$

which has been considered numerically for $q = 2$ in [1], for example.

Remark 3.3. The assumptions that $\alpha_c^{-1} = \alpha_e = \alpha^{2/p}$ and $\alpha_l^{-1} = \alpha_u = \alpha^p$ are made only for technical reasons. They basically originate from the fact that we defined $\mathcal{I}_{1,\beta}$ in a symmetric way; that is, we considered the intersection of the M_0^- th iterate of $\mathcal{I}_0 - (M_0^- - 1)\omega \times C$ and the M_0^+ th inverse iterate of $\mathcal{I}_0 + (M_0^+ + 1)\omega \times E$ with $M_0^+ = M_0^-$ (cf. Definition 4.1). We believe that by allowing different relations between M_0^+ and M_0^- , we could also allow different scaling behaviour in order to apply a statement similar to Theorem 3.1 to $(\theta, x) \mapsto (\theta + \omega, \tanh(\alpha x) - \beta(1 + \cos(2\pi\theta)))$, for example, where the ratio of α_l^{-1}/α_u grows exponentially with α .

Remark 3.4. Combining Theorems 2.2, 2.6, and 3.1, we straightforwardly get conditions which guarantee the occurrence of a non-smooth saddle-node bifurcation. However, it is worth mentioning that besides some minor technical hypothesis, the convexity assumption of Theorem 2.6 is not needed in Theorem 3.1. In other words: the existence of an SNA is in a sense independent of the saddle-node bifurcation framework.

4. Proof of Theorem 3.1

In this section, we prove Theorem 3.1 by showing that there is a point (θ, x) whose positive iterates mostly stay in the expanding region, while its negative iterates mostly stay in the contracting region. This can be achieved if the frequency of the jumps from one region to the other is small enough, which is the idea behind the inductive assumptions $(\mathcal{F}1)_n$ and $(\mathcal{F}2)_n$ (§4.1). These are basically hypotheses on the size of the inductively defined *critical intervals* $\mathcal{I}_n$. By a geometrical argument, we will get upper bounds for these quantities in §4.2. In §4.3, we eventually show that these upper bounds decrease fast enough to guarantee the existence of an SNA.

4.1. Combinatorial considerations: the basic mechanism. We make use of (A1)–(A4) and (A8) to estimate the vertical growth rate of orbits which converge to a sink-source orbit. In order to achieve this, we need to assume some additional inductive assumptions. The verification of these additional assumptions is the goal of the subsequent sections. As a matter of fact, the statements of this section are basically provided in [3, 15], already. For the convenience of the reader and as there are some subtle technical differences, we nevertheless include some of the proofs.

In the following, let $(M_n)_{n \in \mathbb{N}_0}, (K_n)_{n \in \mathbb{N}_0} \in \mathbb{N}^{\mathbb{N}_0}$ be strictly increasing sequences (independent of β) with $M_0 \geq 2$ and $M_n \leq 2K_{n-1}M_{n-1} - 2$ for all $n \in \mathbb{N}$. Set $M_{-1} = 0$.

Definition 4.1. Suppose we have already defined the n th *critical region* $\mathcal{I}_{n,\beta}$. Set:

- $\mathcal{A}_{n,\beta} := (\mathcal{I}_{n,\beta} - (M_n - 1)\omega) \times C$;
- $\mathcal{B}_{n,\beta} := (\mathcal{I}_{n,\beta} + (M_n + 1)\omega) \times E$;
- $\mathcal{I}_{n+1,\beta} := \pi_1(f_\beta^{M_n-1}(\mathcal{A}_{n,\beta}) \cap f_\beta^{-(M_n+1)}(\mathcal{B}_{n,\beta}))$.

Remark 4.2. It is obvious that $\mathcal{I}_{n+1,\beta} \subseteq \mathcal{I}_{n,\beta}$ ($n \in \mathbb{N}_0$). However, note that $\mathcal{I}_{n+1,\beta}$ might be empty even if $\mathcal{I}_{n,\beta} \neq \emptyset$.

For fixed $N \in \mathbb{N}$, we will only consider such $\beta \in \mathcal{B}(0)$ with $f_{\beta, \theta_{-(M_n-1)}}^{M_n-1}(c^+) \geq f_{\beta, \theta_{M_n+1}}^{-(M_n+1)}(e^-)$ for each $\theta \in \mathcal{I}_n$ and $0 \leq n \leq N-1$. We denote the set of these β by $\tilde{\mathcal{B}}(N)$ and set $\tilde{\mathcal{B}}(0) := \mathcal{B}(0)$.

Occasionally, we might suppress the index β . For $n \in \mathbb{N}_0$, set

$$\begin{aligned}\mathcal{Z}_n^- &:= \bigcup_{j=0}^n \bigcup_{l=-(M_j-2)}^0 \mathcal{I}_j + l\omega; \\ \mathcal{Z}_n^+ &:= \bigcup_{j=0}^n \bigcup_{l=1}^{M_j} \mathcal{I}_j + l\omega; \\ \mathcal{V}_n &:= \bigcup_{j=0}^n \bigcup_{l=1}^{M_j+1} \mathcal{I}_j + l\omega; \\ \mathcal{W}_n &:= \bigcup_{j=0}^n \bigcup_{l=-(M_j-1)}^0 \mathcal{I}_j + l\omega.\end{aligned}$$

Moreover, set $\mathcal{I}_{-1,\beta} = \mathcal{I}_{0,\beta}$ and $\mathcal{Z}_{-1}^-, \mathcal{Z}_{-1}^+, \mathcal{V}_{-1}, \mathcal{W}_{-1} = \emptyset$.

In order to be able to control an orbit, we do not want it to visit the critical regions too often. We therefore need to assume that the critical regions are small enough.

Definition 4.3. We say f_β verifies $(\mathcal{F}1)_n$ and $(\mathcal{F}2)_n$ if:

$$(\mathcal{F}1)_n \quad \mathcal{I}_{j,\beta} \cap \bigcup_{k=1}^{2K_j M_j} \mathcal{I}_{j,\beta} + k\omega = \emptyset;$$

$$(\mathcal{F}2)_n \quad (\mathcal{I}_{j,\beta} - (M_j - 1)\omega) \cup \mathcal{I}_{j,\beta} + (M_j + 1)\omega \cap (\mathcal{V}_{j-1} \cup \mathcal{W}_{j-1}) = \emptyset;$$

for $j = 0, \dots, n$ and $n \in \mathbb{N}_0$. If f_β satisfies both $(\mathcal{F}1)_n$ and $(\mathcal{F}2)_n$, we say f_β satisfies $(\mathcal{F})_n$. It is convenient to set $(\mathcal{F})_{-1}$ to be true.

For $\theta \in \mathbb{T}^d$, we denote by $\mathcal{L}_m, \mathcal{R}_m \in \mathbb{N}_0 \cup \{\infty\}$ the smallest integers l, r with $\theta_l \in \mathcal{I}_m$ and $\theta_{-r} \in \mathcal{I}_m + \omega$, respectively.

LEMMA 4.4. (Cf. [15, Lemma 3.4]) *Let $n \in \mathbb{N}_0$. Suppose f_β satisfies $(\mathcal{A}3)$, $(\mathcal{A}4)$ and $(\mathcal{A}8)$ as well as $(\mathcal{F})_{n-1}$ with $\beta \in \tilde{\mathcal{B}}(n)$ and assume*

$$\begin{cases} x \in C, \\ \theta \notin \mathcal{Z}_{n-1}^-. \end{cases} \quad (\mathcal{B}1)_n$$

Furthermore, let $0 < \mathcal{L}^{(1)} < \dots < \mathcal{L}^{(N)} = \mathcal{L}_n$ be all those times $m \leq \mathcal{L}_n$ for which $\theta + m\omega \in \mathcal{I}_{n-1}$. Then $(\theta_{\mathcal{L}^{(i)}+M_{n-1}+2}, x_{\mathcal{L}^{(i)}+M_{n-1}+2})$ satisfies $(\mathcal{B}1)_n$ for each $i = 1, \dots, N-1$ and the following implication holds.

$$x_k \notin C \Rightarrow \theta_k \in \mathcal{V}_{n-1} \quad \text{and} \quad x_k \in [e^-, c^-] \quad (k = 1, \dots, \mathcal{L}_n). \quad (\mathcal{C}1)_n$$

Analogously backwards: instead of $(\mathcal{B}1)_n$, assume

$$\begin{cases} x \in E, \\ \theta \notin \mathcal{Z}_{n-1}^+, \end{cases} \quad (\mathcal{B}2)_n$$

and let $0 < \mathcal{R}^{(1)} < \dots < \mathcal{R}^{(N)} = \mathcal{R}_n$ be all those times $m \leq \mathcal{R}_n$ for which $\theta - m\omega \in \mathcal{I}_{n-1} - \omega$. Then $(\theta_{-\mathcal{R}^{(i)}-M_{n-1}}, x_{-\mathcal{R}^{(i)}-M_{n-1}})$ satisfies $(\mathcal{B}2)_n$ for each $i = 1, \dots, N-1$ and the following implication holds.

$$x_{-k} \notin E \Rightarrow \theta_{-k} \in \mathcal{W}_{n-1} \quad \text{and} \quad x_{-k} \in [e^+, c^+] \quad (k = 1, \dots, \mathcal{R}_n). \quad (\mathcal{C}2)_n$$

Proof. We only consider the forward case; the other case works similarly. Note that for $n = 0$ the statement is true due to (A8).

Assume the statement holds for $n_0 \in \mathbb{N}_0$ and assume (θ, x) satisfies $(\mathcal{B}1)_{n_0+1}$. Trivially, $(\mathcal{B}1)_{n_0+1}$ implies $(\mathcal{B}1)_{n_0}$ such that $x_k \notin C \Rightarrow \theta_k \in \mathcal{V}_{n_0-1}$ and $x_k \in [e^-, c^+]$ for $k \leq \mathcal{L}^{(1)}$. Notice that $(\mathcal{I}_{n_0} - (M_{n_0} - 1)\omega) \cap \mathcal{V}_{n_0-1} = \emptyset$ because of $(\mathcal{F}2)_{n_0}$. Hence, $(\theta_{\mathcal{L}^{(1)}-(M_{n_0}-1)}, x_{\mathcal{L}^{(1)}-(M_{n_0}-1)}) \in \mathcal{A}_{n_0}$ due to $(\mathcal{C}1)_{n_0}$. As $\beta \in \tilde{\mathcal{B}}(n_0 + 1)$, we further have $f_{\beta, \theta_{\mathcal{L}^{(1)}-M_{n_0}+1}}^{M_{n_0}-1}(c^+) \geq f_{\beta, \theta_{\mathcal{L}^{(1)}+M_{n_0}+1}}^{-(M_{n_0}+1)}(e^-)$. If we had $x_{\mathcal{L}^{(1)}} \leq f_{\beta, \theta_{\mathcal{L}^{(1)}+M_{n_0}+1}}^{-(M_{n_0}+1)}(e^-)$, this would imply the existence of $y \in [x_{\mathcal{L}^{(1)}-M_{n_0}+1}, c^+] \subseteq [c^-, c^+]$ with $f_{\beta}^{M_{n_0}-1}(\theta_{\mathcal{L}^{(1)}-M_{n_0}+1}, y) \in f_{\beta}^{-(M_{n_0}+1)}(\{\theta_{\mathcal{L}^{(1)}+M_{n_0}+1}\} \times E)$ meaning that $\theta_{\mathcal{L}^{(1)}} \in \mathcal{I}_{n_0+1, \beta}$, which contradicts the assumptions. Therefore, $x_{\mathcal{L}^{(1)}} \geq f_{\beta, \theta_{\mathcal{L}^{(1)}+M_{n_0}+1}}^{-(M_{n_0}+1)}(e^-)$. By (A4) and the monotonicity, we thus have $x_k \in [e^-, c^+]$ for $k = \mathcal{L}^{(1)}, \dots, \mathcal{L}^{(1)} + M_{n_0} + 1$. Now, $x_{\mathcal{L}^{(1)}+M_{n_0}+1} \notin E$, since otherwise again $\theta_{\mathcal{L}^{(1)}} \in \mathcal{I}_{n_0+1}$, by definition of $\mathcal{I}_{n_0+1}$. (A8) and $(\mathcal{F}2)_{n_0}$ hence yield $x_{\mathcal{L}^{(1)}+M_{n_0}+2} \in C$. By $(\mathcal{F})_{n_0}$, we get that $(\mathcal{I}_{n_0} + (M_{n_0} + 2)\omega) \cap \mathcal{Z}_{n_0}^- = \emptyset$. Thus, $(\theta_{\mathcal{L}^{(1)}+M_{n_0}+2}, x_{\mathcal{L}^{(1)}+M_{n_0}+2})$ verifies $(\mathcal{B}1)_{n_0+1}$. The statement follows by induction. $\square$

Remark 4.5. Suppose $(\mathcal{F})_{n-1}$ and $(\mathcal{F}2)_n$ hold true. As $M_n \leq 2K_{n-1}M_{n-1} - 2$ and $\mathcal{I}_n \subseteq \mathcal{I}_{n-1}$, we have that $(\theta, x) \in \mathcal{A}_n$ satisfies $(\mathcal{B}1)_n$ with $\mathcal{L}_n = M_n - 1$ and $(\theta, x) \in \mathcal{B}_n$ satisfies $(\mathcal{B}2)_n$ with $\mathcal{R}_n = M_n$.

COROLLARY 4.6. Let $n \in \mathbb{N}_0$. Suppose f_β satisfies (A3), (A4), (A8) and $(\mathcal{F})_{n-1}$ as well as $(\mathcal{F}2)_n$ with $\beta \in \tilde{\mathcal{B}}(n)$. Then

$$f_\beta^{M_n-1}(\mathcal{A}_n) \subseteq \mathcal{I}_{n, \beta} \times C \quad \text{and} \quad f_\beta^{-M_n}(\mathcal{B}_n) \subseteq (\mathcal{I}_{n, \beta} + \omega) \times E.$$

Proof. Note that $\mathcal{I}_{n, \beta} - \omega \cap \mathcal{V}_n = \emptyset$, because of $(\mathcal{F}1)_{n-1}$ and $M_n \leq 2K_{n-1}M_{n-1} - 2$. By means of Remark 4.5, the first inclusion follows from Lemma 4.4. The second one follows similarly. $\square$

COROLLARY 4.7. (Cf. [15, Corollary 3.7]) Let $N > n \in \mathbb{N}_0$. Suppose f_β satisfies (A3), (A4), (A8) as well as $(\mathcal{F}1)_{N-1}$, $(\mathcal{F}2)_N$ with $\beta \in \tilde{\mathcal{B}}(N)$. Then $f_\beta^{M_N-M_n}(\mathcal{A}_N) \subseteq (\mathcal{I}_n - (M_n - 1)\omega) \times (c^-, c^+) \subseteq \mathcal{A}_n$ and $f_\beta^{-M_N+M_n}(\mathcal{B}_N) \subseteq (\mathcal{I}_n + (M_n + 1)\omega) \times [e^-, e^+] \subseteq \mathcal{B}_n$.

Proof. Since $\mathcal{I}_{n+1} - (M_n - 1)\omega \cap \mathcal{V}_n = \emptyset$, Lemma 4.4 yields $f_\beta^{M_{n+1}-M_n}(\mathcal{A}_{n+1}) \subseteq \mathcal{A}_n$. Due to (A8), the proof of Lemma 4.4 even yields the slightly stronger inclusion $f_\beta^{M_{n+1}-M_n}(\mathcal{A}_{n+1}) \subseteq (\mathcal{I}_n - (M_n - 1)\omega) \times (c^-, c^+]$. Now, the first result follows by induction. The other relation follows similarly. $\square$

By means of the next statement, we can control the time spent in the contracting and expanding region, respectively. For $n, N \in \mathbb{N}$ set

$$\begin{aligned}\mathcal{P}_n^N(\theta, x) &:= \#\{l \in [n, N-1] \cap \mathbb{N}_0 : x_l \in C \text{ and } \theta_l \notin \mathcal{I}_0\}, \\ \mathcal{Q}_n^N(\theta, x) &:= \#\{l \in [n, N-1] \cap \mathbb{N}_0 : x_{-l} \in E \text{ and } \theta_{-l} \notin \mathcal{I}_0 + \omega\}.\end{aligned}$$

Set

$$b_0 := 1, \quad b_n := \left(1 - \frac{1}{K_{n-1}}\right) b_{n-1} \quad (n \in \mathbb{N}).$$

LEMMA 4.8. (Cf. [15, Lemma 3.8]) *Let $n \in \mathbb{N}_0$. Suppose f_β satisfies (A3), (A4), (A8) and $(\mathcal{F})_{n-1}$ with $\beta \in \mathcal{B}(n)$. Furthermore, assume $(\mathcal{B}1)_n$ and let $0 < \mathcal{L}^{(1)} < \dots < \mathcal{L}^{(N)} = \mathcal{L}_n$ be as in Lemma 4.4. Then, for each $i = 0, \dots, N$, we have*

$$P_k^{\mathcal{L}^{(i)}}(\theta, x) \geq b_n(\mathcal{L}^{(i)} - k) \quad (k = 0, \dots, \mathcal{L}^{(i)}).$$

Analogously backwards: instead of $(\mathcal{B}1)_n$, assume $(\mathcal{B}2)_n$ and let $0 < \mathcal{R}^{(1)} < \dots < \mathcal{R}^{(N)} = \mathcal{R}_n$ be as in Lemma 4.4. Then, for each $i = 0, \dots, N$, we have

$$Q_k^{\mathcal{R}^{(i)}}(\theta, x) \geq b_n(\mathcal{R}^{(i)} - k) \quad (k = 0, \dots, \mathcal{R}^{(i)}).$$

Remark 4.9. For the present work (that is, in order to show the existence of an SNA) it suffices to have the lower bound for $P_k^{\mathcal{L}_n}(\theta, x)$ only. Nevertheless, the estimates for $P_k^{\mathcal{L}^{(i)}}(\theta, x)$ will be needed in order to study further properties of the SNA [6].

Proof. We consider the first inequality; the second one follows similarly. For $n = 0$, the statement follows from (A8).

Assume the statement is true for $n = n_0$ and assume (θ, x) verifies $(\mathcal{B}1)_{n_0+1}$. Due to Lemma 4.4, we have that $(\theta_{\mathcal{L}^{(i)}+M_{n_0}+2}, x_{\mathcal{L}^{(i)}+M_{n_0}+2})$ satisfies $(\mathcal{B}1)_{n_0+1}$ for $i = 1, \dots, N-1$. By the induction hypothesis we thus get the desired estimate for $P_k^{\mathcal{L}^{(i)}}(\theta, x)$ as long as $k \in [\mathcal{L}^{(i)} + M_{n_0} + 2, \mathcal{L}^{(i+1)}]$ for some $0 \leq i \leq N-1$ or as $i = 1$ and $k \in [0, \mathcal{L}^{(1)} - 1]$.

Moreover, by $(\mathcal{F}1)_{n_0}$ we have

$$\mathcal{L}^{(i+1)} - \mathcal{L}^{(i)} \geq 2K_{n_0}M_{n_0}. \quad (4.1)$$

Hence, for all $k \in [\mathcal{L}^{(i)}, \mathcal{L}^{(i)} + M_{n_0} + 1]$ we get

$$\begin{aligned}\mathcal{P}_k^{\mathcal{L}^{(i+1)}}(\theta, x) &\geq \mathcal{P}_{\mathcal{L}^{(i)}+M_{n_0}+2}^{\mathcal{L}^{(i+1)}}(\theta, x) \\ &\geq b_{n_0}(\mathcal{L}^{(i+1)} - (\mathcal{L}^{(i)} + M_{n_0} + 2)) \geq b_{n_0}(\mathcal{L}^{(i+1)} - \mathcal{L}^{(i)} - 2M_{n_0}) \\ &\stackrel{(4.1)}{\geq} b_{n_0+1}(\mathcal{L}^{(i+1)} - \mathcal{L}^{(i)}) \geq b_{n_0+1}(\mathcal{L}^{(i+1)} - k).\end{aligned}$$

Altogether, with $j := \min\{l = 1, \dots, N : \mathcal{L}^{(l)} \geq k\}$ and $N \geq i \geq j$ we therefore have

$$\begin{aligned}P_k^{\mathcal{L}^{(i)}}(\theta, x) &= P_k^{\mathcal{L}^{(j)}}(\theta, x) + \sum_{l=j}^i P_{\mathcal{L}^{(l)}}^{\mathcal{L}^{(l+1)}}(\theta, x) \geq b_{n_0+1} \left(\mathcal{L}^{(j)} - k + \sum_{l=j}^i \mathcal{L}^{(l+1)} - \mathcal{L}^{(l)} \right) \\ &= b_{n_0+1}(\mathcal{L}^{(i)} - k).\end{aligned}$$

□

The following two results can be proved in a similar way to the respective statements in [15].

COROLLARY 4.10. (Cf. [15, Corollary 3.9]) *Let $n \in \mathbb{N}_0$. Suppose f_β satisfies (A3), (A4), (A8) and $(\mathcal{F})_{n-1}$ as well as $(\mathcal{F}2)_n$ with $\beta \in \tilde{\mathcal{B}}(n)$. Further, assume (A1) and let $(\theta, x) \in f_\beta^{M_n}(\mathcal{A}_n)$. Then*

$$\partial_x f_{\beta, \theta}^{-k}(x) \geq (\alpha_c^{b_n} \alpha_u^{1-b_n})^{-k} \quad (0 \leq k \leq M_n).$$

Analogously, instead of (A1) assume (A2) and let $(\theta, x) \in f_\beta^{-M_n}(\mathcal{B}_n)$. Then

$$\partial_x f_{\beta, \theta}^k(x) \geq (\alpha_e^{b_n} \alpha_l^{1-b_n})^k \quad (0 \leq k \leq M_n).$$

Define:

- $b := \lim_{n \rightarrow \infty} b_n$;
- $\alpha_- := \alpha_c^b \alpha_u^{1-b}$;
- $\alpha_+ := \alpha_e^b \alpha_l^{1-b}$.

PROPOSITION 4.11. (Cf. [15, Proposition 3.10]) *Suppose f_β satisfies (A1)–(A4) as well as (A8) and for each $n \in \mathbb{N}$ we have $f_\beta^{M_n}(\mathcal{A}_n) \cap f_\beta^{-M_n}(\mathcal{B}_n) \neq \emptyset$. Moreover, assume $(\mathcal{F})_n$ holds for all $n \in \mathbb{N}$, α_-^{-1} , $\alpha_+ > 1$ and $\beta \in \bigcap_{n \in \mathbb{N}} \tilde{\mathcal{B}}(n) \neq \emptyset$. Then there exists a sink-source orbit in $\mathbb{T}^d \times [e^-, c^+]$ and hence an SNA and an SNR. More precisely,*

$$\{(\theta, x) \in \mathbb{T}^d \times X : (\theta, x) \text{ is a sink-source orbit}\} \supseteq \bigcap_{n \in \mathbb{N}} (f_\beta^{M_n}(\mathcal{A}_n) \cap f_\beta^{-M_n}(\mathcal{B}_n)) \neq \emptyset.$$

4.2. *Geometric considerations.* In this paragraph, we get an upper bound for the size of the n th critical region $\mathcal{I}_{n, \beta}$. So far, we have dealt with $\beta \in \tilde{\mathcal{B}}(n)$ in order to guarantee that the respective orbits stay in the strip $\mathbb{T}^d \times [e^-, c^+]$. Due to the monotonicity in β (provided by (A6)), this amounts to only considering small enough β . On the other hand, $\tilde{\mathcal{B}}(n)$ also contains parameters β which are *too small* such that $\mathcal{I}_{n, \beta} = \emptyset$, which is not desirable either. In order to exclude these parameters as well, we define the set of *admissible parameters* up to order $n \in \mathbb{N}$ by

$$\begin{aligned} \mathcal{B}(n) &:= \{\beta \in \tilde{\mathcal{B}}(n) : f_{\beta, \theta_{-(M_l-1)}}^{M_l-1}(c^-) \leq f_{\beta, \theta_{M_l+1}}^{-(M_l+1)}(e^+) \text{ for some } \theta \in \mathcal{I}_l \text{ and } 0 \leq l \leq n-1\} \\ &= \{\beta \in \tilde{\mathcal{B}}(n) : \mathcal{I}_{l, \beta} \neq \emptyset \text{ for } 0 \leq l \leq n\}, \end{aligned}$$

where we assume $(M_l)_{l=0, \dots, n-1}$ to be given.

PROPOSITION 4.12. *Suppose $(f_\beta)_{\beta \in [0, 1]}$ satisfies (A6) and (A10) and let $\beta < \beta' \in \tilde{\mathcal{B}}(n)$ for some $n \in \mathbb{N}_0$. Then*

$$\mathcal{I}_{n, \beta} \subseteq \mathcal{I}_{n, \beta'}. \quad (4.2)$$

In particular, this implies that $\mathcal{B}(n)$ is an interval.

Proof. For $n = 0$, (4.2) holds by (A10). Assume (4.2) is true for some $n \in \mathbb{N}_0$. For $\beta \in \tilde{\mathcal{B}}(n+1)$, we know $\theta \in \mathcal{I}_{n+1, \beta}$ if and only if $0 \geq f_{\beta, \theta_{-(M_n-1)}}^{M_n-1}(c^-) - f_{\beta, \theta_{M_n+1}}^{-(M_n+1)}(e^+)$. Since $f_{(\cdot)}(\theta, x)$ is non-increasing, $f_{(\cdot), \theta_{-(M_n-1)}}^{M_n-1}(c^-) - f_{(\cdot), \theta_{M_n+1}}^{-(M_n+1)}(e^+)$ is non-increasing, too. Hence, $\theta \in \mathcal{I}_{n+1, \beta}$ implies $\theta \in \mathcal{I}_{n+1, \beta'}$. Now, (4.2) follows by induction. $\square$

FIGURE 2. (a) The geometric idea behind the proof of Lemma 4.13; (b) $\mathcal{I}_{n+1,\beta_-(n+1)}$ is degenerate, with $\beta_-(n+1) = \min \mathcal{B}(n+1)$; (c) $\beta_+(n+1) = \max \mathcal{B}(n+1)$ is the largest parameter such that $\mathcal{I}_{n+1,\beta_+(n+1)}$ is connected. (d) An upper bound for the length of $\mathcal{I}_{n,\beta}$. Note that h only touches $\phi_{n,\beta}^-$ in this picture for illustrative reasons.

Up to now, we basically used monotonicity in β in order to investigate the set of admissible parameters. In order to guarantee that $\mathcal{B}(n)$ is not empty and to control the size of the critical regions $\mathcal{I}_{n,\beta}$, we need subtler geometric information. The intuitive idea of the argument for the smallness of $\mathcal{I}_{n,\beta}$ can be seen by considering $\mathcal{I}_{1,\beta}$: as $f_\beta^j(\mathcal{A}_{0,\beta})$ stays in the contracting region for $j = 0, \dots, M_0 - 1$, the iterates of $\mathcal{A}_{0,\beta}$ become thinner and thinner horizontal strips with each step of the iteration until they meet $\mathcal{I}_{0,\beta} \times C$. Likewise, $f_\beta^{-M_0}(\mathcal{B}_{0,\beta})$ is basically a thin horizontal strip. Iterating $f_\beta^{M_0-1}(\mathcal{A}_{0,\beta})$ once more deforms the previously horizontal strip to a thin strip around a parabola with second derivative at least s because of (A9). This yields an upper bound for the size of $\mathcal{I}_{1,\beta}$; see Figure 2(a).

The smallness of $\mathcal{I}_{n,\beta}$ follows in a similar fashion, but we have to show that even though the iterates of $\mathcal{A}_{n,\beta}$ enter the expanding region for some iterates, the overall effect of the iteration under f is still a contraction.

In order to formalise this intuitive idea, we define the functions

$$\phi_{n,\beta}^\pm(\theta) := f_{\beta,\theta-M_n\omega}^{M_n}(c^\pm) \quad \text{and} \quad \psi_{n,\beta}^\pm(\theta) := f_{\beta,\theta+M_n\omega}^{-M_n}(e^\pm)$$

for $\theta \in \mathcal{I}_{n,\beta} + \omega$, $n \in \mathbb{N}_0$. Note that

$$\begin{aligned} f_\beta^{M_n}(\mathcal{A}_{n,\beta}) &= \{(\theta, x) \in (\mathcal{I}_{n,\beta} + \omega) \times X : x \in [\phi_{n,\beta}^-(\theta), \phi_{n,\beta}^+(\theta)]\}, \\ f_\beta^{-M_n}(\mathcal{B}_{n,\beta}) &= \{(\theta, x) \in (\mathcal{I}_{n,\beta} + \omega) \times X : x \in [\psi_{n,\beta}^-(\theta), \psi_{n,\beta}^+(\theta)]\} \end{aligned}$$

(cf. Figure 2). We introduce a shorthand notation for the following inductive assumptions.

$$\left. \begin{aligned} \mathcal{B}(n) \text{ is a non-empty and closed interval,} \\ \mathcal{I}_{n,\beta} \text{ is closed and convex for } \beta \in \mathcal{B}(n), \\ \phi_{n,\beta}^-(\theta) > \psi_{n,\beta}^+(\theta) \text{ for each } \theta \in \partial \mathcal{I}_{n,\beta} + \omega \text{ and } \beta \in \mathcal{B}(n), \\ \exists \beta_-(n+1) \in \mathcal{B}(n) \text{ and } \exists! \theta_-^n \in \mathcal{I}_{n,\beta_-(n+1)} + \omega : \phi_{n,\beta_-(n+1)}^-(\theta_-^n) = \psi_{n,\beta_-(n+1)}^+(\theta_-^n), \\ \exists \beta_+(n+1) \in \mathcal{B}(n) \text{ and } \exists! \theta_+^n \in \mathcal{I}_{n,\beta_+(n+1)} + \omega : \phi_{n,\beta_+(n+1)}^+(\theta_+^n) = \psi_{n,\beta_+(n+1)}^-(\theta_+^n). \end{aligned} \right\} (\mathcal{I})_n \quad \begin{aligned} (4.3) \\ (4.4) \\ (4.5) \\ (4.6) \\ (4.7) \end{aligned}$$

Moreover, set:

- $H_n^\phi := \sup_{\theta \in \mathcal{I}_n, \beta \in \mathcal{B}(n)} |\phi_{n,\beta}^+(\theta) - \phi_{n,\beta}^-(\theta)|;$
- $H_n^\psi := \sup_{\theta \in \mathcal{I}_n, \beta \in \mathcal{B}(n)} |\psi_{n,\beta}^+(\theta) - \psi_{n,\beta}^-(\theta)|;$
- $v_n^\tau := \inf_{\substack{\theta \in \mathcal{I}_n, \beta \in \mathcal{B}(n) \\ \vartheta \in \mathbb{S}^{d-1}}} \partial_\vartheta^2 \phi_{n,\beta}^\tau(\theta) - \partial_\vartheta^2 \psi_{n,\beta}^{-\tau}(\theta) \quad (\tau \in \{-, +\}).$

LEMMA 4.13. Assume $(\mathcal{I})_n$ holds for some $n \in \mathbb{N}_0$. Then $\mathcal{B}(n+1)$ is non-empty. Further, suppose f_β satisfies (A3), (A4), (A6)–(A8), (A10) and $(\mathcal{F}1)_n$, $(\mathcal{F}2)_{n+1}$ for $\beta \in \mathcal{B}(n+1)$. If $v_n^\pm, v_{n+1}^\pm > 0$, then:

- $(\mathcal{I})_{n+1}$ holds;
- $|\mathcal{I}_{n+1,\beta}| \leq \sqrt{8} \sqrt{(H_n^\phi + H_n^\psi)/v_n^-}$ for $\beta \in \mathcal{B}(n+1)$.

Proof. Note that $\emptyset \neq \mathcal{B}(n+1) = [\beta_-(n+1), \beta_+(n+1)]$ by (4.6), (4.7) as well as Proposition 4.12 and (A6).

$\mathcal{I}_{n+1,\beta}$ is a sublevel set of $\phi_{n,\beta}^- - \psi_{n,\beta}^+$ and hence it is closed. Given two points $\theta_1, \theta_2 \in \mathcal{I}_{n+1,\beta}$, denote by $[\theta_1, \theta_2] \subseteq \mathcal{I}_{n,\beta}$ the line joining the two points. As $\partial_\vartheta^2 \phi_{n,\beta}^-(\theta) - \partial_\vartheta^2 \psi_{n,\beta}^+(\theta) \geq v_n^- > 0$ (with ϑ the unit vector in direction of $\theta_2 - \theta_1$), we have $\phi_{n,\beta}^- - \psi_{n,\beta}^+ \leq 0$ on $[\theta_1, \theta_2]$ and thus convexity of $\mathcal{I}_{n+1,\beta}$.

By applying Corollary 4.7, we see that $[\phi_{n+1,\beta}^-(\theta), \phi_{n+1,\beta}^+(\theta)] \subseteq (\phi_{n,\beta}^-(\theta), \phi_{n,\beta}^+(\theta))$ and $[\psi_{n+1,\beta}^-(\theta), \psi_{n+1,\beta}^+(\theta)] \subseteq (\psi_{n,\beta}^-(\theta), \psi_{n,\beta}^+(\theta))$ for all $\theta \in \mathcal{I}_{n+1,\beta} + \omega, \beta \in \mathcal{B}(n)$. This ensures $\phi_{n+1,\beta}^- > \psi_{n+1,\beta}^+$ on $\partial \mathcal{I}_{n+1,\beta}$ and guarantees that

$$\begin{aligned} \beta_+(n+2) &:= \min\{\beta \in \mathcal{B}(n+1) \mid \exists \theta \in \mathcal{I}_{n+1,\beta} + \omega : \phi_{n+1,\beta}^+(\theta) \leq \psi_{n+1,\beta}^-(\theta)\} \\ &= \min\{\beta \in \mathcal{B}(n+1) \mid \exists \theta \in \mathcal{I}_{n+1,\beta} + \omega : \phi_{n+1,\beta}^+(\theta) = \psi_{n+1,\beta}^-(\theta)\} \end{aligned}$$

as well as

$$\begin{aligned} \beta_-(n+2) &:= \max\{\beta \in \mathcal{B}(n+1) \mid \beta < \beta_+(n+2), \forall \theta \in \mathcal{I}_{n+1,\beta} + \omega : \phi_{n+1,\beta}^-(\theta) \geq \psi_{n+1,\beta}^+(\theta)\} \end{aligned}$$

are well-defined. Using $v_{n+1}^\pm > 0$, we get the uniqueness of the tangent points of $\phi_{n+1,\beta_-(n+1)}^-$ and $\psi_{n+1,\beta_-(n+1)}^+$ as well as of $\phi_{n+1,\beta_+(n+1)}^+$ and $\psi_{n+1,\beta_+(n+1)}^-$ and conclude $(\mathcal{I})_{n+1}$.

Note that $\phi_{n,\beta}^- - \psi_{n,\beta}^+ \geq -(H_n^\phi + H_n^\psi)$ and furthermore, $\partial_\theta^2(\phi_{n,\beta}^- - \psi_{n,\beta}^+) \geq v_n^-$. Suppose $\phi_{n,\beta}^- - \psi_{n,\beta}^+$ assumes its minimum at $\theta_0 \in \mathcal{I}_{n,\beta} + \omega$ and define $h : \theta \mapsto (v_n^-/2)(\theta - \theta_0)^2 - (H_n^\phi + H_n^\psi)$. As $\phi_{n,\beta}^-(\theta) - \psi_{n,\beta}^+(\theta) \leq 0$ if and only if $\theta \in \mathcal{I}_{n,\beta} + \omega$,

we necessarily have that $h(\theta) \leq 0$ for all $\theta \in \mathcal{I}_{n,\beta} + \omega$ such that an upper bound for the size of $\mathcal{I}_{n+1,\beta}$ is given by the distance of the zeros of h (cf. Figure 2(d)). $\square$

Remark 4.14. By means of (A5), we defined $\mathcal{B}(0) \subseteq [0, 1]$ to be a closed interval. Further, we set $\mathcal{I}_{0,\beta} \subseteq \mathbb{T}^d$ to be closed and convex for each $\beta \in \mathcal{B}(0)$. Moreover, with (A1) we have (4.5) and similarly as in the proof of Lemma 4.13 we can define $\beta_-(1)$ and $\beta_+(1)$ such that (4.6) and (4.7) are verified, respectively, assuming that (A9) holds. In other words: $(\mathcal{I})_0$ is true.

We next provide estimates for the quantities used in Lemma 4.13.

LEMMA 4.15. (Cf. [15, Lemma 3.13]) *Let $n \in \mathbb{N}_0$, $\beta \in \tilde{\mathcal{B}}(n)$. Suppose f_β verifies $(\mathcal{I})_n$ as well as (A1)–(A4) and (A8). If $(\mathcal{F})_{n-1}$ and $(\mathcal{F}2)_n$ hold, then $H_{n,\beta}^\phi \leq (\alpha_c^{b_n} \alpha_u^{1-b_n})^{M_n} \cdot |C|$ and $H_{n,\beta}^\psi \leq (\alpha_e^{b_n} \alpha_l^{1-b_n})^{-M_n} \cdot |E|$.*

Proof. Apply Corollary 4.10. $\square$

The following statement is, from a technical point of view, the core part of this work. It provides us with a positive lower bound for $v_n^\pm$ and thereby ensures that we can apply Lemma 4.13. The idea is to show that the second derivative of $\phi_n^\pm(\theta) - \psi_n^\mp(\theta) = f_{\theta-M_n\omega}^{M_n}(c^\pm) - f_{\theta+M_n\omega}^{-M_n}(c^\mp)$ in direction ϑ only differs from $(\partial_\vartheta^2 f_{\theta-\omega})(f_{\theta-M_n\omega}^{M_n-1}(c^\pm))$ by a remainder term, whose supremum goes to zero exponentially fast with increasing α . Since (A9) † provides us with a lower bound s for the second derivative of f with respect to the base coordinates in every direction, this proves the claim.

LEMMA 4.16. *Let $n \in \mathbb{N}_0$, $\beta \in \tilde{\mathcal{B}}(n)$. Suppose f_β satisfies (A1)–(A4), (A8), (A9), (A11)–(A16) and $(\mathcal{F})_{n-1}$ as well as $(\mathcal{F}2)_n$. Let there be $p \geq \sqrt{2}$ and $\alpha > 1$ such that*

$$\alpha_c^{-1} = \alpha_e = \alpha^{2/p}, \quad \alpha_l^{-1} = \alpha_u = \alpha^p$$

and assume $b_n > 5p^2/(2 + 5p^2)$. Then

$$v_n^\pm \geq s - S^2 c \cdot \alpha^{-((2b_n/p) - 5(1-b_n)p)},$$

where $c = c(\alpha, b_n) > 0$ can be chosen to be monotonously decreasing in α and b_n .

Proof. For reasons of readability, we omit the index β in the following. Let us consider $(\partial^2/\partial\vartheta^2)\phi_n^\pm(\theta)$ ($\theta \in \mathcal{I}_n + \omega$ and $\vartheta \in \mathbb{S}^{d-1}$). Set $\theta_0 := \theta - M_n\omega$. Then

$$\begin{aligned} \frac{\partial}{\partial\vartheta}\phi_n^\pm(\theta) &= \partial_\vartheta f_{\theta_0}^{M_n}(c^\pm) = (\partial_\vartheta f_{\theta_{M_n-1}})(x_{M_n-1}) + (\partial_x f_{\theta_{M_n-1}})(x_{M_n-1}) \cdot \partial_\vartheta f_{\theta_0}^{M_n-1}(c^\pm) \\ &= \dots = (\partial_\vartheta f_{\theta_{M_n-1}})(x_{M_n-1}) + \sum_{k=0}^{M_n-2} (\partial_\vartheta f_{\theta_k})(x_k) \cdot (\partial_x f_{\theta_{k+1}}^{M_n-k-1})(x_{k+1}), \end{aligned} \quad (4.8)$$

where we used

$$(\partial_x f_{\theta_{k+1}}^{M_n-k-1})(x_{k+1}) = \prod_{j=k+1}^{M_n-1} (\partial_x f_{\theta_j})(x_j) \quad (k = -1, 0, \dots, M_n-1). \quad (4.9)$$

† Note that indeed $f_{\theta-M_n\omega}^{M_n-1}(c^\pm) \in C$ because of Corollary 4.6.

Differentiating once more gives

$$\begin{aligned} \frac{\partial^2}{\partial \vartheta^2} \phi_n^\pm(\theta) &= (\partial_\vartheta^2 f_{\theta_{M_n-1}})(x_{M_n-1}) + (\partial_x \partial_\vartheta f_{\theta_{M_n-1}})(x_{M_n-1}) \cdot \partial_\vartheta f_{\theta_0}^{M_n-1}(c^\pm) \\ &\quad + \sum_{k=0}^{M_n-2} (\partial_\vartheta f_{\theta_k})(x_k) \cdot \partial_\vartheta (\partial_x f_{\theta_{k+1}}^{M_n-k-1})(x_{k+1}) \\ &\quad + [\partial_\vartheta (\partial_\vartheta f_{\theta_k})(x_k)] \cdot (\partial_x f_{\theta_{k+1}}^{M_n-k-1})(x_{k+1}). \end{aligned}$$

Further,

$$\begin{aligned} \partial_\vartheta (\partial_x f_{\theta_{k+1}}^{M_n-k-1})(x_{k+1}) &= \partial_\vartheta \prod_{j=k+1}^{M_n-1} (\partial_x f_{\theta_j})(x_j) = \sum_{l=k+1}^{M_n-1} \partial_\vartheta (\partial_x f_{\theta_l})(x_l) \prod_{\substack{j=k+1 \\ j \neq l}}^{M_n-1} (\partial_x f_{\theta_j})(x_j) \\ &= \sum_{l=k+1}^{M_n-1} [(\partial_\vartheta \partial_x f_{\theta_l})(x_l) + (\partial_x^2 f_{\theta_l})(x_l) \cdot \partial_\vartheta f_{\theta_0}^l(c^\pm)] \prod_{\substack{j=k+1 \\ j \neq l}}^{M_n-1} (\partial_x f_{\theta_j})(x_j) \end{aligned}$$

and

$$\partial_\vartheta (\partial_\vartheta f_{\theta_k})(x_k) = (\partial_\vartheta^2 f_{\theta_k})(x_k) + (\partial_x \partial_\vartheta f_{\theta_k})(x_k) \cdot \partial_\vartheta f_{\theta_0}^k(c^\pm).$$

Altogether, we have

$$\begin{aligned} \frac{\partial^2}{\partial \vartheta^2} \phi_n^\pm(\theta) &= (\partial_\vartheta^2 f_{\theta_{M_n-1}})(x_{M_n-1}) + (\partial_x \partial_\vartheta f_{\theta_{M_n-1}})(x_{M_n-1}) \cdot \partial_\vartheta f_{\theta_0}^{M_n-1}(c^\pm) \\ &\quad + \sum_{k=0}^{M_n-2} (\partial_\vartheta f_{\theta_k})(x_k) \left(\sum_{l=k+1}^{M_n-1} [(\partial_\vartheta \partial_x f_{\theta_l})(x_l) + (\partial_x^2 f_{\theta_l})(x_l) \cdot \partial_\vartheta f_{\theta_0}^l(c^\pm)] \prod_{\substack{j=k+1 \\ j \neq l}}^{M_n-1} (\partial_x f_{\theta_j})(x_j) \right) \\ &\quad + [(\partial_\vartheta^2 f_{\theta_k})(x_k) + (\partial_x \partial_\vartheta f_{\theta_k})(x_k) \cdot \partial_\vartheta f_{\theta_0}^k(c^\pm)] (\partial_x f_{\theta_{k+1}}^{M_n-k-1})(x_{k+1}). \end{aligned} \quad (4.10)$$

It is our goal to show that the long times spent in the contracting region keep the derivatives small, such that $(\partial_\vartheta^2 f_{\theta_{M_n-1}})(x_{M_n-1})$ becomes the leading term. The part which is the hardest to control is

$$\sum_{k=0}^{M_n-2} (\partial_\vartheta f_{\theta_k})(x_k) \sum_{l=k+1}^{M_n-1} (\partial_x^2 f_{\theta_l})(x_l) \cdot \partial_\vartheta f_{\theta_0}^l(c^\pm) \prod_{\substack{j=k+1 \\ j \neq l}}^{M_n-1} (\partial_x f_{\theta_j})(x_j),$$

with $\partial_\vartheta f_{\theta_0}^l(c^\pm) = \sum_{m=0}^{l-1} (\partial_\vartheta f_{\theta_m})(x_m) \cdot (\partial_x f_{\theta_{m+1}}^{l-m-1})(x_{m+1})$ as in equation (4.8). Using (A11), we see that it is bounded from above by

$$S^2 \sum_{k=0}^{M_n-2} \sum_{l=k+1}^{M_n-1} \sum_{m=0}^{l-1} |(\partial_x^2 f_{\theta_l})(x_l)| (\partial_x f_{\theta_{m+1}}^{l-m-1})(x_{m+1}) \prod_{\substack{j=k+1 \\ j \neq l}}^{M_n-1} (\partial_x f_{\theta_j})(x_j). \quad (4.11)$$

If $m \leq k$, then

$$\begin{aligned}
 & |(\partial_x^2 f_{\theta_l})(x_l)| (\partial_x f_{\theta_{m+1}}^{l-m-1})(x_{m+1}) \prod_{\substack{j=k+1 \\ j \neq l}}^{M_n-1} (\partial_x f_{\theta_j})(x_j) \\
 &= |(\partial_x^2 f_{\theta_l})(x_l)| \prod_{\substack{j=m+1 \\ j \neq l}}^{M_n-1} (\partial_x f_{\theta_j})(x_j) \cdot \prod_{j=k+1}^{l-1} (\partial_x f_{\theta_j})(x_j) \\
 &\leq |(\partial_x^2 f_{\theta_l})(x_l)| \prod_{\substack{j=m+1 \\ j \neq l \\ x_j \in C}}^{M_n-1} \alpha_c \cdot \prod_{\substack{j=m+1 \\ j \neq l \\ x_j \notin C}}^{M_n-1} \alpha_u \prod_{\substack{j=k+1 \\ x_j \notin C}}^{l-1} \alpha_u \\
 &\leq |(\partial_x^2 f_{\theta_l})(x_l)| \prod_{\substack{j=m+1 \\ j \neq l \\ x_j \in C}}^{M_n-1} \alpha_c \cdot \prod_{\substack{j=m+1 \\ j \neq l \\ x_j \notin C}}^{M_n-1} \alpha_u^2 \leq \alpha_c^{b_n(M_n-m-1)} \alpha_u^{2(1-b_n)(M_n-m-1)} \\
 &= \alpha_1^{-(M_n-m-1)},
 \end{aligned}$$

where we used Lemma 4.8[†] and (A14) in the last estimate and where we set $\alpha_1 := \alpha_c^{-b_n} \alpha_u^{-2(1-b_n)} = \alpha^{-2(p(1-b_n)-(b_n/p))}$. For $m > k$ we get an analogous result with m replaced by k . Hence, (4.11) is bounded by

$$\begin{aligned}
 & S^2 \sum_{k=0}^{M_n-2} \sum_{l=k+1}^{M_n-1} \left(\sum_{m=0}^k \alpha_1^{-(M_n-1-m)} + \sum_{m=k+1}^{l-1} \alpha_1^{-(M_n-1-k)} \right) \\
 &\leq S^2 \sum_{k=0}^{M_n-2} \sum_{l=k+1}^{M_n-1} \left(\alpha_1^{-(M_n-1-k)} \sum_{m=0}^k \alpha_1^{-m} + \alpha_1^{-(M_n-1-k)} \sum_{m=k+1}^{l-1} 1 \right) \\
 &\leq S^2 \sum_{k=0}^{M_n-2} \left(\alpha_1^{-(M_n-1-k)} \frac{1}{1-\alpha_1^{-1}} (M_n-k-1) + \alpha_1^{-(M_n-1-k)} \sum_{l=k+1}^{M_n-1} (l-k-1) \right) \\
 &\leq S^2 \frac{2}{1-\alpha_1^{-1}} \sum_{k=0}^{M_n-2} \alpha_1^{-(M_n-1-k)} (M_n-k-1)^2 \leq S^2 \frac{2\alpha_1}{\alpha_1-1} \sum_{l=1}^{M_n-1} l^2 \alpha_1^{-l} \\
 &\leq S^2 \tilde{c}(\alpha_1) \cdot \alpha_1^{-1},
 \end{aligned}$$

where $\tilde{c}(\alpha) := (2\alpha/(\alpha-1)) \sum_{l=1}^{\infty} l^2 \alpha^{-l+1}$ for each $\alpha > 1$ [‡]. Note that $\tilde{c}$ is monotonously decreasing in α . The other addends of (4.10) can be treated in a similar fashion, which eventually gives

$$\frac{\partial^2}{\partial \vartheta^2} \phi_n^{\pm}(\theta) \geq (\partial_{\vartheta}^2 f_{\theta_{M_n-1}})(x_{M_n-1}) - 5S^2 \tilde{c}(\alpha_1) \cdot \alpha_1^{-1} \stackrel{(A9)}{\geq} s - 5S^2 \tilde{c}(\alpha_1) \cdot \alpha_1^{-1}.$$

Now, let us consider $(\partial^2/\partial \vartheta^2) \psi_n^{\pm}(\theta) = f_{\theta+M_n\omega}^{-M_n}(e^{\pm})$ for $\vartheta \in \mathbb{S}^{d-1}$. We proceed similarly as before but this time considering the map f^{-1} instead of f , that is $\theta_k = \theta_0 - k\omega$

[†] Observe that the assumptions of Lemma 4.8 are verified due to Remark 4.5.

[‡] Notice that $\alpha_1 > 1$, since $b_n > 5p^2/(2+5p^2) > p^2/(p^2+1)$.

(with $\theta_0 = \theta + M_n \omega$) and $x_k = f_{\theta_0}^{-k}(e^\pm)$:

$$\begin{aligned}\partial_x f_\theta^{-1}(x) &= \frac{1}{(\partial_x f_{\theta-\omega})(f_\theta^{-1}(x))} \quad (\Rightarrow 0 < \partial_x f_\theta^{-1}(x) < \alpha_e^{-1} \text{ (} x \in E, \theta \notin \mathcal{I}_0 + \omega \text{)}), \\ \partial_\vartheta f_\theta^{-1}(x) &= -\frac{(\partial_\vartheta f_{\theta-\omega})(f_\theta^{-1}(x))}{(\partial_x f_{\theta-\omega})(f_\theta^{-1}(x))} = -(\partial_\vartheta f_{\theta-\omega})(f_\theta^{-1}(x)) \cdot \partial_x f_\theta^{-1}(x),\end{aligned}\tag{4.12}$$

for $(\theta, x) \in f(\mathbb{T}^d \times X) \setminus \{(\theta, x) : Df^{-1} \text{ is singular}\} \supseteq f(\mathbb{T}^d \times [e^-, c^+])$. Hence,

$$\partial_x^2 f_\theta^{-1}(x) = -\frac{(\partial_x^2 f_{\theta-\omega})(f_\theta^{-1}(x)) \cdot \partial_x f_\theta^{-1}(x)}{[(\partial_x f_{\theta-\omega})(f_\theta^{-1}(x))]^2} = -(\partial_x^2 f_{\theta-\omega})(f_\theta^{-1}(x)) \cdot (\partial_x f_\theta^{-1}(x))^3\tag{4.13}$$

such that $|\partial_x^2 f_\theta^{-1}(x)| \leq \alpha_u^2 \alpha_l^{-3}$ for $x \in f_{\theta-\omega}([e^-, c^+])$,

$$\begin{aligned}\partial_\vartheta \partial_x f_\theta^{-1}(x) &= -\frac{(\partial_\vartheta \partial_x f_{\theta-\omega})(f_\theta^{-1}(x)) + (\partial_x^2 f_{\theta-\omega})(f_\theta^{-1}(x)) \cdot \partial_\vartheta f_\theta^{-1}(x)}{[(\partial_x f_{\theta-\omega})(f_\theta^{-1}(x))]^2} \\ &= -(\partial_\vartheta \partial_x f_{\theta-\omega})(f_\theta^{-1}(x)) \cdot (\partial_x f_\theta^{-1}(x))^2 - (\partial_\vartheta f_{\theta-\omega})(f_\theta^{-1}(x)) \cdot \partial_x^2 f_\theta^{-1}(x)\end{aligned}\tag{4.14}$$

and thus $|\partial_\vartheta \partial_x f_\theta^{-1}(x)| \leq 2S\alpha_u^2 \alpha_l^{-3}$ for $x \in f_{\theta-\omega}([e^-, c^+])$. Finally,

$$\begin{aligned}\partial_\vartheta^2 f_\theta^{-1}(x) &= -(\partial_\vartheta^2 f_{\theta-\omega})(f_\theta^{-1}(x)) \cdot \partial_x f_\theta^{-1}(x) \\ &\quad - (\partial_x \partial_\vartheta f_{\theta-\omega})(f_\theta^{-1}(x)) \cdot \partial_\vartheta f_\theta^{-1}(x) \partial_x f_\theta^{-1}(x) \\ &\quad - (\partial_\vartheta f_{\theta-\omega})(f_\theta^{-1}(x)) \cdot \partial_\vartheta \partial_x f_\theta^{-1}(x) \\ &= -(\partial_\vartheta^2 f_{\theta-\omega})(f_\theta^{-1}(x)) \cdot \partial_x f_\theta^{-1}(x) - 2(\partial_\vartheta f_{\theta-\omega})(f_\theta^{-1}(x)) \cdot \partial_\vartheta \partial_x f_\theta^{-1}(x) \\ &\quad - ((\partial_\vartheta f_{\theta-\omega})(f_\theta^{-1}(x)))^2 \cdot \partial_x^2 f_\theta^{-1}(x).\end{aligned}\tag{4.15}$$

As in the forward case, we get

$$\begin{aligned}\frac{\partial^2}{\partial \vartheta^2} \psi_n^\pm(\theta) &= \sum_{k=0}^{M_n-1} (\partial_\vartheta f_{\theta_k}^{-1})(x_k) \\ &\quad \times \left(\sum_{l=k+1}^{M_n-1} [(\partial_\vartheta \partial_x f_{\theta_l}^{-1})(x_l) + (\partial_x^2 f_{\theta_l}^{-1})(x_l) \cdot \partial_\vartheta f_{\theta_0}^{-l}(e^\pm)] \prod_{\substack{j=k+1 \\ j \neq l}}^{M_n-1} (\partial_x f_{\theta_j}^{-1})(x_j) \right) \\ &\quad + [(\partial_\vartheta^2 f_{\theta_k}^{-1})(x_k) + (\partial_\vartheta \partial_x f_{\theta_k}^{-1})(x_k) \cdot \partial_\vartheta f_{\theta_0}^{-k}(e^\pm)] (\partial_x f_{\theta_{k+1}}^{-(M_n-k-1)})(x_{k+1}).\end{aligned}\tag{4.16}$$

Similarly as before, we want to show that the long times spent in the expanding region keep all the derivatives small (as we consider iterates of the inverse map). Since $\partial_\vartheta f_{\theta_0}^{-l}(e^\pm) = \sum_{m=0}^{l-1} (\partial_\vartheta f_{\theta_m}^{-1})(x_m) \cdot (\partial_x f_{\theta_{m+1}}^{-(l-m-1)})(x_{m+1})$, the term which is the hardest to control in (4.16) is

$$\begin{aligned}
& \sum_{k=0}^{M_n-1} (\partial_{\vartheta} f_{\theta_k}^{-1})(x_k) \sum_{l=k+1}^{M_n-1} (\partial_x^2 f_{\theta_l}^{-1})(x_l) \partial_{\vartheta} f_{\theta_0}^{-l}(e^{\pm}) \prod_{\substack{j=k+1 \\ j \neq l}}^{M_n-1} (\partial_x f_{\theta_j}^{-1})(x_j) \\
&= - \sum_{k=0}^{M_n-2} (\partial_{\vartheta} f_{\theta_{k+1}})(x_{k+1}) \partial_x f_{\theta_k}^{-1}(x_k) \sum_{l=k+1}^{M_n-1} (\partial_x^2 f_{\theta_l}^{-1})(x_l) \\
&\quad \times \sum_{m=0}^{l-1} (\partial_{\vartheta} f_{\theta_{m+1}})(x_{m+1}) \partial_x f_{\theta_m}^{-1}(x) (\partial_x f_{\theta_{m+1}}^{-(l-m-1)})(x_{m+1}) \\
&\quad \times \prod_{\substack{j=k+1 \\ j \neq l}}^{M_n-1} (\partial_x f_{\theta_j}^{-1})(x_j).
\end{aligned}$$

An upper bound for this expression reads

$$S^2 \sum_{k=0}^{M_n-2} \sum_{l=k+1}^{M_n-1} \sum_{m=0}^{l-1} |(\partial_x^2 f_{\theta_l}^{-1})(x_l)| \cdot \prod_{n=m}^{l-1} (\partial_x f_{\theta_n}^{-1})(x_n) \prod_{\substack{j=k \\ j \neq l}}^{M_n-1} (\partial_x f_{\theta_j}^{-1})(x_j). \quad (4.17)$$

We deal similarly with (4.17) as we did with (4.11). Suppose $m > k$. Since $(\partial_x f_{\theta}^{-1}(x))^2 < \alpha_l^{-2} < \alpha_u^2 \alpha_l^{-3}$, we get

$$\begin{aligned}
& |(\partial_x^2 f_{\theta_l}^{-1})(x_l)| \prod_{n=m}^{l-1} (\partial_x f_{\theta_n}^{-1})(x_n) \prod_{\substack{j=k \\ j \neq l}}^{M_n-1} (\partial_x f_{\theta_j}^{-1})(x_j) \\
&\leq |(\partial_x^2 f_{\theta_l}^{-1})(x_l)| \prod_{\substack{j=k \wedge j \neq l \\ x_j \in E \wedge \theta_j \notin \mathcal{I}_0 + \omega}}^{M_n-1} (\partial_x f_{\theta_j}^{-1})(x_j) \prod_{\substack{j=k \wedge j \neq l \\ x_j \notin E \vee \theta_j \in \mathcal{I}_0 + \omega}}^{M_n-1} (\partial_x f_{\theta_n}^{-1})^2(x_n) \\
&\stackrel{(A15)}{\leq} \alpha_e^{-b_n(M_n-k)} (\alpha_u^2 \alpha_l^{-3})^{(1-b_n)(M_n-k)} \leq \alpha_2^{-(M_n-k)},
\end{aligned}$$

where $\alpha_2 := \alpha_e^{b_n} (\alpha_u^2 \alpha_l^{-3})^{-(1-b_n)} = \alpha^{(2b_n/p)-5(1-b_n)p}$. For $m \leq k$ we get an analogous result. Hence, (4.17) is bounded by

$$\begin{aligned}
& S^2 \sum_{k=0}^{M_n-2} \sum_{l=k+1}^{M_n-1} \left(\alpha_2^{-(M_n-k)} \sum_{m=0}^k \alpha_2^{-m} + \alpha_2^{-(M_n-k)} \sum_{m=k+1}^{l-1} \right) \\
&\leq S^2 \frac{2\alpha_2}{\alpha_2 - 1} \sum_{k=0}^{M_n-2} \alpha_2^{-(M_n-k)} (M_n - k)^2 \leq S^2 \tilde{c}(\alpha_2) \cdot \alpha_2^{-2},
\end{aligned}$$

with $\tilde{c}$ as in the forward case[†]. Nevertheless, notice that

$$\begin{aligned}
& \left| \sum_{k=0}^{M_n-1} (\partial_{\vartheta}^2 f_{\theta_k}^{-1})(x_k) (\partial_x f_{\theta_{k+1}}^{-(M_n-k-1)})(x_{k+1}) \right| \leq \sum_{k=0}^{M_n-1} S^2 (\partial_x f_{\theta_k}^{-(M_n-k)})(x_k) \\
&\quad + 2S |\partial_{\vartheta} \partial_x f_{\theta_k}^{-1}(x_k)| (\partial_x f_{\theta_{k+1}}^{-(M_n-k-1)})(x_{k+1}) + S^2 \partial_x^2 f_{\theta_k}^{-1}(x_k) (\partial_x f_{\theta_{k+1}}^{-(M_n-k-1)})(x_{k+1}) \\
&\leq 3S^2 \tilde{c}(\alpha_2) \alpha_2^{-1},
\end{aligned}$$

[†] $\alpha_2 > 1$, since $b > 1 - (2/(2 + 5p^2))$.

where we used (4.15) in the first step. Altogether, we eventually get

$$\frac{\partial^2}{\partial \vartheta^2} \psi^\pm(\theta) \leq 6S^2 \tilde{c}(\alpha_2) \cdot \alpha_2^{-1}.$$

Setting $c(\alpha, b_n) := 6\tilde{c}(\alpha^{(2b_n/p)-5(1-b_n)p}) + 5\tilde{c}(\alpha^{(2b_n/p)-2(1-b_n)p})$ yields the desired estimate. $\square$

4.3. Existence of a sink-source orbit. In §4.1, we proved the existence of a sink-source orbit for f_β provided there are strictly increasing sequences $(M_n)_{n \in \mathbb{N}_0}$, $(K_n)_{n \in \mathbb{N}_0} \in \mathbb{N}^{\mathbb{N}_0}$ such that the inductively defined critical regions $\mathcal{I}_{n,\beta}$ are non-empty and satisfy $(\mathcal{F}1)_n$, $(\mathcal{F}2)_n$. By means of the geometric considerations of the last section, we are now able to show that for some β such sequences $(M_n)_{n \in \mathbb{N}_0}$, $(K_n)_{n \in \mathbb{N}_0}$ actually do exist. This finishes the proof of Theorem 3.1.

As a matter of fact, we prove that for some β , the critical regions satisfy a slightly stronger version of $(\mathcal{F}1)_n$, that is, we will show

$$d\left(\mathcal{I}_{j,\beta}, \bigcup_{k=1}^{2K_j M_j} \mathcal{I}_{j,\beta} + k\omega\right) > \varepsilon_j \geq |\mathcal{I}_{j,\beta}| \quad (\mathcal{F}1)'_n$$

for $j = 0, \dots, n$ and $n \in \mathbb{N}_0$ where $\varepsilon_0 := \frac{1}{2}\mathcal{C}(2K_0 M_0)^{-\eta}$ and $\varepsilon_j := \tilde{\mathcal{C}}\tilde{\alpha}^{-M_{j-1}}$ for all $j \in \mathbb{N}$ and some positive constants $\mathcal{C}$ and $\tilde{\alpha} > 1$.

LEMMA 4.17. (Cf. [15, Lemma 3.16]) *Assume $(\mathcal{I})_{n-1}$ for $n \in \mathbb{N}$. Suppose f_β verifies $(A6)$, $(A10)$ and we are given K_l, M_l ($l = 0, \dots, n-1$) such that $(\mathcal{F}1)'_{n-1}$, $(\mathcal{F}2)_{n-1}$ hold for $\beta \in \mathcal{B}(n)$. If $\sum_{j=0}^{n-1} (1/K_j) \leq \frac{1}{6}$, then there exists $M_n \in [K_{n-1}M_{n-1}, 2K_{n-1}M_{n-1} - 2]$ such that $(\mathcal{F}2)_n$ holds for f_β ($\beta \in \mathcal{B}(n)$).*

Proof. By Proposition 4.12, monotonicity of $f_{(\cdot)}(\theta, x)$ yields that f_β ($\beta \in \mathcal{B}(n)$) verifies $(\mathcal{F}2)_n$ if just $f_{\beta_+(n)}$ does. Therefore, we only consider $\beta = \beta_+(n)$ and suppress the index β in the following.

Let $j = 0, \dots, n-1$. Then,

$$\mathcal{I}_n - (M_n - 1)\omega \cap \bigcup_{l=-(M_j-1)}^{M_j+1} \mathcal{I}_j + l\omega \neq \emptyset$$

implies

$$\mathcal{I}_j - (M_n - 1)\omega \cap \mathcal{I}_j + l\omega \neq \emptyset,$$

for some $l \in \{-M_j + 1, -M_j + 2, \dots, M_j + 1\}$. By $(\mathcal{F}1)'_{n-1}$,

$$\begin{aligned} & \#\{q \in [K_{n-1}M_{n-1}, 2K_{n-1}M_{n-1} - 2] \cap \mathbb{N} \mid \mathcal{I}_j - (q-1)\omega \cap \mathcal{I}_j + l\omega \neq \emptyset\} \\ & \leq \frac{K_{n-1}M_{n-1} - 2}{2K_j M_j}. \end{aligned}$$

Hence,

$$\begin{aligned} & \#\left\{q \in [K_{n-1}M_{n-1}, 2K_{n-1}M_{n-1} - 2] \cap \mathbb{N} \mid \mathcal{I}_j - (q-1)\omega \cap \bigcup_{l=-(M_j-1)}^{M_j+1} \mathcal{I}_j + l\omega \neq \emptyset\right\} \\ & \leq (2M_j + 1) \frac{K_{n-1}M_{n-1} - 2}{2K_j M_j}. \end{aligned}$$

For the number of $q \in \{K_{n-1}M_{n-1}, K_{n-1}M_{n-1} + 1, \dots, 2K_{n-1}M_{n-1} - 2\}$ with $\mathcal{I}_j + (q+1)\omega \cap \bigcup_{l=-(M_j-1)}^{M_j+1} \mathcal{I}_j + l\omega \neq \emptyset$, we get the same upper bound. Therefore,

$$\begin{aligned} & \# \left\{ q \in [K_{n-1}M_{n-1}, 2K_{n-1}M_{n-1} - 2] \cap \mathbb{N} \mid (\mathcal{I}_n - (q-1)\omega \cup \mathcal{I}_n \right. \\ & \quad \left. + (q+1)\omega) \cap \bigcup_{j=0}^{n-1} \bigcup_{l=-(M_j-1)}^{M_j+1} \mathcal{I}_j + l\omega \neq \emptyset \right\} \\ & \leq 2(K_{n-1}M_{n-1} - 2) \sum_{j=0}^{n-1} \frac{2M_j + 1}{2K_j M_j} \leq 3(K_{n-1}M_{n-1} - 2) \sum_{j=0}^{n-1} \frac{1}{K_j}. \end{aligned}$$

Thus, if $\sum_{j=0}^{n-1} (1/K_j) \leq \frac{1}{6}$, there is $M_n \in [K_{n-1}M_{n-1}, 2K_{n-1}M_{n-1} - 2] \cap \mathbb{N}$ such that $(\mathcal{F}2)_n$ holds. $\square$

Given $\alpha > 1$ and $b_1 = 1 - 1/K_0$, set

$$v := s - c(\alpha, b_1^2) S^2 \alpha^{-(2b_1^2/p - 5(1-b_1^2)p)},$$

where $c(\alpha, b_1^2)$ is as in Lemma 4.16. Theorem 3.1 follows from the following statement.

THEOREM 4.18. *Suppose ω is Diophantine of type $(\mathcal{C}, \eta)$ and $(f_\beta)_{\beta \in [0,1]}$ satisfies (A1)–(A16). Let there be $p \geq \sqrt{2}$ and $\alpha > 1$ with*

$$\alpha_c^{-1} = \alpha_e = \alpha^{2/p}, \quad \alpha_l^{-1} = \alpha_u = \alpha^p.$$

Further, assume $2|\mathcal{I}_{0,\beta}| < \mathcal{C}(2K_0M_0)^{-\eta}$ for some $K_0, M_0 \in \mathbb{N}_{\geq 2}$ and assume $v > 0$. Then there exists $\alpha_0 = \alpha_0(v, K_0, M_0, p, |C|, |E|, \eta, \mathcal{C})$ such that if $\alpha > \alpha_0$, there is $\beta_c \in [0, 1]$ such that f_{β_c} has a sink-source orbit in $\mathbb{T}^d \times [e^-, c^+]$, and hence an SNA and an SNR.

Remark 4.19. We can choose α_0 to depend monotonously decreasing on v . Further, note that since we assume $v > 0$, we necessarily have $K_0 > 2 + 5p^2$.

Proof. $(\mathcal{F}1)'_n$ is necessarily satisfied if $2|\mathcal{I}_{n,\beta}| \leq 2\varepsilon_n < d(k\omega, 0)$ ($k = 1, \dots, 2K_nM_n - 2$). Note that since ω is Diophantine of type $(\mathcal{C}, \eta)$, we have

$$\mathcal{C}(2K_nM_n - 2)^{-\eta} < d(k\omega, 0) \tag{4.18}$$

for $k = 1, \dots, 2K_nM_n - 2$. Hence, $(\mathcal{F}1)'_0$ holds by the assumptions. Therefore, Lemma 4.17 together with Remark 4.14 yields the existence of $M_1 \in [K_0M_0, 2K_0M_0 - 2]$ such that $(\mathcal{F}2)_1$ holds for $\beta \in \mathcal{B}(1)$. Lemma 4.16 gives $v_0^\pm, v_1^\pm \geq v > 0$ such that Lemma 4.13 yields $(\mathcal{I})_1$. By means of Lemma 4.13 together with Lemma 4.15, we get

$$|\mathcal{I}_{1,\beta}| \leq \tilde{\mathcal{C}} \alpha_c^{M_0/2} \leq \tilde{\mathcal{C}} \tilde{\alpha}^{M_0}$$

where $\tilde{\mathcal{C}} := \sqrt{8(|C| + |E|)/v}$ and $\tilde{\alpha} := \alpha^{(-b/p + p(1-b)/2)}$.

Let us set $(K_n)_{n \in \mathbb{N}_0} := (K_0\kappa^n)_{n \in \mathbb{N}_0}$ for some $\kappa \in \mathbb{N}_{\geq 2}$ large enough to guarantee that $b > b_1^2$, and hence $v_n^\pm \geq v$ for all $n \in \mathbb{N}$. Then, since $M_1 \in [K_0M_0, 2K_0M_0 - 2]$, we have that for $n = 1$ (4.18) is bounded from below by

$$\mathcal{C}(2K_1M_1)^{-\eta} \geq \frac{\mathcal{C}}{(4\kappa K_0^2 M_0)^\eta}.$$

Therefore, if α is large enough, $(\mathcal{F}1)'_1$ is verified.

Let $n \in \mathbb{N}_{\geq 2}$. Suppose $(\mathcal{I})_{n-1}$ and $(\mathcal{F}1)'_{n-1}$ as well as $(\mathcal{F}2)_{n-1}$ (for $\beta \in \mathcal{B}(n)$) hold with $K_j = K_0 \kappa^j$ and $M_j \in [K_{j-1}M_{j-1}, 2K_{j-1}M_{j-1} - 2]$ for $j = 1, \dots, n-1$. As for $n = 1$, Lemma 4.17 yields $M_n \in [M_{n-1}K_{n-1}, 2M_{n-1}K_{n-1} - 2]$ such that $(\mathcal{F}2)_n$ holds. Now, $(\mathcal{I})_n$ follows similarly as in the case $n = 1$. By means of Lemma 4.13 and Lemma 4.15, we get

$$\begin{aligned} 2|\mathcal{I}_{n,\beta}| &\leq 2\tilde{\mathcal{C}}\alpha_c^{M_{n-1}b_{n-1}/2}\alpha_u^{M_{n-1}(1-b_{n-1})/2} \leq 2\tilde{\mathcal{C}}\alpha^{(-b/p+p(1-b)/2)M_{n-1}} \\ &= 2\tilde{\mathcal{C}}\tilde{\alpha}^{M_{n-1}} \leq \frac{\mathcal{C}}{(4\kappa^{2n-1}K_0^2M_{n-1})^\eta} \end{aligned}$$

and thereby $(\mathcal{F}1)'_n$ for $\beta \in \mathcal{B}(n)$, where the last inequality holds for all $n \in \mathbb{N}$ if α is large enough.

By induction, we thus see that there are sequences $(M_n)_{n \in \mathbb{N}_0}$ and $(K_n)_{n \in \mathbb{N}_0}$ such that $(\mathcal{I})_n$ is true for all $n \in \mathbb{N}_0$. Moreover, with these sequences we get $(\mathcal{F})_n$ (and actually $(\mathcal{F}1)'_n$) for each $n \in \mathbb{N}$ and $\beta \in \mathcal{B}(n)$. Applying Proposition 4.11 finishes the proof. $\square$

Acknowledgements. I would like to thank Tobias Jäger for pointing out this problem to me. Moreover, I am very grateful for the fruitful discussions with him and Alejandro Passeggi as well as Maik Gröger.

This work was supported by the Emmy-Noether-grant ‘Low-dimensional and Nonautonomous Dynamics’ (Ja 1721/2-1) of the German Research Council.

REFERENCES

- [1] V. Anagnostopoulou and T. Jäger. Nonautonomous saddle-node bifurcations: Random and deterministic forcing. *J. Differential Equations* **253**(2) (2012), 379–399.
- [2] L. Arnold. *Random Dynamical Systems*. Springer, Berlin, 1998.
- [3] K. Bjerklov. Dynamics of the quasi-periodic Schrödinger cocycle at the lowest energy in the spectrum. *Comm. Math. Phys.* **272** (2007), 397–442, doi:10.1007/s00220-007-0238-y.
- [4] K. Bjerklov and M. Saprykina. Universal asymptotics in hyperbolicity breakdown. *Nonlinearity* **21**(3) (2008), 557–586, QC 20100525.
- [5] M. Crucifix. Why could ice ages be unpredictable? *Clim. Past* **9** (2013), 1053–1098.
- [6] G. Fuhrmann, M. Gröger and T. Jäger. Non-smooth saddle-node bifurcations II: Dimensions of the SNA, in preparation, 2014.
- [7] P. Glendinning. The nonsmooth pitchfork bifurcation. *Discrete Contin. Dyn. Syst. Ser. B*. 2004, pp. 457–464.
- [8] C. Grebogi, E. Ott, S. Pelikan and J. A. Yorke. Strange attractors that are not chaotic. *Phys. D* **13** (1984), 261–268.
- [9] M. Gröger and T. Jäger. Dimensions of attractors in pinched skew products. *Comm. Math. Phys.* **320**(1) (2013), 101–119.
- [10] A. Haro and R. de la Llave. Manifolds on the verge of a hyperbolicity breakdown. *Chaos* **16**(1) (2006), 013120.
- [11] M. Herman. Une méthode pour minorer les exposants de Lyapounov et quelques exemples montrant le caractère local d’un théorème d’Arnold et de Moser sur le tore de dimension 2. *Comment. Math. Helv.* **58** (1983), 453–502, doi:10.1007/BF02564647.
- [12] T. Jäger. The creation of strange non-chaotic attractors in non-smooth saddle-node bifurcations. *Mem. Amer. Math. Soc.* **201** (2009), 1–106.
- [13] T. Jäger. On the structure of strange non-chaotic attractors in pinched skew products. *Ergod. Th. & Dynam. Sys.* **27** (2007), 493–510.
- [14] T. Jäger. Quasiperiodically forced interval maps with negative schwarzian derivative. *Nonlinearity* **16**(4) (2003), 1239–1255.
- [15] T. Jäger. Strange non-chaotic attractors in quasiperiodically forced circle maps. *Comm. Math. Phys.* **289** (2009), 253–289, doi:10.1007/s00220-009-0753-0.

- [16] G. Keller. A note on strange nonchaotic attractors. *Fund. Math.* **151** (1996), 139–148.
- [17] J. Kim and Y. Kim. Strange nonchaotic attractors in quasiperiodically driven Izhikevich neuron models. *J. Korean Phys. Soc.* **60**(4) (2012), 660–664.
- [18] V. Millionščikov. Proof of the existence of irregular systems of linear differential equations with quasi periodic coefficients. *Differ. Uravn.* **5**(11) (1969), 1979–1983.
- [19] T. Nguyen, T. Doan, T. Jäger and S. Siegmund. Saddle-node bifurcations in the quasiperiodically forced logistic map. *Internat. J. Bifur. Chaos* **21** (2011), 1427–1438.
- [20] C. Núñez and R. Obaya. A non-autonomous bifurcation theory for deterministic scalar differential equations. *Discrete Contin. Dyn. Syst. Ser. B* **9** (2008), 701–730.
- [21] H. Osinga, J. Wiersig, P. Glendinning and U. Feudel. Multistability and nonsmooth bifurcations in the quasiperiodically forced circle map. *Internat. J. Bifur. Chaos* **11** (2001), 3085–3105.
- [22] C. Pötsche. *Bifurcations in Nonautonomous Dynamical Systems: Results and Tools in Discrete Time*. Servizo de Publicacións da Universidade de Vigo, Vigo, 2011, pp. 163–212.
- [23] D. Sonechkin and N. Ivachtchenko. On the role of quasiperiodic forcing in the interannual and interdecadal climate variations. *CLIVAR Exchanges* **6** (2001), 5–6.
- [24] J. Stark. Transitive sets for quasi-periodically forced monotone maps. *Dyn. Syst.* **18**(4) (2003), 351–364.
- [25] A. J. Veraart, E. J. Faassen, V. Dakos, E. H. van Nes, M. Lürling and M. Scheffer. Recovery rates reflect distance to a tipping point in a living system. *Nature* **481** (2012), 357–359.
- [26] R. Vinograd. A problem suggested by N. R. Erugin. *Differ. Uravn.* **11**(4) (1975), 632–638.
- [27] L.-S. Young. Lyapunov exponents for some quasi-periodic cocycles. *Ergod. Th. & Dynam. Sys.* **17** (1997), 483–504.