Dieses Dokument ist eine Zweitveröffentlichung (Verlagsversion) /

This is a self-archiving document (published version):

I. Boehm, J. A. King, F. Bernardoni, D. Geisler, M. Seidel, F. Ritschel, T. Goschke, J.-D. Haynes, V. Roessner, S. Ehrlich

Subliminal and supraliminal processing of reward-related stimuli in anorexia nervosa

Erstveröffentlichung in / First published in:

Psychological Medicine. 2018, 48 (5), S. 790 – 800 [Zugriff am: 15.04.2020]. Cambridge University Press. ISSN 1469-8978.

DOI: <https://doi.org/10.1017/S0033291717002161>

Diese Version ist verfügbar / This version is available on:

<https://nbn-resolving.org/urn:nbn:de:bsz:14-qucosa2-707542>

„Dieser Beitrag ist mit Zustimmung des Rechteinhabers aufgrund einer (DFGgeförderten) Allianz- bzw. Nationallizenz frei zugänglich.“

This publication is openly accessible with the permission of the copyright owner. The permission is granted within a nationwide license, supported by the German Research Foundation (abbr. in German DFG).
www.nationallizenzen.de/

Subliminal and supraliminal processing of reward-related stimuli in anorexia nervosa

I. Boehm^{1,2}, J. A. King^{1,2}, F. Bernardoni^{1,2}, D. Geisler^{1,2}, M. Seidel^{1,2}, F. Ritschel^{1,2}, T. Goschke³, J.-D. Haynes⁴, V. Roessner⁵ and S. Ehrlich^{1,2*}

¹Division of Psychological and Social Medicine and Developmental Neurosciences, University Hospital Carl Gustav Carus, Technische Universität Dresden, Dresden, Germany

²Department of Child and Adolescent Psychiatry, Eating Disorders Research and Treatment Center, University Hospital Carl Gustav Carus, Technische Universität Dresden, Dresden, Germany

³Department of Psychology, Technische Universität Dresden, Dresden, Germany

⁴Bernstein Center for Computational Neuroscience, Charité Universitäts-Medizin, Berlin, Germany

⁵Department of Child and Adolescent Psychiatry, University Hospital Carl Gustav Carus, Technische Universität Dresden, Dresden, Germany

Background. Previous studies have highlighted the role of the brain reward and cognitive control systems in the etiology of anorexia nervosa (AN). In an attempt to disentangle the relative contribution of these systems to the disorder, we used functional magnetic resonance imaging (fMRI) to investigate hemodynamic responses to reward-related stimuli presented both subliminally and supraliminally in acutely underweight AN patients and age-matched healthy controls (HC).

Methods. fMRI data were collected from a total of 35 AN patients and 35 HC, while they passively viewed subliminally and supraliminally presented streams of food, positive social, and neutral stimuli. Activation patterns of the *group × stimulation condition × stimulus type* interaction were interrogated to investigate potential group differences in processing different stimulus types under the two stimulation conditions. Moreover, changes in functional connectivity were investigated using generalized psychophysiological interaction analysis.

Results. AN patients showed a generally increased response to supraliminally presented stimuli in the inferior frontal junction (IFJ), but no alterations within the reward system. Increased activation during supraliminal stimulation with food stimuli was observed in the AN group in visual regions including superior occipital gyrus and the fusiform gyrus/parahippocampal gyrus. No group difference was found with respect to the subliminal stimulation condition and functional connectivity.

Conclusion. Increased IFJ activation in AN during supraliminal stimulation may indicate hyperactive cognitive control, which resonates with clinical presentation of excessive self-control in AN patients. Increased activation to food stimuli in visual regions may be interpreted in light of an attentional food bias in AN.

Received 13 October 2016; Revised 5 July 2017; Accepted 5 July 2017; First published online 23 August 2017

Key words: Anorexia nervosa, cognitive control, eating disorder, food, reward processing, social, subliminal stimulation.

Introduction

Anorexia nervosa (AN) is characterized by extreme restriction of caloric intake despite severe bodily emaciation. This asceticism is not limited to food, but also expressed in denial of other primary needs and desires, such as intimate and personal relationships (Schmidt & Treasure, 2006). In search for neurobiological mechanisms underlying these puzzling behaviors, researchers

have begun to focus on the reward system (O'Hara *et al.* 2015). The reward system, encompassing cortical as well as subcortical structures including striatum, amygdala, hypothalamus, insula, and orbitofrontal cortex (OFC), processes (incentive) salience, (dis) pleasure, and positive reinforcement (Berridge & Kringelbach, 2015). However, when investigating the processing of reward-related stimuli, the reward system should not be considered in isolation. Several studies using an array of tasks suggest that cognitive processes can modulate (enhance or inhibit) the experience of reward-related stimuli (Goschke, 2014; Botvinick & Braver, 2015). Brain regions that have been associated with such processes including attention and inhibition are the lateral prefrontal cortex (LPFC) and the parietal cortex (Botvinick & Cohen,

* Address for correspondence: S. Ehrlich, M.D., Division of Psychological and Social Medicine and Developmental Neurosciences, University Hospital Carl Gustav Carus, Technische Universität Dresden, Dresden, Germany and Translational Developmental Neuroscience Section, Fetscherstraße 74, Dresden 01307, Germany. (Email: transden.lab@uniklinikum-dresden.de)

2014; Cole *et al.* 2014). When choosing between healthy and tasty food, the neural integration of competing goal values in OFC has been shown to be modulated by activity in the LPFC (Hare *et al.* 2009); a brain region broadly implicated in top-down biasing (Miller & Cohen, 2001) and self-control (Heatherton & Wagner, 2011). Its role as a key regulator in the control of food choice is also underlined by a transcranial magnetic stimulation study, which demonstrated changed preference ratings for high-caloric food items after inhibitory stimulation to the LPFC (Camus *et al.* 2009).

Previous studies on reward processing in AN have reported alterations in reward-related brain regions (Wagner *et al.* 2007b; Wierenga *et al.* 2015) but also in the cognitive control system (Ehrlich *et al.* 2015; King *et al.* 2016). While some of these studies have focused on monetary rewards, neuroimaging studies applying taste- or food-related stimuli in AN confirmed alterations in both networks, albeit as increased and decreased activation (Joos *et al.* 2011; Kim *et al.* 2012; Frank *et al.* 2013; Sanders *et al.* 2015). A recent study (Foerde *et al.* 2015) suggested that alterations in the circuit between LPFC and dorsal striatum may underpin maladaptive food choice in AN. Although processing of social reward in AN has received relatively less attention, two recent functional imaging studies reported reduced blood oxygenation level-dependent (BOLD) responses in the prefrontal (Via *et al.* 2015) and parietal (McAdams *et al.* 2015) regions when processing positive social feedback. In summary, these findings remain inconclusive and we lack a clear understanding of reward-related processes in AN.

Despite the relative lack of substantiated findings, different etiological models have been developed to create a comprehensive framework describing interactions between the reward system and the cognitive control system and their contribution to the pathogenesis of AN. A recent model by O'Hara *et al.* (2015) proposes increased reward responsiveness in the striatal dopamine system toward AN-related cues and behaviors. As in other models (Walsh, 2013; Steinglass & Walsh, 2016), it is postulated that a hyper-responsive striatal dopamine system promotes a learning process of pathological behavior. Over time, these learned behaviors may turn into robust habits and impede the treatment of the illness. In contrast, others (Kaye *et al.* 2009; Brooks *et al.* 2011a) have emphasized the role of an overactive prefrontal control system in AN and hypothesize that this system inhibits motivational drives or compensates for primary deficits in limbic regions.

To gain a deeper understanding of reward and cognitive control processes in AN, it would be helpful to examine the role of the reward system uncoupled from modulation by the cognitive control system

during reward processing. One opportunity that may offer a certain degree of decoupling is subliminal stimulation (Zedelius *et al.* 2011; Mudrik *et al.* 2014). By definition, subliminal processing is a condition where bottom-up stimulation is less likely to provoke robust and outlasting fronto-parietal responses (Dehaene *et al.* 2006). In line with that, research using masked priming paradigms suggests that primarily supraliminal stimuli gain access to a 'global workspace', which can be used as input for a variety of higher cognitive processes such as inhibition (Baars, 2002; Dehaene *et al.* 2006). Meta-analytic data of healthy and clinical samples (Brooks *et al.* 2012b) indicate that subliminal stimulation with arousing stimuli provokes activation in subcortical regions including the amygdala and thalamus independently from higher order cortical regions, but also the cingulo-insular network thought to integrate autonomic and sensory information to determine their relevance (Seeley *et al.* 2007). However, another meta-analysis unexpectedly did not find subcortical regions to be associated with subliminal processing (Meneguzzo *et al.* 2014). Subliminal stimulation nevertheless remains a potentially powerful tool for investigating the early response of the reward system without strong modulating influence of the fronto-parietal control system in health and disease (Childress *et al.* 2008; Brooks *et al.* 2012b). Accordingly, previous work in AN showed that supraliminal but not subliminal distractor stimuli interfered with working memory task performance (Dickson *et al.* 2008) as well as performance in a Stroop task (Sackville *et al.* 1998). In a new paradigm, we present subliminal and supraliminal reward-related stimuli to differentiate early bottom-up subcortical responses from later, top-down cortical responses. If AN patients were predominantly characterized by abnormalities in the reward system, one would expect alterations in the corresponding brain regions during subliminal stimulation, while alterations in consequence of a regulation mechanism of the cognitive control system would only be evident under the supraliminal stimulation. In the latter case, modulating influence of the cognitive control system, which is hypothesized to be hyperactive during the supraliminal processing of reward-related stimuli in AN, would become visible by increased functional connectivity between brain areas of the cognitive control and reward system.

Methods

Participants

The sample of the current study consisted of a total of 70 female volunteers: 35 patients with acute AN

according to the Diagnostic and Statistical Manual of Mental Disorders, Fifth Edition (DSM-V) (12–29 years old) and 35 female healthy controls (HC) (12–29 years old). Within the AN group, 33 were of the restrictive and two of the binge/purging subtype; three had comorbid psychiatric disorders (one patient with depressive disorders, one with anxiety disorder, and one with obsessive compulsive disorder). All patients were admitted to eating disorder programs of a university child and adolescent psychiatry and psychosomatic medicine department and were assessed within 96 h after beginning a behaviorally oriented nutritional rehabilitation program. HC participants had to be of normal weight and eumenorrheic and without any history of psychiatric illness. HC were recruited through advertisement among pupils and university students.

We applied several additional exclusion criteria for each group (see online Supplementary Material 1.1) – most importantly psychotropic medication within 4 weeks prior to the study, binge eating, or diagnosis of bulimia nervosa, substance abuse, neurologic or medical conditions.

This study was approved by the local Institutional Review Board, and all participants (and if underage their guardians) gave written informed consent.

Clinical measures

For all participants, current diagnoses of eating disorders were ascertained by evaluation of the expert form of the Structured Interview of AN and Bulimia Nervosa (Fichter & Quadflieg, 1999). Eating disorder-specific psychopathology was assessed with the short version of the Eating Disorders Inventory (EDI-2; Paul & Thiel, 2005).

Self-reported appetite, including the subscale ‘hunger’, was measured with the use of a visual analog scale (Blundell et al. 2010; see online Supplementary Material 1.2).

Task

During functional magnetic resonance imaging (fMRI), participants passively viewed streams of food stimuli, social stimuli, and neutral stimuli presented either subliminally or supraliminally (referred to as ‘condition’) with an Acer projector P5290DLP capable of presenting stimuli for 17 ms. The paradigm was divided into four equally long blocks in which the first two involved subliminal stimulation and the last two supraliminal stimulation. Subliminal blocks were shown first as familiarity of the stimuli may promote conscious processing even under subliminal stimulation condition (Mudrik et al. 2014). Every block consisted of nine mini-blocks (three of each stimulus

category in pseudo-randomized order), each composed of 10 trials. In the subliminal trials, the stimuli were presented for 17 ms, followed by a mask for 150 ms, and a cross-hair (fixation) for 1309 ms. In the supraliminal trials, stimuli were presented for 500 ms, followed by a cross-hair presented for 973 ms (see online Supplementary Material 1.3).

Stimuli

Thirty neutral and 30 social stimuli were selected from the International Affective Picture System (Lang et al. 1999) and the database EmoPics (Wessa et al. 2010). Neutral stimuli showed, e.g. plants or household items, while social stimuli showed happy social sceneries (e.g. playing children). The 30 food stimuli employed in this study originated from a dataset by Kroemer et al. (2013). The picture selection was based on the following criteria: (1) social stimuli needed to be free of eating disorder-relevant content (e.g. pictures showing women in bikini were excluded), (2) only neutral and social stimuli with valence and arousal ratings that were similar in AN patients and HC (established in an independent pilot study) were selected, (3) the stimuli of all three conditions had to be similar regarding entropy of intensity distribution and brightness to exclude that the perception is biased by differing stimulus properties. The mask used in the subliminal trials was created by scrambling fractioned pieces of the original stimuli and were similar with respect to entropy and brightness to the original stimuli.

Visibility test

After a short break, a second task was carried out in the scanner during structural scanning (to simulate similar conditions to those during the main task) to measure the participant’s ability to recognize subliminal stimuli. This visibility test adapts the structure of the above described subliminal trial, but was followed by a forced choice, where the participants had to identify by button press the preceding subliminally presented stimulus (target) out of three distractor stimuli, which were displayed at the same time for 3000 ms.

Structural and functional image acquisition

Data were acquired between 8:00 and 9:00 after an overnight fast using a standard 3 T Siemens Trio, equipped with a standard 12-channel head coil. T1-weighted structural brain scans were acquired with rapid acquisition gradient echo sequence with the following parameters: number of slices = 176; repetition time = 1900 ms; echo time = 2.26 ms; flip angle = 9°; slice thickness = 1 mm; voxel size = 1 mm × 1 mm × 1

mm; field-of-view (FoV) = 256 mm × 224 mm; bandwidth = 2004 Hz/pixel.

Functional images were acquired by using a gradient-echo T2*-weighted echo planar imaging with the following parameters: tilted 30° toward AC–PC line (to reduce signal dropout in orbitofrontal regions); number of volumes = 190; number of slices = 40; repetition time = 2410 ms; echo time = 25 ms; flip angle of 80°; 3.4 mm in-plane resolution; slice thickness of 2 mm (1 mm gap resulting in a voxel size of 3 × 3 × 2 mm); FoV = 192 × 192 mm; bandwidth of 2112 Hz/pixel.

Image data preprocessing

Functional and structural images were processed using SPM8 toolbox (<http://www.fil.ion.ucl.ac.uk/spm/>) within the Nipype framework (<http://nipy.sourceforge.net/nipype/>). The slice time corrected functional data were realigned and registered to their mean. The realigned files were coregistered to the subject's structural brain image. A DARTEL template was created using structural images from all subjects (Ashburner, 2007). The echo planar image (EPI) volumes were then normalized to Montreal Neurological Institute (MNI) space using the DARTEL template and corresponding flow field. The resulting data were smoothed with an isotropic 8 mm full-width at half maximum (FWHM) Gaussian kernel.

We evaluated the quality of the fMRI data by manual inspection and using artifact detection tools (Whitfield-Gabrieli *et al.* 2009). Volumes that exceed an intensity threshold of three standard deviations or a threshold of 2 mm normalized movement in any direction were classified as outliers [motion outlier: AN patients = 3.21 (6.96), HC = 8.35 (19.85); intensity outlier: AN patients = 6.73 (3.90), HC = 8.35 (4.89)]. The two groups did not differ regarding numbers of motion and intensity outliers [motion outlier: $T(65) = 1.41$, $p = 0.17$; intensity outlier: $T(65) = 1.50$, $p = 0.14$].

Statistical analyses

On the first level, a general linear model was fitted to the hemodynamic response to each of the six combinations of stimulation conditions and stimulus types: supraliminal food stimuli (food_{supra}), subliminal food stimuli (food_{sublim}), supraliminal social stimuli (social_{supra}), subliminal social stimuli (social_{sublim}), supraliminal neutral stimuli (neutral_{supra}), and subliminal neutral stimuli (neutral_{sublim}) using boxcar functions with duration of 15 s (epoch-related design). Additionally, six realignment parameters and outlier volumes identified by quality control as described above were included as nuisance regressors of no interest. On the second level, a linear mixed model

including a two-level within-subject variable (stimulation condition: supraliminal and subliminal), a three-level within-subject variable (stimulus type: food, neutral, and social), and a binary between-subject variable (group: AN patients, HC) were estimated using the GLMflex toolbox (http://mrtools.mgh.harvard.edu/index.php/GLM_Flex). To confirm that the task worked as intended, we first examined stimulus type-specific activation patterns within both the supraliminal and the subliminal stimulation conditions by calculating the contrasts social_{sublim} > neutral_{sublim}, food_{sublim} > neutral_{sublim} as well as social_{supra} > neutral_{supra} and food_{supra} > neutral_{supra} in HC. To address our main research question regarding differences between the groups when processing the different stimulus types under the two stimulation conditions, we examined the *group × stimulation condition × stimulus type* interaction effect. If AN patients were predominantly characterized by abnormalities in the reward system, we would predict alterations in those brain regions during subliminal stimulation condition, while alterations in consequence of top-down regulation mechanisms would be expected to be mirrored by alterations in cognitive control areas under the supraliminal stimulation condition.

To guard against type I errors, results were corrected based on permutations using the updated version of 3DClustSim, released August 2016 (http://afni.nimh.nih.gov/pub/dist/doc/program_help/3dClustSim.html). This tool runs Monte Carlo simulations to estimate the cluster size at which the false-positive probability is below an α level of 0.05 for a voxelwise p value level, which was set at $p = 0.005$. The simulations resulted in a threshold for minimal cluster size of $\kappa = 468$ (whole-brain).

Region-of-interest (ROI) analyses

To further specify group differences revealed by the whole-brain analysis of the *group × stimulation condition × stimulus type* interaction, we extracted condition-specific β values for each participant from all voxels belonging to significant clusters using MarsBaR toolbox for SPM (Brett, Anton, Valabregue, & Poline, 2002). To explore what factors contribute to the group differences revealed by the whole-brain analysis of the *group × stimulation condition × stimulus type* interaction effect, extracted β values were subjected to a linear mixed model using SPSS. Predictors in the model included the fixed-effects group, stimulation condition, stimulus type and their interaction, and a by-subject random adjustment for the intercept. Bonferroni correction for multiple comparisons was used for *post hoc* tests.

The individual sensitivity toward subliminal stimuli in the visibility test was estimated using d' , a measure

of the ability to discriminate between target items and distractor items in a four-alternative forced choice task (Hacker & Ratcliff, 1979).

Associations between BOLD activity and clinical variables, as well as the individual sensitivity (d') to detect subliminal targets were investigated using Pearson's r separately in each group. The latter association is of interest, as one might assume that the neural response during subliminal stimulation might be modulated by the individual ability to perceive subliminal targets.

Connectivity analysis

To follow-up on results revealed by the analysis described above, we investigated changes in functional connectivity between a brain region identified as a putative source of top-down control signals, the inferior frontal junction (IFJ; see results), using the generalized psychophysiological interaction approach (gPPI; McLaren *et al.* 2012). This approach allowed us to assess whether the effect of stimulus type (psychological factors: food_{supra}, neutral_{supra}, social_{supra}) is modulated by the activity of a source region (physiological factor). We chose to investigate the supraliminal condition only since potential modulating effects of prefrontal brain regions were only evident during this condition. The investigation of alterations in functional connectivity comprised two steps: First, based on existing hypothesis of hyperactive cognitive control over limbic brain regions in AN (Kaye *et al.* 2009), we explored changes in functional connectivity between the source region IFJ and limbic target regions: ventral striatum, amygdala, OFC, and insula. Second, an exploratory approach with no *a priori* defined target regions of the top-down control signal (see online Supplementary Material 1.4).

Results

Sample characteristics

As shown in Table 1, there were no group differences for age, IQ, and d' (parameter of the visibility test). As expected, AN patients had a lower body mass index, higher EDI-2 total scores, and reported less hunger at the time of the scanning session than HC.

Imaging results

Stimulus-specific activation patterns were found for both, the supraliminal as well as the subliminal stimulation condition, i.e. increased BOLD responses in reward-related brain regions, such as the ventral striatum for social *v.* neutral stimuli types in HC (see online Supplementary Material 2.1).

Table 1. Sociodemographic and clinical variables of the two groups

	AN (<i>n</i> = 35)	HC (<i>n</i> = 35)	<i>t</i>	<i>P</i>
Age	16.25 ± 3.46	16.31 ± 3.39	−0.08	0.94
BMI	14.59 ± 1.50	20.46 ± 2.06	−13.62	<0.001
BMI-SDS	−3.23 ± 1.18	−0.05 ± 0.72	−13.62	<0.001
IQ	114.70 ± 11.57	112.86 ± 7.64	0.77	0.45
Age of onset	14.38 ± 1.93	N.A.	–	–
Duration of current episode (in month)	12.83 ± 19.47	N.A.		
EDI-2 total	211.95 ± 42.94	143.77 ± 30.49	7.66	<0.001
d'	0.08 ± 0.34	0.09 ± 0.17	0.13	0.90
Hunger (subjective)	3.84 ± 3.01	6.61 ± 2.20	4.33	<0.001

d' = index of sensitivity toward subliminal stimuli (a value of zero indicates a performance at chance level; d' values of HC are not different from zero [$T(34) = 1.451$; $p = 0.153$] but d' values of AN patients [$T(34) = 3.242$; $p = 0.003$]), Hunger was assessed with the use of a visual analogue scale ranging from 'not at all' (value of 0) to 'extremely' (value of 10). Group differences were tested using Student's t tests. Displayed are means ± standard deviations; AN, anorexia nervosa; HC, healthy controls; BMI, body mass index; BMI-SDS, body mass index standard deviation score; IQ, intelligence quotient; EDI-2, Eating Disorder Inventory 2.

Analysis of the critical group × stimulation condition × stimulus type interaction showed significant clusters in a region of IPFC centered at the IFJ (Fig. 1a), the superior occipital gyrus (SOG; Fig. 1b) and the fusiform gyrus/parahippocampal gyrus (FFG/PG; Fig. 1c). To explore what factors contribute to the group differences, parameter estimates (β values) were extracted and subjected to a linear mixed model. Analysis of activation within these clusters revealed that the three-way interaction was driven by distinct factors in the respective ROIs. In the IFJ, while a two-way group × stimulation condition interaction (see Table 2) driven by increased activity in AN *v.* HC in the supraliminal condition was evident [$T(96.37) = -3.14$; $p = 0.001$], the three-way interaction effect was driven by a differential influence of stimulus type in HC. Specifically, HC showed an increased activity of the IFJ for social_{supra} in comparison to neutral_{supra} stimuli [$T(340) = 6.27$; $p < 0.001$] and food_{supra} [$T(340) = 8.93$; $p < 0.001$]. In sum, this indicates that, within the given task context, in HC the IFJ responds stronger to social stimuli, while this brain region is strongly active irrespectively of the stimulus type in patients with AN.

In the SOG, the three-way interaction was driven by a differential influence of the stimulus type under supraliminal stimulation within AN, while no

Fig. 1. Left, significant (family-wise error-corrected) clusters revealed by the whole-brain three-way *group* $\times$ *stimulation condition* $\times$ *stimulus type* interaction analysis: (a) cluster at IFJ [46, 6, 26]; (b) cluster at SOG [−30, −90, 28]; (c) cluster at FFG/PHG [−26, −90, 28]; right mean activation (β -estimates $\pm$ S.E.M.) extracted from the respective significant clusters identified by the whole-brain analysis are plotted for both groups under each stimulation condition and for each stimulus type; ** = significant difference of extracted β estimates at $p = 0.001$; IFJ, inferior frontal junction; SOG, superior occipital gyrus; FFG/PHG, fusiform gyrus/parahippocampal gyrus.

Table 2. Main effects and interaction effect of the linear mixed model (based on extracted β -values for each relevant cluster)

	IFJ <i>F</i> (<i>p</i>)	SOG <i>F</i> (<i>p</i>)	FFG/PHG <i>F</i> (<i>p</i>)
Group	5.07 (0.028)	0.10 (0.756)	0.30 (0.585)
Stimulation condition	85.89 (<0.001)	558.00 (<0.001)	974.241 (<0.001)
Stimulus type	12.40 (<0.001)	8.07 (<0.001)	10.907 (<0.001)
Group $\times$ stimulation condition	12.07 (0.001)	0.73 (0.394)	0.55 (0.459)
Group $\times$ stimulus type	8.85 (<0.001)	3.45 (0.033)	3.36 (0.036)
Stimulation condition $\times$ stimulus type	2.18 (0.115)	6.35 (0.002)	2.80 (0.062)
Group $\times$ stimulation condition $\times$ stimulus type	11.722 (<0.001)	6.073 (0.003)	10.907 (<0.001)

IFJ, inferior frontal junction; SOG, superior occipital gyrus; FFG/PHG, fusiform gyrus/parahippocampal gyrus.

differential effect of stimulus type under supraliminal stimulation was evident in HC. In detail, AN patients showed an increased response to food_{supra} in comparison to neutral_{supra} [$T(340) = 3.27$; $p = 0.004$] and social_{supra} [$T(340) = 6.75$; $p < 0.001$] as well as to social_{supra} in comparison to neutral_{supra} [$T(340) = -4.02$; $p = 0.004$]. A similar pattern was found for the FFG/PHG, the three-way interaction was modulated by the type of stimulus under supraliminal stimulation in AN, while HC showed no differential response to stimulus type under supraliminal

stimulation condition. Within the AN group, the FFG/PHG showed increased activity for food_{supra} in comparison to neutral_{supra} [$T(340) = 4.98$; $p < 0.001$] and social_{supra} [$T(340) = 5.52$; $p < 0.001$] but not for social_{supra} in comparison to neutral_{supra}. Control analyses accounting for (1) the relatively broad age range (online Supplementary Material 2.2) and (2) patients comorbidities ($n = 3$; online Supplementary Material 2.3) and (3) AN binge/purge subtype ($n = 2$; online Supplementary Material 2.4), revealed statistically identical results.

Association of hemodynamic activity, clinical variables, and sensitivity index (d')

Given the broad role of the IFJ in executive functions (Brass *et al.* 2005; Derrfuss *et al.* 2005) and the hypothesis of elevated cognitive control in AN (Kaye *et al.* 2009), we next inspected relationships between neural responses in this region and eating disorder symptoms as gauged by the EDI-2. Since we established a group difference in hemodynamic activity during supraliminal stimulation condition irrespective of stimulus type, we explored the association between the average BOLD activity and EDI-2 in patients with AN, which was found to be significant ($r = 0.402$, $p = 0.017$).

Next, we tested for a potential association between d' and the IFJ response during each stimulus type under subliminal stimulation condition. Interestingly, IFJ activation during subliminal stimulation condition of social stimuli was associated with the probability to detect subliminal targets ($r = 0.496$, $p = 0.002$) in HC. This indicates that with increasing individual visibility under subliminal stimulation condition neural pattern indicative for supraliminal processing emerge.

Connectivity analyses

To explore whether the IFJ serves as a putative source region of top-down control during the supraliminal stimulation condition, we conducted a gPPI analysis using the IFJ as seed region and $\text{food}_{\text{supra}}$, $\text{social}_{\text{supra}}$ and $\text{neutral}_{\text{supra}}$ as psychological factors. The exploratory approach with no *a priori* defined target regions, as well as the hypothesis-based approach with the four limbic target regions revealed no significant group differences in functional connectivity.

Discussion

The present study analyzed neural responses to subliminally and supraliminally presented reward-related food and social stimuli in AN. The objective of our experimental design and analysis strategy was to determine whether differences in brain response to these stimuli are predominantly present in subcortical (bottom-up) or cortical (top-down) networks. As a first step, we confirmed that the subliminal and supraliminal processing of rewarding stimuli elicited the expected reward-related neural activation pattern in HC (Dehaene *et al.* 2006; Brooks *et al.* 2011b; Zedelius *et al.* 2011; Mudrik *et al.* 2014; Wang *et al.* 2016).

Contrasting the two groups, we found no difference under the subliminal stimulation condition, suggesting that adolescent (non-chronic) AN patients may not have particularly strong alterations in the bottom-up neural response to rewarding stimuli. In response to the supraliminal stimulation condition, we found

activation to be elevated in the AN group in a region of IPFC associated with cognitive control, the IFJ (Brass *et al.* 2005; Derrfuss *et al.* 2005). IFJ activation was increased in the AN group irrespective of stimulus type – suggestive of a sustained response – which was associated with the level of eating disorder symptoms. HC, however, showed comparably elevated activity in this region only to social stimuli. These findings are in accordance with studies reporting no interference with working memory or performance monitoring when distracting food stimuli are presented subliminally (Brooks *et al.* 2012a), while supraliminal distractors, irrespective of salience, seem to interfere with working memory performance in patients with AN (Dickson *et al.* 2008). The authors of the latter study argued that this might indicate that AN patients are more easily distracted due to their extreme attention to details – an interpretation which dovetails with our finding of a sustained elevated IFJ activation.

In contrast to the notion of altered reward processing in AN (O'Hara *et al.* 2015), no group differences were evident in the reward system, neither during subliminal nor supraliminal stimulation. This seems surprising given that previous studies using food stimuli or monetary rewards in AN demonstrate alterations in those brain areas (Wagner *et al.* 2007a; Cowdrey *et al.* 2011). A potential explanation for this discrepancy might be that alterations in the reward system evolve with the chronic manifestation of the disorder and potential alterations in this system have not (yet) developed in our primarily young non-chronic sample (85% younger than 18 years; 74% with a duration of illness shorter than 1 year). Indeed, other studies investigating adolescent AN reported no difference in the reward system during processing of monetary rewards (Bischoff-Grethe *et al.* 2013; Ehrlich *et al.* 2015; King *et al.* 2016). This resonates with the argument postulated by the reward-centered model of AN that patients do not suffer from a generalized inability to experience reward, but that processes within this system change with the progress of the disorder (O'Hara *et al.* 2015). However, to investigate this notion in more detail, future studies contrasting chronic and non-chronic patients are needed.

As mentioned above, we found a neural pattern suggestive of sustained increased IFJ responsivity in AN during supraliminal processing, which was associated with eating disorder psychopathology. The IFJ, a region located at the junction of the inferior frontal sulcus and the inferior precentral sulcus (Derrfuss *et al.* 2005), plays a pivotal role in a variety of cognitive control functions, including selective attention (Baldauf & Desimone, 2014) and updating task representations (Derrfuss *et al.* 2005). This finding provides support for the notion of AN as model disorder for excessive

cognitive control (Kaye *et al.* 2009). However, according to the framework of Kaye *et al.* (2009) and Brooks *et al.* (2011a), one could have expected that an overactive cognitive control system in AN has a top-down modulatory effect on limbic brain regions implicated in motivational drives. In this study, we found: (1) no alteration in functional connectivity between the IFJ and the reward system and (2) the elevated neural response of the IFJ in patients with AN was not limited to reward-related stimuli. The unspecific response of the IFJ may indicate an attempt of sustained control in AN patients once a meaningful picture is presented, while in HC such a cognitive control reaction is only evident in response to emotionally salient stimuli. One might also speculate that AN patients are less able to disengage cognitive control reactions.

The IFJ has also been associated with attentional allocation. Studies have demonstrated neural synchrony between the IFJ and posterior sensory brain areas (Zanto *et al.* 2010; Sundermann & Pfeiderer, 2012). Within this network, the IFJ is proposed as the source of attentional modulation, facilitated by long-distance projections, linking the frontal cortex with the visual association areas. By this means, top-down modulation facilitates neural excitability toward selected stimuli, thus forming perceptual biases (Baldauf & Desimone, 2014; Xu, 2014). Interestingly, we found increased activity in brain areas associated with visual processing, namely the SOG (dorsal visual stream) and the FFA/PHG (ventral visual stream), during supraliminal processing of food stimuli in AN patients. These findings can be interpreted in light of an attentional bias to food stimuli that has been argued to be a maintenance factor for AN (Shafran *et al.* 2007). Nevertheless, we found no evidence for altered coupling between the IFJ and these brain areas when processing food stimuli. Second, although we discovered increased activation in visual areas in AN when processing supraliminally presented food stimuli, BOLD activity in the IFJ was elevated when processing all stimulus types.

Limitations

Our study has to be interpreted in light of the following limitations. First, it is plausible that the lack of group differences in the subliminal stimulation condition may be due to a relative insufficiency of our task to evoke reward-related neural processes. This possibility seems unlikely, given that subliminal stimulation did elicit an expected pattern of stimulus-specific activation in HC (see online Supplementary Material 2.1). Second, when studying AN patients, we cannot distinguish between predisposing factors and consequences

of the acute undernutrition. Future studies will need to investigate recovered patients or apply longitudinal study designs. Third, HC were scanned at different phases of their menstrual cycle. Variance in sex hormone levels could have influenced our ability to uncover potential group differences in reward-related brain regions (Toffoletto *et al.* 2014).

A strength of our study, however, is the large homogenous sample of young acute AN patients, who have had a short duration of illness.

Conclusions

In summary, this study suggests that adolescent AN patients may not (yet) show alterations in the brain reward system, while we found a hyperactivity of a brain region implicated in cognitive control during the supraliminal processing of all stimulus types. The latter finding is remarkable, since the passive viewing task does not require any control functions *per se*. These findings thus lend further support for the hypothesis of sustained and elevated (self-) control in AN (Kaye *et al.* 2009; Brooks *et al.* 2011a). Since we also found increased neural response in visual brain areas to food stimuli in patients, our findings may also be interpreted in light of biased attention toward food stimuli or increased attention to details (weak central coherence; Lopez *et al.* 2008). These findings open up new avenues for cognitive behavioral or neuromodulatory therapeutic approaches which try to target the tendency of AN patients to 'stay in control' even when it is not necessary (Pop-Jordanova, 2000; Wildes & Marcus, 2011; McClelland *et al.* 2013).

Supplementary material

The supplementary material for this article can be found at <https://doi.org/10.1017/S0033291717002161>.

Acknowledgements

This work was supported by the Deutsche Forschungsgemeinschaft (EH 367/5-1 & SFB 940) and the Technische Universität Dresden. The authors thank the Center for Information Services and High Performance Computing (ZIH) at TU Dresden for generous allocations of computer time.

Declaration of Interest

In the last two years, Dr Roessner has received lecture fees from Eli Lilly, Janssen-Cilag, Medice, and Novartis and was a member of advisory boards of Eli Lilly, Novartis. All other authors declare that there are no conflicts of interest in relation to the subject of this study.

References

- Ashburner J (2007). A fast diffeomorphic image registration algorithm. *Neuroimage* **38**, 95–113.
- Baars BJ (2002). The conscious access hypothesis: origins and recent evidence. *Trends in Cognitive Sciences* **6**, 47–52.
- Baldauf D, Desimone R (2014). Neural mechanisms of object-based attention. *Science* **344**, 424–427.
- Berridge KC, Kringelbach ML (2015). Pleasure systems in the brain. *Neuron* **86**, 646–664.
- Bischoff-Grethe A, McCurdy D, Grenesko-Stevens E, Irvine LE, Wagner A, Wendy Yau W-Y, Fennema-Notestine C, Wierenga CE, Fudge JL, Delgado MR (2013). Altered brain response to reward and punishment in adolescents with anorexia nervosa. *Psychiatry Research: Neuroimaging* **214**, 331–340.
- Blundell J, De Graaf C, Hulshof T, Jebb S, Livingstone B, Lluch A, Mela D, Salah S, Schuring E, Van Der Knaap H, Westerterp M (2010). Appetite control: methodological aspects of the evaluation of foods. *Obesity Reviews* **11**, 251–270.
- Botvinick M, Braver T (2015). Motivation and cognitive control: from behavior to neural mechanism. *Annual Review of Psychology* **66**, 83–113.
- Botvinick MM, Cohen JD (2014). The computational and neural basis of cognitive control: charted territory and new frontiers. *Cognitive science* **38**, 1249–1285.
- Brass M, Derrfuss J, Forstmann B, von Cramon DY (2005). The role of the inferior frontal junction area in cognitive control. *Trends in Cognitive Sciences* **9**, 314–316.
- Brett M, Anton J, Valabregue R, Poline J (2002). Region of interest analysis using an SPM toolbox. Paper presented at 8th International Conference on Functional Mapping of the Human Brain, Sendai, Japan.
- Brooks S, O'Daly O, Uher R, Schiöth H, Treasure J, Campbell I (2012a). Subliminal food images compromise superior working memory performance in women with restricting anorexia nervosa. *Consciousness and Cognition* **21**, 751–763.
- Brooks S, Owen G, Uher R, Friederich H, Giampietro V, Brammer M, Williams S, Schiöth H, Treasure J, Campbell I (2011a). Differential neural responses to food images in women with bulimia versus anorexia nervosa. *PLoS ONE* **6**, e22259.
- Brooks S, Prince A, Stahl D, Campbell I, Treasure J (2011b). A systematic review and meta-analysis of cognitive bias to food stimuli in people with disordered eating behaviour. *Clinical Psychology Review* **31**, 37–51.
- Brooks SJ, Savov V, Allzén E, Benedict C, Fredriksson R, Schiöth HB (2012b). Exposure to subliminal arousing stimuli induces robust activation in the amygdala, hippocampus, anterior cingulate, insular cortex and primary visual cortex: a systematic meta-analysis of fMRI studies. *NeuroImage* **59**, 2962–2973.
- Camus M, Halelamien N, Plassmann H, Shimojo S, O'Doherty J, Camerer C, Rangel A (2009). Repetitive transcranial magnetic stimulation over the right dorsolateral prefrontal cortex decreases valuations during food choices. *European Journal of Neuroscience* **30**, 1980–1988.
- Childress AR, Ehrman RN, Wang Z, Li Y, Sciortino N, Hakun J, Jens W, Suh J, Listerud J, Marquez K, Franklin T, Langleben D, Detre J, O'Brien CP (2008). Prelude to passion: limbic activation by 'unseen' drug and sexual cues. *PLoS ONE* **3**, e1506.
- Cole MW, Repovš G, Anticevic A (2014). The frontoparietal control system A central role in mental health. *The Neuroscientist* **20**, 652–664.
- Cowdrey FA, Park RJ, Harmer CJ, McCabe C (2011). Increased neural processing of rewarding and aversive food stimuli in recovered anorexia nervosa. *Biological Psychiatry* **70**, 736–743.
- Dehaene S, Changeux J-P, Naccache L, Sackur J, Sergent C (2006). Conscious, preconscious, and subliminal processing: a testable taxonomy. *Trends in Cognitive Sciences* **10**, 204–211.
- Derrfuss J, Brass M, Neumann J, von Cramon DY (2005). Involvement of the inferior frontal junction in cognitive control: meta-analyses of switching and stroop studies. *Human Brain Mapping* **25**, 22–34.
- Dickson H, Brooks S, Uher R, Tchanturia K, Treasure J, Campbell I (2008). The inability to ignore: distractibility in women with restricting anorexia nervosa. *Psychological Medicine* **38**, 1741–1748.
- Ehrlich S, Geisler D, Ritschel F, King JA, Seidel M, Boehm I, Breier M, Clas S, Weiss J, Marxen M, Smolka MN, Roessner V, Kroemer NB (2015). Elevated cognitive control over reward processing in recovered patients with anorexia nervosa. *Journal of Psychiatry and Neuroscience* **40**, 307–315.
- Fichter M, Quadflieg N (1999). *SIAB. Strukturiertes Inventar fuer Anorektische und Bulimische Essstoerungen Nach DSM-IV und ICD-10*. Huber: Bern.
- Foerde K, Steinglass JE, Shohamy D, Walsh BT (2015). Neural mechanisms supporting maladaptive food choices in anorexia nervosa. *Nature Neuroscience* **18**, 1571–1573.
- Frank G, Shott ME, Hagman JO, Mittal VA (2013). Alterations in brain structures related to taste reward circuitry in ill and recovered anorexia nervosa and in bulimia nervosa. *American Journal of Psychiatry* **170**, 1152–1160.
- Goschke T (2014). Dysfunctions of decision-making and cognitive control as transdiagnostic mechanisms of mental disorders: advances, gaps, and needs in current research. *International Journal of Methods in Psychiatric Research* **23**, 41–57.
- Hacker MJ, Ratcliff R (1979). A revised table of d' for M-alternative forced choice. *Perception & Psychophysics* **26**, 168–170.
- Hare TA, Camerer CF, Rangel A (2009). Self-control in decision-making involves modulation of the vmPFC valuation system. *Science* **324**, 646–648.
- Heatherton TF, Wagner DD (2011). Cognitive neuroscience of self-regulation failure. *Trends in Cognitive Sciences* **15**, 132–139.
- Joos AAB, Saum B, van Elst LT, Perlov E, Glauche V, Hartmann A, Freyer T, Tüscher O, Zeeck A (2011). Amygdala hyperreactivity in restrictive anorexia nervosa. *Psychiatry Research: Neuroimaging* **191**, 189–195.
- Kaye W, Fudge J, Paulus M (2009). New insights into symptoms and neurocircuit function of anorexia nervosa. *Nature Reviews Neuroscience* **10**, 573–584.

- Kim KR, Ku J, Lee J-H, Lee H, Jung Y-C (2012). Functional and effective connectivity of anterior insula in anorexia nervosa and bulimia nervosa. *Neuroscience Letters* **521**, 152–157.
- King JA, Geisler D, Bernardoni F, Ritschel F, Böhm I, Seidel M, Mennigen E, Ripke S, Smolka MN, Roessner V, Ehrlich S (2016). Altered neural efficiency of decision making during temporal reward discounting in anorexia nervosa. *Journal of the American Academy of Child & Adolescent Psychiatry* **55**, 972–979.
- Kroemer NB, Krebs L, Kobiella A, Grimm O, Vollstädt-Klein S, Wolfensteller U, Kling R, Bidlingmaier M, Zimmermann US, Smolka MN (2013). (Still) longing for food: insulin reactivity modulates response to food pictures. *Human Brain Mapping* **34**, 2367–2380.
- Lang PJ, Bradley MM, Cuthbert BN (1999). *International Affective Picture System (IAPS): Instruction Manual and Affective Ratings*, vol. 2. Center of Research in Psychophysiology, University of Florida: Gainesville, Florida, USA.
- Lopez C, Tchanturia K, Stahl D, Booth R, Holliday J, Treasure J (2008). An examination of the concept of central coherence in women with anorexia nervosa. *The International Journal of Eating Disorders* **41**, 143–152.
- McAdams CJ, Lohrenz T, Montague PR (2015). Neural responses to kindness and malevolence differ in illness and recovery in women with anorexia nervosa. *Human Brain Mapping* **36**, 5207–5219.
- McClelland J, Bozhilova N, Nestler S, Campbell IC, Jacob S, Johnson-Sabine E, Schmidt U (2013). Improvements in symptoms following neuronavigated repetitive transcranial magnetic stimulation (rTMS) in severe and enduring anorexia nervosa: findings from two case studies. *European Eating Disorders Review* **21**, 500–506.
- McLaren DG, Ries ML, Xu G, Johnson SC (2012). A generalized form of context-dependent psychophysiological interactions (gPPI): a comparison to standard approaches. *NeuroImage* **61**, 1277–1286.
- Meneguzzo P, Tsakiris M, Schioth HB, Stein DJ, Brooks SJ (2014). Subliminal versus supraliminal stimuli activate neural responses in anterior cingulate cortex, fusiform gyrus and insula: a meta-analysis of fMRI studies. *BMC Psychology* **2**, 52–63.
- Miller EK, Cohen JD (2001). An integrative theory of prefrontal cortex function. *Annual Review of Neuroscience* **24**, 167–202.
- Mudrik L, Faivre N, Koch C (2014). Information integration without awareness. *Trends in Cognitive Sciences* **18**, 488–496.
- O'Hara CB, Campbell IC, Schmidt U (2015). A reward-centred model of anorexia nervosa: a focussed narrative review of the neurological and psychophysiological literature. *Neuroscience & Biobehavioral Reviews* **52**, 131–152.
- Paul T, Thiel A (2005). *Eating Disorder Inventory-2 (EDI-2): Deutsche Version*. Hogrefe: Goettingen.
- Pop-Jordanova N (2000). Psychological characteristics and biofeedback mitigation in preadolescents with eating disorders. *Pediatrics International* **42**, 76–81.
- Sackville T, Schotte DE, Touyz SW, Griffiths R, Beumont P (1998). Conscious and preconscious processing of food, body weight and shape, and emotion-related words in women with anorexia nervosa. *International Journal of Eating Disorders* **23**, 77–82.
- Sanders N, Smeets PA, van Elburg AA, Danner UN, van Meer F, Hoek HW, Adan RA (2015). Altered food-cue processing in chronically ill and recovered women with anorexia nervosa. *Frontiers in Behavioral Neuroscience* **9**, 46.
- Schmidt U, Treasure J (2006). Anorexia nervosa: valued and visible. A cognitive-interpersonal maintenance model and its implications for research and practice. *British Journal of Clinical Psychology* **45**, 343–366.
- Seeley WW, Menon V, Schatzberg AF, Keller J, Glover GH, Kenna H, Reiss AL, Greicius MD (2007). Dissociable intrinsic connectivity networks for salience processing and executive control. *The Journal of Neuroscience* **27**, 2349–2356.
- Shafraan R, Lee M, Cooper Z, Palmer R, Fairburn C (2007). Attentional bias in eating disorders. *International Journal of Eating Disorders* **40**, 369–380.
- Steinglass JE, Walsh BT (2016). Neurobiological model of the persistence of anorexia nervosa. *Journal of Eating Disorders* **39**, 267–275.
- Sundermann B, Pfeleiderer B (2012). Functional connectivity profile of the human inferior frontal junction: involvement in a cognitive control network. *BMC Neuroscience* **13**, 119.
- Toffoletto S, Lanzenberger R, Gingnell M, Sundström-Poromaa I, Comasco E (2014). Emotional and cognitive functional imaging of estrogen and progesterone effects in the female human brain: a systematic review. *Psychoneuroendocrinology* **50**, 28–52.
- Via E, Soriano-Mas C, Sánchez I, Forcano L, Harrison BJ, Davey CG, Pujol J, Martínez-Zalacáin I, Menchón JM, Fernández-Aranda F, Cardoner N (2015). Abnormal social reward responses in anorexia nervosa: an fMRI study. *PLoS ONE* **10**, e0133539.
- Wagner A, Aizenstein H, Mazurkewicz L, Fudge J, Frank G, Putnam K, Bailer U, Fischer L, Kaye W (2007a). Altered insula response to taste stimuli in individuals recovered from restricting-type anorexia nervosa. *Neuropsychopharmacology* **33**, 513–523.
- Wagner A, Aizenstein H, Venkatraman V, Fudge J, May J, Mazurkewicz L, Frank G, Bailer U, Fischer L, Nguyen V (2007b). Altered reward processing in women recovered from anorexia nervosa. *American Journal of Psychiatry* **164**, 1842–1849.
- Walsh BT (2013). The enigmatic persistence of anorexia nervosa. *The American Journal of Psychiatry* **170**, 477–484.
- Wang KS, Smith DV, Delgado MR (2016). Using fMRI to study reward processing in humans: past, present, and future. *Journal of Neurophysiology* **115**, 1664–1678.
- Wessa M, Kanske P, Neumeister P, Bode K, Heissler J, Schönfelder S (2010). Emopics: subjektive und psychophysiologische evaluation neuen bildmaterials für die klinisch-biopsychologische forschung. Available from Michele Wessa, michele.wessa@med.uni-heidelberg.de
- Whitfield-Gabrieli S, Thermenos HW, Milanovic S, Tsuang MT, Faraone SV, McCarley RW, Shenton ME, Green AI, Nieto-Castanon A, LaViolette P (2009). Hyperactivity and hyperconnectivity of the default network in schizophrenia and in first-degree relatives of persons with schizophrenia.

- Proceedings of the National Academy of Sciences of the United States of America* **106**, 1279–1284.
- Wierenga CE, Bischoff-Grethe A, Melrose AJ, Irvine Z, Torres L, Bailer UF, Simmons A, Fudge JL, McClure SM, Ely A, Kaye WH** (2015). Hunger does not motivate reward in women remitted from anorexia nervosa. *Biological Psychiatry* **77**, 642–652.
- Wildes JE, Marcus MD** (2011). Development of emotion acceptance behavior therapy for anorexia nervosa: a case series. *International Journal of Eating Disorders* **44**, 421–427.
- Xu Y** (2014). Inferior frontal junction biases perception through neural synchrony. *Trends in Cognitive Sciences* **18**, 447–448.
- Zanto TP, Rubens MT, Bollinger J, Gazzaley A** (2010). Top-down modulation of visual feature processing: the role of the inferior frontal junction. *NeuroImage* **53**, 736–745.
- Zedelius CM, Veling H, Aarts H** (2011). Boosting or choking—how conscious and unconscious reward processing modulate the active maintenance of goal-relevant information. *Consciousness and Cognition* **20**, 355–362.