Aus der Klinik und Poliklinik für Neurologie
Direktor: Herr Prof. Dr. Heinz Reichmann

The societal economic costs of disability worsening and relapses in patients with multiple sclerosis

Dissertationsschrift
Zur Erlangung des akademischen Grades
Doctor of Philosophy (Ph.D.)
vorgelegt
der Medizinischen Fakultät Carl Gustav Carus
der Technischen Universität Dresden

von

Nils-Henning Neß, M.Sc.

aus Wiesbaden

Dresden 2020

1. Gutachter:

2. Gutachter:

Tag der mündlichen Prüfung:

Table of contents

List of figures	I
List of tables	II
Abbreviations	III
1 Introduction	1
1.1 Multiple sclerosis	1
1.1.1 Pathogenesis	1
1.1.2 Clinical manifestation	1
1.1.3 Descriptive epidemiology	2
1.1.4 Clinical hallmarks and outcome measures	2
1.1.5 Clinical course	4
1.1.6 Disease modifying therapies	5
1.1.7 Natural history	6
1.2 Cost of illness research	7
1.2.1 Classification of costs	7
1.2.2 Costing methodologies	8
1.2.3 Process of cost estimation	10
1.3 Cost of multiple sclerosis	14
1.3.1 Instruments for resource use measurement	14
1.3.2 Review of cost of multiple sclerosis studies in Germany	16
1.4 Hypotheses and objective	20
2 Data source	21
2.1 Study setting	21
2.2 Clinical and economic data	22
2.3 Study population	24
2.4 Ethical considerations	24
2.5 Data management	25
3 Methods	26
3.1 Questionnaire development and validation	26
3.1.1 Development process	26
3.1.2 Identification of resource consumption	27
3.1.3 Quantification of resource consumption	28
3.1.4 Valuation of resource consumption	29
3.2 Assessment of psychometric properties	32
3.3 Definition of clinical outcome measures	34
3.4 Assessment of relapse associated costs	35
3.5 Statistical analyses	36
4 Results	38
4.1 The Multiple Sclerosis Health Resource Survey	38

4.1.1	Study population	38
4.1.2	Validation of the questionnaire	39
4.1.3	Resource utilization	42
4.2	Societal economic impact of disability worsening	43
4.2.1	Study population	43
4.2.2	Descriptive cost analyses	45
4.2.3	Associations between disability worsening and costs	46
4.3	Societal economic impact of relapses	49
4.3.1	Study population	49
4.3.2	Relapse associated costs	52
4.3.3	Longitudinal analysis of relapse associated costs	58
5	Discussion	60
5.1	Summary of results	60
5.2	Societal economic impact of disability worsening	61
5.3	Societal economic impact of relapses	62
5.4	Discussion of the Multiple Sclerosis Health Resource Survey	63
5.5	Generalizability of the study results	66
5.6	Limitations of the study	66
6	Conclusion	68
7	Summary	69
7.1	English version	69
7.2	German version	72
	Bibliography	75
	Supplement	89
	Acknowledgement	95
	Anlage 1	96
	Anlage 2	98

List of figures

Figure 1: Characteristics and limitations of the Expanded Disability Status Scale.	4
Figure 2: Classification of the clinical courses of multiple sclerosis.	5
Figure 3: Overview of costing methodologies.	9
Figure 4: Timeline of the pharmacoeconomic (sub)studies PEARL and PANGAEA.	21
Figure 5: Identified direct and indirect cost items for patients with MS.	27
Figure 6: Schematic of the assessment of clinical descriptors.	34
Figure 7: Assessment of changes in resource utilization (prior-, during and post-relapse).	36
Figure 8: Effects (IRR, 95%-CI) of PIRA and RAW on the quarterly economic costs compared to DAF.	47
Figure 9: Comparison of patients with in- and outpatient hospitalizations within two years.	48
Figure 10: Love plot summarizing the covariate balance before and after PS matching.	51
Figure 11: Histogram comparing the density of PS before and after matching.	52
Figure 12: Resource costs (3 months) for relapse inactive (n=597) and relapse active (RAW: n=149; not RAW n=448) patients (before, during and after relapse).	53
Figure 13: Quarterly costs (in €) of direct and indirect resource items before during and post relapse.	59
Figure 14: Development process of the MS-HRS.	65

List of tables

Table 1: Frequency of symptoms in patients with MS by disease duration.	1
Table 2: Exemplary resource items by cost categories.	8
Table 3: Perspective of cost estimation and relevant cost categories.	11
Table 4: Recommended items for a resource use measure instrument.	15
Table 5: Study characteristics of cost of illness studies conducted in Germany since 2010.	16
Table 6: Comparison of annual resource costs of studies conducted in Germany since 2010.	17
Table 7: Clinical and patient reported data obtained during the studies.	22
Table 8: Core Items of the MS-HRS.	28
Table 9: Valuation of resource costs applied in the MS-HRS (price year 2011).	30
Table 10: Defined Daily Dose net costs in the respective years for DMTs multiplied with 365.	31
Table 11: Validity and reliability measurements assessed for the MS-HRS.	33
Table 12: Baseline characteristics of the study population included in the validation process (N=2207).	39
Table 13: Comparison of baseline characteristics of included and excluded patients for the validation process.	40
Table 14: Direct (non-)medical and indirect costs (excl. DMTs) stratified by EDSS scores.	41
Table 15: Correlation coefficients for convergent and discriminant validity.	41
Table 16: Health resource utilization in persons with MS per quarter (N=2207).	42
Table 17: Baseline characteristics of the study population for the analysis of associations between disability worsening and societal economic costs (N=1959).	44
Table 18: Mean±SD quarterly cost in Euro (€) within two years of study time (N=1959).	46
Table 19: Comparison of baseline characteristics of included and excluded patients for the relapse costs analyses.	49
Table 20: Characteristics of the study population for the analysis of relapse associated societal economic costs (N=1882).	50
Table 21: Total, direct and indirect relapse costs (Δ) stratified by patient and disease characteristics.	54
Table 22: Detailed breakdown of relapse costs (Δ) stratified by patient and disease characteristics.	56
Table 23: Sensitivity analysis for relapse costs (Δ) stratified by patient and disease characteristics.	57

Abbreviations

APCQ	Annotated Patient Cost Questionnaire
ARR	Annualized relapse rate
AVO	Avonex®
BeT	Betaferon®
BMI	Body mass index
BRACE	Beta interferons and glatiramer acetate
CDW	Confirmed disability worsening
CDW-6	Disability worsening with confirmation after six months
CNS	central nervous system
COI	Cost of illness
COP	Copaxone®
CRF	Case report form
CSRI	Client Service Receipt Inventory
DAF	Disease activity free
DIRUM	Database of Instruments for Resource Use Measurement
DMT	disease-modifying therapies
DSS	disability status scale
EDSS	Expanded disability status scale
EMA	European Medicines Agency
EQ-5D	EuroQol-5 Dimensions
EQ-5D-3L	EuroQol-5 Dimensions 3 level version
EXT	Extavia®
FINGO	Fingolimod
GA	Glatiramer acetate
GIL	Gilenya®
GLM	generalized linear model
GLMM	generalized linear mixed model
HRQOL	Health-related quality of life
ICC	intraclass correlation coefficient
IFN- β	interferon beta
IFN- β -1a	interferon beta-1a
IFN- β -1b	interferon beta-1b
IQR	interquartile range
IQWiG	Institute for Quality and Efficiency in Health Care
IRR	Incidence rate ratios
MRI	Magnetic resonance imaging
MS	Multiple sclerosis
MSDS3D	Multiple sclerosis documentation system 3D
MS-HRS	Multiple Sclerosis Health Resource Utilization Survey
OTC	Over-the-counter
PANGAEA	Post-authorization non-interventional German safety study of Gilenya®
PEARL	Prospective Pharmacoeconomic Cohort Evaluation
PIRA	progression independent of relapse activity
PP	primary progressive
PR	Progressive relapsing
PRIMUS-A	Patient Reported Indices for Multiple Sclerosis-Activity limitations
PRIMUS-L	Patient Reported Indices for Multiple Sclerosis-Health-related quality of life
PRO	Patient-reported outcomes

PS	Propensity score
RAW	Relapse-associated worsening
REB	Rebif®
RR	Relapsing-remitting
RRMS	Relapsing-remitting multiple sclerosis
RWW	Relapse without worsening
RWW	Relapse without worsening
SD	Standard deviation
SHI	Statutory health insurance
SP	Secondary progressive
SPMS	secondary progressive multiple sclerosis
SPSS	Statistical Pack-age for Social Science
TSQM	Treatment Satisfaction Question-naire for Medication
TYS	Tysabri®
UKNDS	UK Neurological Disability Scale
WPAI	Work Productivity and Activity Impairment Question-naire
WPAI:GH	Work Productivity and Activity Impairment Question-naire: General Health
WPAI:SHP	Work Productivity and Activity Impairment Question-naire: General Health: Specific Health Problem

1 Introduction

1.1 Multiple sclerosis

1.1.1 Pathogenesis

Multiple sclerosis (MS) is a chronic disease characterized by inflammatory demyelination and neurodegeneration occurring either focally or diffusely throughout the white and grey matter in the central nervous system (CNS) (Lassmann et al., 2012). MS is presumed to be of autoimmune origin, however the exact pathogenesis is not fully understood (Breuer et al., 2019). It is considered that genetic and environmental factors contribute to an increased risk of developing MS (Ascherio and Munger, 2016). Those factors with most supporting evidence are the Epstein-Barr virus, smoking cigarettes and vitamin D deficiency (Amato et al., 2017; O'Gorman et al., 2012; Pugliatti et al., 2008). Furthermore, there is a geographic variation in the distribution of MS. It is more common in regions with temperate climate than in those with tropical climate (Simpson et al., 2011; Wallin et al., 2019).

1.1.2 Clinical manifestation

MS can cause a variety of neurological symptoms, since any region of the CNS can be affected by the disease (Ghasemi et al., 2017). However, in the course of the disease, patterns of abnormalities emerge: MS commonly manifests with sensory, motor, visual or cerebellar dysfunctions as well as fatigue (MSIF, 2013). In later stages of the disease patients may also be affected by bowel, bladder and sexual dysfunction (Stuke et al., 2008). Table 1 lists the frequency of common symptoms by disease duration.

Symptoms	Disease duration <2 years (%)	Disease duration > 15 years (%)
Neuropsychological		
Fatigue	40.6	67.6
Cognitive dysfunction	19.6	40.6
Depression	23.6	38.0
Motoric dysfunction		
Spasticity	17.3	75.8
Ataxia/ tremor	24.4	56.5
Others		
Pain	24.6	42.2
Bladder dysfunction	20.5	74.0
Bowel dysfunction	4.8	31.1
Sexual dysfunction	8.0	27.3
Others	2.1	3.9

Table 1: Frequency of symptoms in patients with MS by disease duration. Reproduced with modifications from Stuke et al. (2008).

1.1.3 Descriptive epidemiology

Approximately 2.2 million people worldwide are affected by MS. The risk of developing the disease is more than two times higher in women than in men (Compston and Coles, 2008; Wallin et al., 2019). So far, nationwide population-based data on the prevalence (number of patients alive at a specific time per 100,000 person) of MS are not available in Germany and estimates are based on extrapolations from regional studies. Prevalence was estimated at 289 cases per 100,000 insured individuals in 2010, with 70% of those affected being female according to an analysis of statutory health insurance (SHI) data (Petersen et al., 2014). Globally, the prevalence of MS increases which might be explained by a decrease in disease-related mortality (Wallin et al., 2019).

The incidence rate (number of new cases per 100,000 persons in one year) increases over time partially due to improvements in diagnostic techniques and a heightened awareness of the disease (Compston and Coles, 2008). The incidence rate is estimated at 8.0 cases per 100,000 person annually in Germany (Fasbender and Kölmel, 2008). Disease onset is mostly reported between 20-40 years of age and peaks at 34.4 years. Although the accumulation of disability is not the primary cause of death, life expectancy is on average 6-12 years shorter compared to the general population (Grytten Torkildsen et al., 2008; Kingwell et al., 2012; Ragonese et al., 2008).

1.1.4 Clinical hallmarks and outcome measures

1.1.4.1 Relapses

Relapses and disability worsening are the clinical hallmarks of MS (Giovannoni et al., 2016). A relapse is defined as a worsening or appearance of new neurological symptoms during acute or subacute disease activity in the absence of fever or infection lasting more than 24 hours followed by a period of partial or complete recovery (Lublin et al., 2014; McDonald et al., 2001). Relapses have a substantial impact on a variety of functioning on physical and mental health scales (Healy et al., 2012). Prevention of relapses is a main treatment target of disease modifying therapies (DMTs). In clinical trials, relapse-based outcomes e.g. annualized relapse rate (ARR, defined as the number of confirmed relapses per patient adjusted for duration of follow-up) are a sound reflection of short term efficacy (van Munster and Uitdehaag, 2017). Further relapse-based outcomes are the average number of relapses per patient, proportion of (non-) relapsing patients and the cumulative probability of (non-) relapse events.

1.1.4.2 Disability worsening

Disability worsening (confirmed disability worsening, CDW) refers to an accumulation of MS-related functional disability sustained over a predefined period usually three or six months (Kalincik et al., 2015; Uitdehaag, 2018; Wiendl and Meuth, 2015). It is objectified as changes in disability, most commonly defined as: ≥ 1.5 -point increase for a reference-score of 0; ≥ 1 -point increase for a reference-score of 1.0-5.0; and ≥ 0.5 point increase for a reference score greater than 5.5 and as measured on the Expanded Disability Status Scale (EDSS) (Wiendl and Meuth, 2015). CDW can be either the result of relapse associated worsening (RAW) or progression independent of relapse activity (PIRA) (Kappos et al., 2018b). While RAW can result from poor recovery from multiple or severe relapse activities, PIRA may indicate a progressive clinical course characterized by neurodegenerative processes independent of inflammation (Kappos et al., 2018a; Lublin et al., 2014).

The disability status scale (DSS) was published in the 1950s and subsequently modified until it was reissued in 1983 as EDSS (Kurtzke, 1955; Kurtzke, 1983). The EDSS is the most commonly used measure of neurological disability in MS clinical trials. Disability is assessed by capturing limitations in functional systems including vision, bowel, bladder, brainstem, the pyramidal system, the cerebellar system, the cerebral and the sensory system. In addition, it covers ambulatory endurance and the ability to perform activities of daily living. The scale ranges from 0 (normal neurological examination) to 10 (death due to MS) with half point steps between 1 and 10. The calculation of scores is subject to criticism (Cohen et al., 2012). While scores of 0-4.0 measure impairments in functional systems, scores above 4.0 are reflecting disability (van Munster and Uitdehaag, 2017). Thereby, ambulatory endurance and the need of assistive devices primarily determine scores between 4.0-7.0 and scores between 7.0-9.5 are based on the ability to carry out activities of daily living (van Munster and Uitdehaag, 2017). Further shortcomings are the non-linearity leading to a bimodal distribution and the limited ability to capture cognitive impairments. Figure 1 gives an overview of characteristics and caveats of the EDSS (Hyland and Rudick, 2011; Twork et al., 2010; Vickrey et al., 1995).

Figure 1: Characteristics and limitations of the Expanded Disability Status Scale. Reproduced with modifications from van Munster and Uitdehaag (2017).

1.1.5 Clinical course

Four clinical courses of MS can be distinguished and are visualized in figure 2: Relapsing-remitting (RR), secondary progressive (SP), primary progressive (PP) and progressive relapsing (PR) (Lublin et al., 2014). The majority of patients (85%-90%) initially experience a RR course characterized by episodic symptoms (relapses) with full or partially recovery (remitting) (MSIF, 2013). Up to 80% of patients transit to the SP course within 20 years marked by continuous neurological decline with or without relapses (MSIF, 2013). So far, no precise clinical, imaging, immunological or pathological criteria exist to define the transition point from RRMS to SPMS (Lublin et al., 2014). Approximately 10-15% of people with MS exhibit an initial steady progressive deterioration in neurological functions with relapses (PR) or without relapses (PP) (Lublin et al., 2014; Miller and Leary, 2007). The pathological features of inflammation and neurodegeneration are present in all clinical subtypes but vary qualitatively and quantitatively over time (Grigoriadis and van Pesch, 2015; Lassmann et al., 2012). While inflammation is more distinct in the acute relapsing phases of the disease, the key mechanism of disease progression might be a neurodegenerative process that is independent of inflammation (Correale et al., 2017; Dendrou et al., 2015; Meuth et al., 2008).

Figure 2: Classification of the clinical courses of multiple sclerosis. Reproduced with modifications from Grigoriadis and van Pesch (2015).

1.1.6 Disease modifying therapies

There is no curative treatment for MS available. Treatments aim at reducing the risk of relapses and slowing disability worsening (Montalban et al., 2018). Since the approval of interferon in 1993, neurologists and patients have seen a variety of therapeutic options including injectable and oral formulations approved by the European Medicines Agency (EMA) for the treatment of MS (De Angelis et al., 2018). First line DMTs for low and moderate active RRMS include interferon beta-1b (IFN- β -1b), interferon beta-1a (IFN- β -1a), glatiramer acetate (GA), teriflunomide and dimethyl fumarate while fingolimod is approved for treatment of patients who are insufficiently treated with first line therapies. Treatment options for patients with highly active MS include apart from first line treatments fingolimod, natalizumab, alemtuzumab, cladribine and ocrelizumab (Rommer et al., 2019). In the following analyses, patients were treated either with IFN- β or with GA. These medications are introduced shortly.

The mechanisms of action of IFN- β and GA are multifactorial and not fully understood. It is evident that broad immunomodulatory effects are generated on multiple levels of the immune response (Aharoni, 2013; Kieseier, 2011). The safety and efficacy profile of IFN- β and GA including clinical and magnetic resonance imaging (MRI) measures show only small differences (La Mantia et al., 2016). Both treatments are associated with a $\approx 30\%$ reduction in risk of confirmed relapses compared to placebo (Khan et al., 2013; Killestein and Polman, 2011). The efficacy of IFN- β in reducing confirmed disability worsening is divergent (Kappos et al., 2004). An European study showed that IFN- β significantly slowed confirmed disability worsening while an American study could not confirm this effect (Kappos and European Study Group on Interferon β -1b in Secondary Progressive MS, 1998; The North American Study Group on Interferon beta-1b in Secondary

Progressive MS, 2004). In summary, response to IFN- β and GA is highly heterogeneous and in some patients, disease activity is insufficiently controlled (Ziemssen et al., 2015a).

Fingolimod, the first available oral agent, is a sphingosine 1-phosphate receptor modulator that prevents the egress of autoreactive lymphocytes from lymph nodes, leading to reduced infiltration of potentially autoaggressive lymphocytes into the CNS (Chun and Hartung, 2010). Fingolimod reduced the annualized relapse rate by 48-60% compared to placebo and by 38-52% compared to INF- β 1a (Calabresi et al., 2014; Cohen et al., 2010; Kappos et al., 2010). A significant effect on confirmed disability worsening after three and six months was observed in one out of three clinical phase II trials (Calabresi et al., 2014; Cohen et al., 2010; Kappos et al., 2010). In RWE, analyses it was further confirmed that switching therapy from IFN- β and GA to fingolimod rather than to another injectable is beneficial in terms of fewer relapses, more favorable disability outcomes and greater treatment persistence (Alsop et al., 2017; He et al., 2015).

1.1.7 Natural history

The natural history of MS is determined by the frequency of relapses as well as disability worsening. The average relapse rate is age- and time-dependent. It is higher in young adulthood and decreases significantly in the course of the disease from 1.85 ARR in the first year after onset to 0.2 ARR after 19 years (Patzold and Pocklington, 1982). In RRMS the ARR is considered to be 0.5 on average (Confavreux et al., 1980). After 18 years about 50% of the patients have reached a disability level, so they need a walking aid to walk 100m (EDSS 6.0) (Scalfari et al., 2010). However, more recent investigations have shown that disability worsening accrued more slowly than previously reported with median time to EDSS 6.0 of 28 years (Tremlett et al., 2006).

1.2 Cost of illness research

Cost of illness (Col) refers to the economic framework that is used to estimate the costs of a particular disease to society (Drummond, 2015; Rice, 1967). Col studies offer valuable insights for improving societal welfare in different manners. First, they can help to detect the drivers of diagnosis and treatment costs of a specific disease (Simoens, 2009). Second, Col studies may help ranking diseases according to their burden to society and hence to determine priorities in medical research (Drummond, 1992; Tarricone, 2006). They are particularly valuable in addition to epidemiological studies. For example, morbidity and mortality are essential factors to estimate the burden of disease but offer limited insights on the consequences of impaired health (WHO, 2009). Col studies capture the economic consequences like productivity losses or escalation in out of pocket payments that are potentially fatal in chronic, progressive diseases like MS. Third, data from Col studies can be used in full economic evaluation in which both costs and consequences of two or more interventions are evaluated (Simoens, 2009). The value of Col research is reflected in its frequent use of policy makers and institutions like the World Health Organization (WHO).

Chapters 1.2.1-1.2.3 provide a methodological review of Col studies. The application of the methods is subsequently described in the empirical part of the thesis.

1.2.1 Classification of costs

Col studies stratify costs into direct medical, direct non-medical and indirect costs. Direct medical costs refer to resource utilization resulting directly from health care interventions, for example physician consultations, in- and outpatient stays and medication (Jo, 2014). Direct non-medical costs arise from resource utilization with a supportive character for the health system such as transportation to the hospital, informal care from family members and friends or reconstruction in a patients home due to disability caused by the illness (Jo, 2014). Indirect costs refer to productivity loss due to a disease (Segel, 2006). They result from not showing up for work (absenteeism) or from reduced productivity at work (presenteeism). Indirect costs are particularly high in chronic illnesses like MS. An overview of exemplary resource items is shown in table 2.

Direct medical resources		Direct non-medical resources	Indirect resources
Inpatient care Outpatient care Home health care Physician visits at Specialists, General practitioner Ancillary services Nurses Physical therapists Complementary medicine	(Non-)Prescription medication Imaging methods Laboratory tests Rehabilitation Trainings/ education	Social services Childcare Housekeeping Investments Modifications at house/ car Time (searching, traveling, waiting etc.)	Sick leave Disability pension Presenteeism

Table 2: Exemplary resource items by cost categories.
Reproduced with modifications from Jo (2014).

So-called intangible costs describe the burden resulting from pain and suffering measured by reduction in quality of life. Intangible costs are not economic costs and usually not monetized (Larg and Moss, 2011). According to the Institute for Quality and Efficiency in Health Care (IQWiG) intangible costs should be reported as a negative benefit on the outcome side in health economic assessments (IQWiG, 2017).

1.2.2 Costing methodologies

Costing methodologies can be described by the underlying process of resource identification (microcosting- vs. gross costing) and resource valuation (top-down vs. bottom-up) (Luce et al., 1996; Tan et al., 2009).

Microcosting involves the “direct enumeration and costing out of every input consumed in the treatment of a particular patient” (Luce et al., 1996). For example, in hospital services laboratory tests, medical consumables and surgical interventions might be identified (Ljungqvist et al., 2017). In contrast, gross costing identifies only a limited number of cost components. For example, inpatient stays might be used as a surrogate for hospital services. Both, micro- and gross costing can be applied as top-down or bottom-up approaches.

The top-down approach calculates average service or unit costs by dividing total costs by the number of cost drivers (e.g. number of patients) (Raulinajtys-Grzybek and Świdarska, 2015). The bottom-up approach values each service provided with the resources used, resulting in patient specific service or unit costs (Cunnam et al., 2016; Raulinajtys-Grzybek and Świdarska, 2015). Combining valuation and identification meth-

odologies gives four costing methodologies as shown in figure 3: Bottom-up microcosting, top-down microcosting, bottom-up gross costing and top-down gross costing (Tan et al., 2009).

		Identification of resources	
		Low accuracy	High accuracy
Valuation of resources	Low accuracy	Top-down gross costing	Top-down microcosting
	High accuracy	Bottom-up gross costing	Bottom-up microcosting

Figure 3: Overview of costing methodologies.
Reproduced with modifications from Tan, et al. (2009).

Bottom-up microcosting is believed to be the gold standard since costs are assigned to individual services at a level that reflects the patients' actual consumption of resources (Raulinajtys-Grzybek and Świdarska, 2015; Ruger and Reiff, 2016). This methodology enables statistical analyses to detect differences in costs between subgroups as well as sensitivity analyses to handle uncertainty (Smith and Rudmik, 2013). On the downside, its feasibility is challenging since bottom-up microcosting is time and cost consuming (Tan et al., 2009).

A frequently used alternative is the top-down microcosting approach that assigns total costs to a detailed enumeration of services provided. It is more feasible compared to bottom-up microcosting (Tan, et al., 2009). Nevertheless, statistical analysis of patient- or disease strata are more restricted.

In bottom-up gross costing resources are valued highly accurate but identified at an aggregated level. At the least accurate level, top-down gross costing assigns aggregated values to average patients (Tan et al., 2009).

Gross costing approaches have the disadvantage that the less resource items are identified, the more equal patients are handled even though they obviously have dissimilar costs (Tan, 2009). In complex diseases like MS, patients' needs cover a wide range of costly resources and a gross costing approach is inadequate. Due to the aforementioned advantages, bottom-up microcosting is used in the following study.

1.2.3 Process of cost estimation

The process of cost estimation includes the identification of resource consumption linked with a particular disease, the quantification of these resources as well as the valuation with corresponding prices weights to obtain costs.

1.2.3.1 Identification of resource use - relevance of the perspective

Identification of resources is highly dependent on the study perspective (Segel, 2006). Commonly studies are conducted from the perspective of the society, SHI or SHI plus patients (Table 3) (IQWiG, 2017).

The societal perspective is the broadest encompassing all direct (non-)medical and indirect costs regardless who is bearing them. Generally, the societal perspective is preferred by economists since it minimizes potential biases of narrower views (Dagenais et al., 2008; Graf von der Schulenburg et al., 2007; van Lier et al., 2018). For example from a societal perspective, transfer payments (e.g. sickness benefits) are not considered as resources. They are only shifted from one group (workers) to another (workers on sick leave) without changing the value of resources within the society (Russell et al., 1996).

The SHI perspective considers all costs that are refunded by the SHI. Included are transfer payments (e.g. sickness benefits) which represents a loss of resources from the viewpoint of the SHI. The SHI plus patients-perspective additionally includes patients' own expenses but transfer payments are not considered.

Various sources can be used to identify cost-causing events. During the cost identification process, it is recommended to use a decision tree showing the pathways of all events of a patient's journey. Previous studies, pretrial data collections, pilot tests, modellings, expert opinion and guidelines can be used to build the decision tree (Johnston et al., 1999).

	Direct medical costs		Direct non-medical costs		Indirect costs	Transfer costs
	Refund-able	Not refund-able	Refund-able	Not refund-able	-	-
Society	✓	✓	✓	✓	✓	X
SHI plus patients	✓	✓	✓	✓	X	X
SHI	X	X	✓	X	X	✓

Table 3: Perspective of cost estimation and relevant cost categories.
Abbreviations: statutory health insurance (SHI).
Reproduced with modifications from IQWiG (2017).

1.2.3.2 Quantification resource consumption

Resource use data are often taken secondarily from sources like administrative datasets or they are collected primarily alongside clinical trials. Administrative data are mostly standardized, reliable and accessible and therefore superior to primary data in specific cases (Franklin and Thorn, 2019). However, data accuracy might be limiting, as several healthcare services can – but should not – be combined under one caption. Furthermore, administrative data underlie the limitations that no societal perspective can be adopted since not all relevant resources are recorded (e.g. informal care or presenteeism) (Martin et al., 2012). Furthermore, administrative data do not contain clinical data, making it difficult to identify reasons for cost differences in subpopulations.

Diaries and questionnaires are frequently used to gather data primary alongside clinical trials (O'Sullivan et al., 2005). Patient diaries are completed after services have been received or on regular sequence of short time intervals (e.g. per day). Therefore, diaries are limiting the risk of recall biases which might occur in questionnaires (Ridyard et al., 2015). Although, resource use diaries are well suited for cross-sectional and short-term data collections, considerable problems regarding dropout rates and thoroughness in reporting are challenging for data collections expanding twelve months (Fowler and Floyd, 1993). Therefore, questionnaires might better suit health economic data collection with time periods longer than one year (Schweikert et al., 2008).

Self-reported questionnaires can be distinguished in disease-specific and generic instruments (Leggett et al., 2016). While disease-specific instruments are adapted to the specific needs of a patient population, generic instruments target more broad population. Especially in complex conditions like MS, disease specific measures are necessary to collect relevant resources comprehensively.

To avoid issues of recall bias in self-reported health resource surveys recall periods should be determined based on cognitive theories. Recall biases have two main causes: First, more common and less striking events are less likely to be recalled precisely over a long period. Second, long recall periods cause telescoping effects in which events outside the targeted period are incorrectly assigned in the recall period.

1.2.3.3 Valuation of resource consumption

Resource valuation is the assignment of an economic price to a resource item. In studies conducted from a societal perspective resource prices should reflect the societal opportunity costs. These are defined as the value of the lost benefits, as resources are not available for its best alternative use (van Lier et al., 2018).

Valuation of direct medical costs

Societal opportunity costs and market prices are equal on perfectly competitive markets. Nevertheless, the healthcare market is imperfect for the sake of statutory regulations. Therefore, societal opportunity costs have to be approximated in most cases using e.g. published data for standardized cost units (Bock et al., 2015). For medical procedures (examinations, lab tests and diagnostic imaging) it is recommended to use bargaining prices e.g. the physicians' Fee Schedule that is negotiated between the National Association of Statutory Health Insurance Physicians and the Central Federal Association of Health Insurance Funds (IQWiG, 2008).

When medication costs are a large proportion of the direct costs, it is highly recommended to use an accurate medication cost method (Heslin et al., 2018; Kobelt et al., 2017b). A recommended approach would be to (i) calculate costs per milligram for each medication on the basis of most cost-efficient pack sizes and (ii) apply these unit costs to the reported doses (Heslin et al., 2018).

Valuation of informal care costs

In contrast to formal care, informal care is provided by family members, friends and neighbors (Oliva-Moreno et al., 2019). Informal care is especially relevant in diseases that cause significant restrictions on a person's autonomy and therefore a distinctive demand for care (Jimenez-Martin and Prieto, 2012). Even though informal care is often provided without monetary compensation there are hidden costs for the caregivers such as a decrease in labor supply (Oliva-Moreno et al., 2017). The two mainly used methods to value informal care costs in CoI studies are the opportunity costs method (69% of cases) and the proxy good method (27% of cases) also known as replacement cost approach (Oliva-Moreno et al., 2017). In the former, informal care time is valued at the

opportunity costs out of the perspective of the caregiver (Landfeldt et al., 2019). The opportunity costs are mostly based on average wages or derived from estimating techniques like conjoint analysis or contingent valuation (Landfeldt et al., 2019). The proxy good method values informal care activities at a shadow market price like a housekeeper for housekeeping service. It is frequently debated which of the two methods should be used (Pike and Grosse, 2018). For analysis from a societal perspective the opportunity cost method is the recommended technique as these are an adequate approximation (Landfeldt et al., 2019).

Valuation of indirect costs

The two main approaches to estimate indirect costs are the friction cost method and the human capital approach. In the latter lost productivity is valued at the opportunity cost of labor using the cost of employment e.g. the national mean wage plus employer's cost and social fees (Landfeldt et al., 2019). Using the friction cost method productivity losses only occur during the friction period, the time it takes to replace a worker taken from the pool of unemployed (Landfeldt et al., 2019). While the friction cost method underlies the highly unrealistic assumption that labor supply is fully elastic, the human capital approach is criticized for not taking into account the possibility of replacement even if the unemployment rate is significantly high (Jo, 2014; Pike and Grosse, 2018). A recent review revealed that more than 90% of CoI-studies evaluated productivity loss using the human capital approach which is also in line with most countries health technology assessment recommendations (Pike and Grosse, 2018).

Prices from different periods should be adjusted using country specific discount rates and the formula:

$$Discount\ factor = \frac{Price\ index_{base\ year}}{Price\ index_{data\ year}} \quad (I)$$

1.3 Cost of multiple sclerosis

The economic burden associated with MS is tremendous for patients, caregivers, healthcare providers and society. The chronic and mostly progressive nature of MS leads to long-term treatment with expensive DMTs. Continuous monitoring of clinical and MRI activity as well as further examinations are needed to determine response to DMTs (Gasperini et al., 2019). Additional management of new or worsening symptoms might require interdisciplinary collaboration of a wide range of specialists including neurologists, physiotherapists or occupational therapists. In later stages of the disease, productivity restrictions and informal care lead to further demand for societal resources.

The following chapters summarize the state of research on the cost of MS. Chapter 1.3.1 reviews instruments used to date to assess the economic burden. Chapter 1.3.2 reviews relevant Col studies with a focus on disease worsening and relapses burden.

1.3.1 Instruments for resource use measurement

Despite the long tradition of health economic analysis in MS, there is no validated instrument available to measure resource costs associated with MS (D'Amico et al., 2019; DIRUM, 2019; Leggett et al., 2016). Even though Kobelt et al. (2019b) recently recognized the need for an instrument for resource use measurement in MS, validating and publishing an instrument still remains a task to fulfil. Considerable research time is repeatedly spent to develop study-specific instruments. Nevertheless, the psychometric properties of these instruments are scarcely analyzed or re-validated if used in a modified manner (Franklin and Thorn, 2019; Ridyard and Hughes, 2010). To address this issue, the DIRUM database was established to provide an open-access database of instruments for resource use measurement (DIRUM, 2019).

Besides disease-specific instruments, universal instruments exist (McCrone et al., 2008). The Client Service Receipt Inventory (CSRI) might be the most widely spread universal instrument with over 200 modifications for different disease areas, healthcare systems, languages and types of applications (Beecham and Knapp, 1992; Thorn et al., 2013). The original questionnaire gathers data on the individual background, accommodation and living situation, employment, social security benefits, healthcare services and informal care (Beecham and Knapp, 1992; Thorn et al., 2013). Psychometric properties have been assessed in different settings with overall good results (Byford et al., 2007; Leggett et al., 2016; Roick et al., 2001). Furthermore, the Annotated Patient Cost Questionnaire (APCQ) is a universal instrument (Thompson and Wordsworth, 2001). It comprises a collection of questions to simplify the transfer of the APCQ into different contexts (Thorn et al., 2013). Even though the questionnaire might perform well, it is not widely adopted

yet (Thorn et al., 2018). However, the usefulness of both of the aforementioned questionnaires is limited in complex diseases like MS because MS patients' needs extend the resources that are captured in those instruments.

Standardized instruments additionally exist that concentrate on specific cost components (Landfeldt et al., 2019; Reilly et al., 1993). For example the Work Productivity and Activity Impairment Questionnaire (WPAI) is a widely spread instrument to assess absenteeism, presenteeism, overall work performance and non-work activities for general health (WPAI:GH) or specific health problems (WPAI:SHP) including MS (WPAI:MS) (Chen et al., 2018). The psychometric properties of the WPAI have been analyzed in several chronic diseases with overall good results (Lambert et al., 2014; Reilly et al., 2004; Reilly et al., 2010).

Recently, a list of core items was published by an expert consensus that should be included in instruments for resource use measurement from the provider perspective (Thorn et al., 2018). The list gives a helpful orientation for the development of a new instrument (Table 4). However, it does not take into account disease specific resources.

Type of care	Item
Hospital care	Number of hospital admissions (inpatient stays/ day case)
	Length of stay (e.g. dates/ number of nights)
	Number of hospital outpatient appointments
Emergency Care	Number of visits to A&E
	Number of hospital visits after A&E
GP surgery/ health clinic/ community setting	Number of appointments
	Type of professional
Home Care	Number of visits
	Type of professionals
Medication	Name/class of medication

Table 4: Recommended items for a resource use measure instrument.
Abbreviations: accident and emergency (A&E); general practitioner (GP).
Reproduced with modifications from Thorn et al. (2018).

1.3.2 Review of cost of multiple sclerosis studies in Germany

Three academically highly relevant Col studies analyzed the societal economic costs of MS comprehensively in Germany since 2010 (Table 5) (Flachenecker et al., 2017; Karampampa et al., 2012b; Reese et al., 2011). I do not consider studies published prior to 2010 due to the major changes in therapy and diagnostic options in recent years.

	Flachenecker 2017	Karampampa 2012	Reese 2011
Sample size (n)	5474	244	144
Year of price level	2015	2009	2006
Age (mean±SD)	51.8±11.0	41.2±9.9	41.7±11.3
Female gender (%)	74%	69%	69%
EDSS (mean±SD)	4.0±2.5	1.8±1.8	3.5±2.0
Phenotype of MS (%)	RRMS: 46% SPMS: 28% PPMS: 17% UNKNOWN: 3%	RRMS: 65% SPMS: 13% PPMS: 8% UNKNOWN: 14%	RRMS: 67% SPMS: 28% PPMS: 5%

Table 5: Study characteristics of cost of illness studies conducted in Germany since 2010.

Abbreviations: Expanded Disability Status Scale (EDSS).

Reproduced with modifications from Flachenecker et al. (2017); Karampampa et al. (2012b); Reese et al. (2011).

The most recent and largest study (n=5457) was conducted by Flachenecker et al. and reported the German results of a Col study rolled out throughout Europe. In comparison to Karampampa (2012) and Reese (2011) participants were older (51.8±11.0 vs. 41.2±9.9 vs. 41.7±11.3) and more severely disabled (EDSS 4.0±2.5 vs. 1.8±1.8 vs. 3.5±2.9). Karampampa et al. (2012b) provided results for Germany of a European study with the sole characteristic that costs were also reported depending on the clinical course (RRMS, SPMS, PPMS). Reese et al. (2011) recruited 144 patients for their study. Contrary to the questionnaire-based methodology of Flachenecker and Karampampa, data were collected in a structured face-to-face interview. Table 6 summarizes the results of these studies.

	Flachenecker 2017 ^{2, 4} annual costs in €	Karampampa 2012 ^{1,2, 5} annual costs in €	Reese 2011 ^{1,3} annual costs in €
Direct medical	18,958-16,521	18,395-32,928	24,037
Inpatient	1236-6783	674-13,406	4833
Outpatient	126-231	515-3875	
Physician consultations	1066-2317	589-2127	2211
Investigations	324-201	107-68	--
DMTs	15,819-4891	15,204-11,900	16,486
Medication	387-2098	1305-1552	507
Direct non-medical	952-18,717	1262-23,198	877
Informal care	721-15,280	901-18,360	--
Investments	231-3,437	361-4838	877
Indirect	8190-21,586	3317-11,931	21,538
Sick leave	723-108	1605-994	2037
Presenteeism	--	--	667
Disability pension, early retirement	7467-21,478	1712-10,937	18,834

Table 6: Comparison of annual resource costs of studies conducted in Germany since 2010.

¹ Cost data are inflated to index year 2015 using the consumer price index and the formula stated in chapter 1.2.3.3. Discount rates are 8.5% and 14.2% for prices from 2006 and 2009, respectively.

² Costs depend on disease severity (EDSS 0-3 to EDSS 7-9).

³ Contrary to the original categorization, investments were allocated to direct non-medical costs instead to direct medical costs.

⁴ To ensure comparability, community services were not considered in direct non-medical costs

⁵ To ensure comparability, professional care was not considered in direct medical costs.

Reproduced with modifications from Flachenecker et al. (2017); Karampampa et al. (2012b); Reese et al. (2011).

Direct medical costs

Considering the differing methodologies, study design as well as patient and disease characteristics, the three Col studies are difficult to compare. However, tendencies can be deduced with caution. In the following paragraphs, the results of the above-mentioned studies are discussed.

Annual direct medical costs ranged between 16,521€-32,928€ depending on disease severity as measured by EDSS. In mildly disabled patients, DMTs represented the largest share of direct medical costs (15,204€-16,486€). In severely disabled patients, the share of DMTs decreased (4891€-11,900€) partially explained by the lack of treatment options for a progressive clinical course in later stages of the disease.

Costs of inpatient hospitalization accounted for the second largest share of annual direct medical costs. They were mainly incurred in neurological wards (Flachenecker et al., 2017). The economic burden rose significantly with increasing disability (1236€ (EDSS 0-3)-13,406€ (EDSS 7-9)). Nevertheless, annual costs in the same EDSS categories varied widely (674€-1236€ (EDSS 0-3); 6783€-13,406€ (EDSS 7-9)), possibly due to different study methodologies.

Annual costs for outpatient physician consultations ranged between 1066€-2317€ depending on the degree of disability. The most frequent consultations were with neurologists, physiotherapists and general practitioners (Flachenecker et al., 2017).

Besides DMTs, symptomatic treatments contributed mainly to medication costs. Annual costs ranged with EDSS from 387€-1305€ (EDSS 0-3) to 1552€-2098€ (EDSS 7-9). Diagnostic services accounted for a small share of direct medical costs and ranged between 68€ (EDSS 0-3) and 324€ (EDSS 7-9).

Direct non-medical

Annual direct non-medical costs ranged between 952€-1262€ (EDSS 0-3) and 18,717€-23,198€ (EDSS 7-9). Severely disabled patients were more dependent on informal care, which resulted in a large cost difference compared to mildly disabled patients (721€-901€ (EDSS 0-3)-15,280€-18,360€ (EDSS 7-9)). Annual costs for investments (e.g. walking and mobility aids) and house or car modifications rose with increasing EDSS from 231€-361€ (EDSS 0-3) up to 3437€-4838€ (EDSS 7-9) (Flachenecker et al., 2017).

Indirect costs

The unfavorable diagnostic age caused annual indirect costs ranging between 3317€-8190€ (EDSS 0-3) and 11,931€-21,586€ (EDSS 7-9). Increasing disability had two opposite effects: While annual costs for absenteeism decreased from 723€-1605€ (EDSS 0-3) to 108€-994€ (EDSS 7-9), costs for disability pension and early retirement increased from 1712€-7467€ (EDSS 0-3) to 10,937€-21,478€ (EDSS 7-9). Increases in indirect costs were multifactorial: In the short term, fatigue and reduced mobility were associated with an increased number of sick leave (Wickstrom et al., 2014). In the long term, female gender, ≥ 51 years of age, moderate/ high disease severity and impaired manual dexterity were associated with higher rates of disability pension (Chruzander et al., 2016). Reese et al. (2011) further reported annually costs of presenteeism of 667€ per patient regardless of disease severity.

Relapse associated costs

The economic burden associated with relapses is significant (Naci et al., 2010). Almost all societal resources are increasingly used. The cost drivers during relapses are hospitalizations, physician consultations, informal care and sick leave (Hawton and Green, 2016; Kobelt et al., 2017b). However, relapse costs can vary with relapse intensity ranging from a single symptom lasting over a few hours up to a wide range of neurological dysfunctions remaining over weeks.

O'Brien et al. (2003) analyzed the direct medical costs associated with relapses stratified by relapse intensity. They reported relapse associated direct costs of 243 US\$, 1847US\$, and 12,870US\$ for mild, moderate and severe relapses, respectively. O'Brien et al. (2003) concluded that costs are more than 6-fold higher if relapses are treated in inpatient settings.

Flachenecker et al. (2017) and Karampampa et al. (2012b) calculated relapse costs as the difference between quarterly costs of patients with and without relapses with results of 2468€ (price level 2015) and 5249€ (price level 2009), respectively. Patient and disease characteristics of these distinct populations were not presented. However, differences in relapse costs between subgroups are likely since relapses and the associated costs might vary across and within individuals with respect to disease severity, disease duration, gender or therapy (Derfuss et al., 2016; Hawton and Green, 2016; Kobelt et al., 2006b; Turner et al., 2019).

1.4 Hypotheses and objective

MS is an incurable and progressive condition leading to tremendous suffering for patients and an enormous societal economic burden (Kobelt et al., 2017b; Wallin et al., 2019). The key clinical hallmarks of MS are relapses and disability worsening both associated with significantly increased utilization of direct and indirect resources (Naci et al., 2010). However, implications of relapses and disability worsening on societal economic costs have scarcely been studied. This is partly due to a lack of longitudinal data that are required for comprehensive analyses (Naci et al., 2010). Consequently, this thesis focuses on how clinical hallmarks in patients with MS affect the societal economic costs of the disease. The two hypotheses below are derived to address this research aim.

Regarding disability worsening it is known that the overall economic burden increases as disease severity increases (Karampampa et al., 2012a; Kobelt et al., 2017b). Furthermore, the distribution of costs changes with increasing disease severity (Naci et al., 2010). While DMTs are the main cost driver in the early stages of the disease, productivity losses account for the majority of costs for more severely disabled patients. However, the societal economic implications regarding disability worsening remain not well understood. The aforementioned observations lead to the first hypothesis: Societal economic costs are significantly higher in patients with disability worsening compared to patients being free of disease activity.

Research to date has shown that the economic burden of MS increases in periods of relapses. However, results are inconsistent between studies. This may arise from different implications of relapses with respect to disease severity, age, disease duration or therapy. Kobelt et al. (2006b) suggested that relapses have a stronger implication in patients with limited disability. However, a comprehensive test of this hypothesis is still needed and thus constitutes the second hypothesis of this thesis.

My second hypothesis is that young patients and those with a short disease duration have higher societal economic relapse costs than older patients and those with longer disease duration.

Hence, this thesis determines priorities in medical research to reduce the societal economic burden in patients with MS. A prerequisite is an instrument with sufficient psychometric properties to assess the societal economic costs. Despite the long tradition of health economic evaluations in MS, there was no such instrument available (chapter 1.3.1). Consequently, I first use and validate the instrument termed Multiple Sclerosis Health Resource Survey (MS-HRS).

2 Data source

Pooled two-year data were extracted from two single arm, prospective, multicenter, non-interventional long term studies conducted in Germany. The Post-Authorization Non-Interventional German safety study of GilEnyA (PANGAEA) assessed the long term safety and efficacy of fingolimod. Beta interferons and glatiramer acetate (BRACE) therapies were assessed in the Prospective Pharmacoeconomic Cohort Evaluation (PEARL) (Vormfelde et al., 2016; Ziemssen et al., 2015b). Additionally, both studies collected pharmacoeconomic data.

2.1 Study setting

Recruitment into PANGAEA took place after market approval for fingolimod in April 2011 until December 2013 for patients receiving 0.5 mg fingolimod daily. The observational period of the main and pharmacoeconomic substudy ended after 60 months in December 2018 and after 24 months in December 2015, respectively. Recruitment into PEARL took place between September 2010 and March 2011 for patients receiving BRACE therapies for at least 30 days. The observational phase of the study ended after 24 months in March 2013. As the participants in both studies were recruited in Germany from overlapping centers and during an overlapping timeframe, the environmental and pharmacoeconomic conditions are expected to be similar (Alsop et al., 2017; Vormfelde et al., 2016; Ziemssen et al., 2015b). An overview of the timeline is shown in figure 4.

Figure 4: Timeline of the pharmacoeconomic (sub)studies PEARL and PANGAEA.

2.2 Clinical and economic data

Study data were collected during routine physician consultations at baseline, every 3 months during follow up and at the final visit after 24 months resulting in nine data entry points per patient. Demographic and clinical data were collected using interviews, examinations and medical records. Patient-reported outcomes (PROs) were completed by participants on a regular basis using a (electronic) case report form (CRF) in the presence of a health care professional.

Data on resource utilization, treatment satisfaction (Treatment Satisfaction Questionnaire for Medication (TSQM-9)) and compliance with the medication were collected every three months in the pharmacoeconomic (sub)study (Atkinson et al., 2004). Every six months patients reported their health-related quality of life (HRQoL) based on the EuroQol-5 Dimensions (EQ-5D) and the Patient Reported Indices for Multiple Sclerosis (PRIMUS). Every 12 months activity limitations and neurological disability were assessed using the PRIMUS and the UK Neurological Disability Scale (UKNDS), respectively (Doward et al., 2009; Kind, 1990; Sharrack and Hughes, 1999). Data obtained in the studies are listed in table 7.

	Baseline visit	Follow-up	Final visit (24 months)
Clinical Data			
EDSS	✓	Every 6 months	✓
Relapse information	✓	Every 3 months	✓
Patient Reported Outcomes			
Consumption of resources	✓	Every 3 months	✓
UKNDS	✓	At 12 months	✓
EQ-5D, PRIMUS-A, PRIMUS-L	✓	Every 6 months	✓
Compliance questionnaire	✓	Every 3 months	✓
TSQM-9	✓	Every 3 months	✓

Table 7: Clinical and patient reported data obtained during the studies.

The EQ-5D is a generic instruments for assessing HRQOL with high psychometric properties (Fisk et al., 2005; Kind, 1990). The assessment is carried out by the test persons based on five dimensions: mobility, independence, everyday activities, pain and anxiety. These dimensions are evaluated on a three-level scale (EQ-5D-3L). The lowest value

indicates no problems, the highest value extreme problems. Additionally, the respondent evaluates his HRQOL on a visual analogue scale ranging from zero to 100. This provides additional information that is not reflected in the previous dimensions.

The PRIMUS measures MS-specific the HRQOL (PRIMUS-L) and activity limitations (PRIMUS-A) (Doward et al., 2009). Depending on the short or long form of the questionnaire, 45 or 22 dichotomous questions are recorded to determine the quality of life, where a high sum value indicates a low quality of life. The activity questionnaire comprises 15 questions related to physical tasks on which respondents assess the degree to which they are capable of performing the tasks. A high score indicates a high activity restriction (McKenna et al., 2010). The reliability and validity of the three scales is rated as high, with the reliability for the scales for recording quality of life and activity restrictions is higher compared to the symptom scale (McKenna et al., 2010).

The UKNDS measures the functional status of the patient in 13 mutually exclusive domains (Heesen et al., 2007; Sharrack and Hughes, 1999). These are cognition, mood, vision, speech, swallowing, upper limb function, mobility, bladder function, bowel function, sexual function, fatigue, pain and "other" as a collective term of other unrecorded dimensions such as dizziness or cramps. Each domain is assessed by the respondent based on three to eight questions. The degree of impairment is converted from zero (normal function) to five (total loss of function) depending on the answers to an ordinal scale (Sharrack and Hughes, 1999). Rossier and Wade (2002) proved that the UKNDS is a reliable and valid instrument.

The TSQM-9 is a psychometrically validated, generic measure to assess the patients' treatment satisfaction with their medication (Atkinson et al., 2004). The TSQM-9 is based on the three scales that assess the patients' perception of medication effectiveness, convenience and global satisfaction. Each scale comprises three items with answers rated from one (low) to seven (high) on a Likert scale. Higher scores on the TSQM-9 indicate a higher treatment satisfaction (Atkinson et al., 2004).

2.3 Study population

Patients were eligible to participate if they were diagnosed with RRMS and if they received their medication in accordance with the summary of product characteristics. To be admitted to PANGAEA, patients had to be prescribed fingolimod. For admission to PEARL, patients must have been continuously treated with interferon (Avonex®, Betaferon®, Extavia®, Rebif®) or glatiramer acetate (Copaxone®) for at least 30 days. DMTs had to be prescribed independently of study participation. Furthermore, informed consent had to be given prior to study inclusion. There were no exclusion criteria except for the contraindications mentioned in the respective summary of the treatment information. Sample sizes in PANGAEA and PEARL were calculated in order to obtain representative data on the most commonly prescribed DMTs (BRACE and fingolimod). Furthermore, geographical representativeness was ensured by the regional distribution of participating practices and centers.

2.4 Ethical considerations

The PROs and clinical outcome measures evaluated in the analysis were obtained exclusively from the PEARL and PANGAEA. Prior to study inclusion, informed consent to participate and to pass on their pseudonymized data was obtained from patients. Data were collected in the presence of a physician or a medical assistant at the MS centers and practices. The persons involved in the evaluation had no access at any time to information that could have violated the anonymity of the study participants. The studies were financed by Novartis Pharma GmbH and voted positively by the Ethics Committee of the University of Dresden. PANGAEA and PEARL are registered as a non-interventional studies under CFTY720DDE02 and CNVF233ADE08, respectively (VFA, 2010; VFA, 2011).

2.5 Data management

Data of PANGAEA and PEARL were initially available in sas7bdat format in several data sets, each referring to one questionnaire, so that UKNDS, PRIMUS, TSQM-9, EQ-5D, PRIMUS-A, PRIMUS-L and consumption of resources were separated. The databases were structured horizontally, with each row representing one observation. Using the Statistical Package for Social Science (SPSS) function "merge files", I combined the separated data sets. The variables "Internal_Patient_ID" and "Visit_ID" were used as identification criteria for the doubtless merger of the data.

Whenever possible, the Multiple sclerosis documentation system 3D (MSDS3D) was used to ensure the highest possible data quality (Haase et al., 2018b). Additionally, all values were checked for plausibility using descriptive statistics. For variables with implausible maximum and minimum values, ranges were defined that resulted from the possible range of the observation period. For example, a resource used daily could be used a maximum of 92 times in a three-month period. Values outside this range were treated as absent.

Statistical methods for imputation of missing values were not used. Missing values were only replaced for binary questions, if the actual value could be derived beyond doubt from the answer. When asked about the amount of an occupational disability pension, no value was entered for subjects without such a pension during data collection. These fields were manually set to zero. The same procedure was used for the question of the number of visits to the doctor: if a patient indicated that he had visited a doctor in the last three months, missing entries were replaced by zero. SPSS version 25 was used for data management.

3 Methods

The following chapters are excerpts of manuscripts submitted to or published in peer review journals:

- *Chapters 3.1 and 3.2 are based on an article titled “Multiple Sclerosis Health Resource Utilization Survey (MS-HRS): Development and Validation Study” by Ness et al. (2020a).*
- *Chapter 3.3 is based on an article titled “Differentiating societal costs of disability worsening in multiple sclerosis” by Ness et al. (2019a).*
- *Chapter 3.4 is based on an article titled “Real-World Evidence on the Societal Economic Relapse Costs in Patients with Multiple Sclerosis” by Ness et al. (2020b).*

3.1 Questionnaire development and validation

3.1.1 Development process

A German expert group consisting of neurologists, healthcare administrators, psychologists and MS nurses developed the first version of the Multiple Sclerosis Health-resource Utilization Survey (MS-HRS) at the University Hospital Dresden in 2009. Following intensive feedback from physicians, nurses and patients, a second more time-efficient version was created in 2016. The current adaptive version of the MS-HRS in its German and translated English version is in line with the recommendations on the core elements of a standardized resource questionnaire (Thorn et al., 2018).

3.1.2 Identification of resource consumption

Relevant resources were identified from the societal perspective. Costs were categorized into direct medical (inpatient stays, day admissions (outpatient), physician consultations, examinations, over-the-counter (OTC) medication, medical consumables and professional care), direct non-medical costs (informal care, investments/ purchases) as well as indirect costs. Within indirect costs, I considered absenteeism, either short term (sick leave) or long-term (disability pension), but also presenteeism, which refers to impaired performance during work. (Figure 5).

Figure 5: Identified direct and indirect cost items for patients with MS. Reproduced without modifications from Ness et al. (2019b).

Information on resource consumption were obtained from studies, guidelines, textbook knowledge, administrative and accounting data and expert opinions (Fowler, 1995; Fowler and Floyd, 1993; Kobelt et al., 2006b; Reilly et al., 1993). Within the expert group it was decided which resource items are both relevant and nontrivial in magnitude (Luce et al., 1996). Keeping multiple use cases in mind, I divided the tool in a core and an additional set of items. Core elements are constitutive for the health-resource utilization model and sufficient to capture the scope of resource utilization whereas the additional questions should foster the understanding of resource utilization (Table 8).

Domain	Items	Valuation
Direct medical costs		
Inpatient care/ day admission	Days in hospital, rehabilitation and nursing ward	Standardizes cost units ¹⁶
Ambulant consultations	Visits at general practitioners, neurologists, others specialist (Urologists, ophthalmologists, psychiatrists), MS-nurses, physical therapists, psychologists, occupational therapists, opticians, speech therapists, acupuncturists and other alternative healing methods	Standardizes cost units ¹⁶
Investigations/ Diagnostics	MRI, CT, spinal tap, blood test, others	Physicians' Fee Schedule ¹⁶
OTC medication	Medication, recommending doctor	Patient reported
Disease-modifying therapies		Annual therapy costs as calculated from medication report ^{18-24,26}
Home help and support of professionals	Professional help, household care, personal assistance	Human capital approach
Medical consumables	Medical consumables	Patient reported
Direct non-medical costs		
Investments, equipment	House and car modifications, walking aids, wheelchair (manual, electric), scooter, sickbed, others, payer, grants	Patient reported
Informal care	Time taken for preparation of meals, climbing stairs, personal care, administration of drugs, transport, others; reduction of working hours of relatives.	Human capital approach
Indirect costs		
Employment and labour productivity	Full or part time work, sick leaves, reduced working time, change of work, loss of earning	Human capital approach
Employment and labour productivity	Absence hours (MS, others), total working hours, productivity scale	Human capital approach

Table 8: Core Items of the MS-HRS.

Abbreviations: over-the-counter (OTC), magnetic resonance imaging (MRI), computed tomography (CT).

Reproduced with layout modifications from Ness et al. (2020a).

3.1.3 Quantification of resource consumption

I used the most accurate method of micro-costing defined as “direct enumeration and costing out of every input consumed in the treatment of a particular patient” to quantify resources (Raftery, 2000). Recall periods of resource utilization have been determined to avoid over- and underreporting. I considered that longer recall periods lead to telescoping effects in which events that occurred outside the time frame are mistakenly included in the intended period (Stull et al., 2009). Furthermore, I took into account that more frequent events and those that are less salient, are less likely to be recalled accurately over a long time span. In conclusion, the recall period may differ with respect to the nature of the cost item but should never exceed twelve months. It is noticeable that

unified recall periods across all items of the survey made the questionnaire more intelligible for patients. In the results, I reported all calculated costs as per quarter.

3.1.4 Valuation of resource consumption

Evaluations conducted from the societal perspectives are intended to reflect the societal opportunity costs, which are equal to market prices on perfectly competitive markets. Nevertheless, markets in the healthcare sector are imperfect due to statutory regulations. Hence, societal opportunity costs have to be estimated in most cases. Bock et al. (2015) approximated societal opportunity costs based on published accounting data of individual actors in the health care system or on regular publications of official statistics. These data were used wherever available. An overview of all direct (non-)medical and indirect resource items is given in table 9.

Due to a lack of data, few valuations were calculated from existing values. Visiting a nurse was rated with the lowest monetary value of 16.42€. Furthermore, a psychologist contact was monetarily valued as visiting a psychotherapist. The valuation for other specialists (e.g. psychiatrist, urologist, and optician) was calculated as the mean of the given examples. Societal opportunity costs for investigations were approximated from the Physicians' Fee Schedule (KBV, 2011).

Indirect costs (absenteeism, early retirement, presenteeism) were valued using the human capital approach (van den Hout, 2010). Absenteeism was defined as not showing up for work while presenteeism refers to reduced work productivity due to health problems. For short term absence (sick leave), I multiplied the average daily working hours with the average labor costs in Germany (29.90€/h) (Bock et al., 2015). Data for this calculation were obtained from the Federal Bureau of Statistics (BPB, 2011; IAB, 2011).

$$\frac{1406.2 \text{ working hours (2011)}}{233 \text{ working days}} * 29.90\text{€ hourly salary (2011)} \quad (\text{II})$$

For long term absence (disability pension), I multiplied the average daily labor costs (127.38€/day) with the average number of working days (233 days) and the disability percentage (0-100%) leading to a maximum amount of 29,656.76€ per year.

To value direct non-medical informal care I used the opportunity cost method. The informal caregiver's forgone benefits were approximated by multiplying the hours spent on informal care by the opportunity costs of leisure time (21.09€/h). The maximum of hours dedicated to informal care was set to 60 hours per week according to the German policies.

To ensure the comparability of cash flows, prices from different periods were adjusted to the 2011 price level using harmonized consumer price index published by the Federal Statistical Office as recommended for Germany (IQWiG, 2017). This applies to patients' self-reported medication, investments, medical consumables and DMTs. Consumption of remaining resources was valued with prices from the year 2011 in order to generate comparable costs within the validation population.

Domain (# of questions)	Valuation	Domain (# of questions)	Valuation
Direct medical costs			
Inpatient		acupuncturists	18.24€
Hospital	593.04 €	other alternative healing methods	27.40€
Rehabilitation	121.85 €	Investigations/ diagnostics	
Nursing	69.80 €	(MRI)	120.21 €
Outpatient		CT	73.78 €
Hospital	385.48 €	Lumbar puncture	38.90 €
Rehabilitation	46.68 €	Blood tests	1.10 €
Nursing	46.15 €	Others	60.66 €
Ambulant consultations		OTC medication	Patient reported
General practitioner	20.06 €	Disease modifying therapy	Table 10
Neurologist	44.72 €	Medical consumables	Patient reported
Other specialists	34.73 €	Home Help/ professional care	27.57 €
Nurse	16.42 €	Direct non-medical costs	
Physiotherapist	16.42 €	Equipment, aids, modifications	Patient reported
Psychologist	78.08 €	Informal care	21.09 €
Occupational therapist	37.51 €	Indirect costs	
Optician	34.78 €	Sick leave	127.38 €
Speech therapists	38.59 €	Disability pension (full year)	29,656.76 €

Table 9: Valuation of resource costs applied in the MS-HRS (price year 2011). Abbreviations: over-the-counter (OTC), magnetic resonance imaging (MRI), computed tomography (CT).

Reproduced with layout modifications from Ness et al. (2020a).

To calculate annual costs for DMTs, I used Defined Daily Dose net-costs multiplied by 365 days (Zeller, 2012). Net costs account for statutory manufacturer discounts as well as pharmacy discounts. Table 10 lists pharmaceutical prices for BRACE therapies, Gilenya and Tysabri for the years 2010-2017.

Year	AVO	REB	BET	EXT	COP	GIL	TYS
2010	18069.2 7	22626.4 6	17977.1 4	15528.4 5	16623.4 5	n.a.	24625.1 0
2011	18611.3 5	23305.2 5	18516.4 5	15994.3 0	17122.1 5	26698.5 8 ^a	25363.8 5
2012	18182.1 9	22900.4 6	18132.0 7	15604.6 8	16377.9 3	25907.5 6	24586.5 8
2013	17981.0 0	22765.3 5	17815.2 9	15431.9 3	16197.0 0	22571.4 4	24214.4 1
2014	19437.3 1	24432.9 8	17601.9 5	15441.4 8	17035.6 1	18965.3 6	26194.9 2
2015	19333.9 6	24155.2 5	17517.7 0	15370.2 5	16733.3 2	20393.7 3	24280.7 5
2016	19247.4 2	23987.2 7	17384.0 9	14680.2 5	16648.4 8	21516.7 2	23321.0 6
2017	18865.1 0	23529.3 8	17047.8 1	15715.4 9	16352.2 6	20784.6 8	22902.0 2

Table 10: Defined Daily Dose net costs in the respective years for DMTs multiplied with 365. If no information was available, the last known prices were adjusted with the consumer price index. All prices in Euro.

Abbreviations: Disease modifying therapies (DMTs), Betaferon® (BET), Rebif® (REB), Avonex® (AVO), Copaxone® (COP), Extavia® (EXT), Gilenya® (GIL) and Tysabri® (TYS).

^a as listed in the technology assessment report (Novartis, 2011).

Reproduced with layout modifications from Ness et al. (2020a).

3.2 Assessment of psychometric properties

3.2.1.1 Practicability and reliability

Practicability was determined by the response rate of patients completing the MS-HRS. In addition, I compared characteristics of completers vs. non-completers to avoid selection bias. For reliability and validity analyses, I focused on the core elements of the survey (Table 8). As I did not intend to evaluate DMTs at this step, costs for DMTs are not part of methodological evaluation.

For reliability analysis, I estimated test-retest reliability of the total costs in a group of stable patients over three months. Stable patients were defined as not having a relapse or increase of the EDSS during the three-month retest period and another three months prior to the assessment. Thresholds for intraclass correlation coefficient (ICC) were applied as recommended ($ICC < 0.5$: poor reliability; $0.5 \leq ICC < 0.75$: moderate; $0.75 \leq ICC < 0.9$: good; $ICC > 0.90$: excellent) (Koo and Li, 2016). Guttman's Lambda-2 and Lambda-6 will be reported for the monetarized-standardized items of health-resource utilization model with respect to the heterogeneous structure providing a lower bound estimate of the consistency of the pricing approach. In an ordinary setting for test construction, desired levels would lay above 0.7. Since the survey is not based on latent constructs but instead on real costs, consistency analysis is not part of the primary evaluation of reliability but rather provides additional insights for the model and the cost components.

3.2.1.2 Validity

Repeated expert consensus meetings of health economists, psychologists and neurologists were conducted to secure face validity in terms of consistency and completeness. Construct validity was assessed as convergent and discriminant validity via correlations with associated patient-reported outcomes (PROs) and known-groups analyses by EDSS (ranges: 0-1.0; 1.5-2.5; 3.0-4.0; 4.5-5.5). The selected PRO measures were the EQ-5D, the UKNDS and the PRIMUS (Doward et al., 2009; Graf von der Schulenburg et al., 1998; Greiner et al., 2005; Sharrack and Hughes, 1999). I expected to find significantly higher costs in groups with higher EDSS scores and correlations above 0.40 for convergent validity. As I did not intend to evaluate DMTs, the TSQM should present correlations clearly below 0.3 for discriminant validity (Atkinson et al., 2005). An overview of assessed psychometric concepts is shown in table 11.

Psychometric concepts	Definition	Analysis
Practicability	Response rate of persons completing the questionnaire (DeVon et al., 2007)	Response rate
Test-retest reliability	Temporal stability of a test from one measure to the other within the same person (Drost, 2011)	ICC; additionally: Lambda-2, Lambda-6
Face validity	The instrument encompasses all items to represent the concept to be measured (DeVon et al., 2007)	Expert group consensus
Construct Validity	<p>Degree to which an instrument measures the construct it is intended to measure (DeVon et al., 2007).</p> <p>Convergent validity Correspondence between constructs that are theoretically similar (DeVon et al., 2007).</p> <p>Discriminative validity Capability to differentiate between constructs that are theoretically similar (DeVon et al., 2007).</p>	Correlation tests

Table 11: Validity and reliability measurements assessed for the MS-HRS.
Abbreviations: Intraclass Correlation Coefficients (ICC).

3.3 Definition of clinical outcome measures

Changes in disability were assessed at a regular basis using the EDSS every six months during clinical visits. CDW measured by EDSS was defined as: ≥ 1.5 -point increase for a reference-score of 0; ≥ 1 -point increase for a reference-score 1.0-5.0; and ≥ 0.5 point increase for a reference score greater than 5.5. Due to its higher sensitivity, I implemented a roving reference score instead of a fixed baseline reference score to detect CDW (Kappos et al., 2018a). A roving EDSS score is independent of the baseline so that EDSS assessments at baseline, months six or twelve can be applied as reference. Increase in EDSS and confirmation of the increase can take place ≥ 6 and ≥ 12 months apart from the reference score, respectively. In order to have sufficient data to determine CDW and in accordance with earlier studies patients with at least three EDSS assessments were included (Lorscheider et al., 2016).

CDW events in which no relapse occurred between reference and confirmation assessment were declared as PIRA (Figure 6). If a relapse occurred between reference and confirmation assessment, the event was classified as RAW. Furthermore, I distinguished RAW from RWW applying to patients, who recover from relapses without having an increase in accumulated disability as defined by CDW. Patients achieving freedom of relapse activity and CDW were defined as DAF. Relapses were assessed every 3 months according to the clinical judgement of physicians during routine clinical visits.

Figure 6: Schematic of the assessment of clinical descriptors.

A roving EDSS reference score was used to detect CDW-6 events within two years. CDW-6 was defined as increases in EDSS of ≥ 1.5 points for reference score between 1.0-5.5 or ≥ 0.5 points for reference score ≥ 6.0 . Physicians assessed EDSS and relapses every six and three months, respectively.

Reproduced without modifications from Ness et al. (2019a).

3.4 Assessment of relapse associated costs

Patients were considered as 'relapse active' if they had at least one relapse during the two-year follow-up whereas 'relapse inactive' refers to patients without relapses during the follow-up time. I calculated relapse costs as the cross-sectional difference in mean quarterly costs between relapse-active patients during relapse quarters and relapse-inactive patients. For the relapse costs analysis, I applied a propensity score (PS) matching to compensate for differences between relapse active and relapse inactive patients. Patients were matched in a 1:1 ratio using a nearest neighbor matching. PS were calculated as the probability to be part of the relapse active group according to the following baseline predictors: EDSS score, time since diagnosis of MS, disease-modifying therapies (DMT), age and gender. These variables were selected for matching as previous studies showed that a shorter disease duration and lower EDSS are associated with higher relapse activity, that women tend to experience more relapses (Kalincik, 2015). Furthermore, DMTs are associated with a varying effectiveness in reducing relapse rate (Li et al., 2019).

In addition to the cross-sectional approach of calculating relapse costs, I analyzed relapse associated costs longitudinally. Each relapse follow-up was re-baselined and costs for corresponding intervals before (prior relapse) and after this interval (post relapse) were calculated additionally (Figure 7). On an intraindividual basis, I calculated the difference between quarterly costs prior relapse and quarterly cost during relapse. For the analyses, I excluded 77 persons who had less than three visits. The excluded persons did not statistically significant differ in their sociodemographic or disease characteristics. Relapse associated costs are additionally reported stratified by body mass index (BMI). The BMI was determined by weight and height of the patients and classified as under- and normal weight ($<25 \text{ kg/m}^2$), overweight including pre-obesity (25 kg/m^2 - 30 kg/m^2) and obesity ($\text{BMI}>30 \text{ kg/m}^2$) (WHO, 1995).

Figure 7: Assessment of changes in resource utilization (prior-, during and post-relapse). Abbreviations: MS- Health Resource Survey (MS-HRS).

3.5 Statistical analyses

Continuous values were reported as arithmetic means and standard deviations (SD, $\pm$). Ordinal values were described by median interquartile range (IQR) and nominal values as percentages (%).

For the validation of the questionnaire, one-way random ICC were used to calculate estimates for the test-retest reliability. All other correlations were calculated as Spearman's rank correlation coefficient. Kruskal-Wallis-H-tests were conducted for known group analyses. Mann-Whitney-U-tests with adjustments by the Bonferroni correction for multiple tests were applied for pairwise comparisons.

Statistical analyses of healthcare resource use and cost data has several challenges. The data are often skewed, can have multimodal distributions and heavy tails (Mihaylova et al., 2011). To take care of these issues I applied a Generalized Linear Mixed Models (GLMM) with a negative binomial distribution and a log link function for model-based analysis with monetary outcomes.

The influence of PIRA and RAW on (i) total costs excluding DMTs, (ii) direct medical costs excluding DMTs, (iii) direct non-medical costs and (iv) indirect costs were reported through incidence rate ratios (IRR) and corresponding 95% confidence intervals (95% CI). For example, an IRR of 1.5 indicates that a specific group had a 50% higher quarterly resource consumption averaged over two years and compared to patients that were referred to as achieving DAF, while holding the other variables constant in the model. Patients achieving DAF served as the reference group for PIRA and RAW in the statistical models. Multivariate models were adjusted for sex, age, EDSS, data source and time, which indicated the sequence of assessments per patient.

Statistical differences in relapse costs between patient and disease strata were derived from Generalized Linear Models (GLM) with a negative binomial distribution and the respective cost category as the dependent variable. The models included the characteristic (gender, age, BMI, disease duration, EDSS or therapy) of interest as well as a variable indicating the relapse status (relapse active vs. relapse inactive) and the interaction of both. For example, to assess statistical differences in total relapse costs regarding disease duration, we included the disease duration variable, the variable indicating the relapse status and the interaction of these two variables as independent variables.

All reported p-values were compared to an alpha-error level of 5%. No imputations were made for missing values. Statistical analyses were performed using SPSS statistics for Windows version 25 and R version 3.5.1.

4 Results

The following chapters are excerpts of manuscripts submitted to or published in peer review journals:

- *Chapters 4.1 is based on an article titled “Multiple Sclerosis Health Resource Utilization Survey (MS-HRS): Development and Validation Study” by Ness et al. (2020a).*
- *Chapter 4.2 is based on an article titled “Differentiating societal costs of disability worsening in multiple sclerosis” by Ness et al. (2019a).*
- *Chapter 4.3 is based on an article titled “Real-World Evidence on the Societal Economic Relapse Costs in Patients with Multiple Sclerosis” by Ness et al. (2020b).*

4.1 The Multiple Sclerosis Health Resource Survey

4.1.1 Study population

In total, 2207 of 2388 patients completed the questionnaire at baseline and were therefore included in the validation process. The population was in average 41.73 years old and was mostly female (73.10%) (Table 12). Employed patients (61.76%) were working predominantly full time (60.79%).

In terms of relapses within the previous year, active (54.96%) and non-active (45.05%) patients were balanced in the population. Participants reported an average disease duration of 7.54 ± 6.11 years and an EDSS score of 2.43 ± 1.57 .

Baseline Characteristics		
Sociodemographic		
	Female gender (n (%))	1609 (73.10)
	Age (mean±SD)	41.73±10.19
	Education	
	intermediate/ advanced school-leaving certificate	439 (20.92)
	vocational training/ university degree	1410 (67.21)
	Other	249 (11.87)
	Employment	
	percentage employed	1347 (61.76)
	percentage full time	814 (60.79)
Disease Characteristics		
	Number of relapses in previous year, n (%)	
	0	1192 (54.96)
	1	612 (28.22)
	2	267 (12.31)
	≥3	98 (4.52)
	Duration of disease since diagnosis, years	
	mean±SD	7.54±6.11
	median (IQR)	6.00 (3.00, 11.00)
	EDSS score	
	mean±SD	2.43±1.57
	median (IQR)	2.00 (1.00, 3.50)

Table 12: Baseline characteristics of the study population included in the validation process (N=2207).

Reproduced with layout modifications from Ness et al. (2020a).

4.1.2 Validation of the questionnaire

4.1.2.1 Practicability

Overall, 2207 out of 2388 patients completed the questionnaire (response rate: 92.40%). Therefore, the answers of 2207 patients could be used to calculate total costs and all other parts of the health-resource model. Non-completing patients (n=181) did not differ in their gender distribution, age or EDSS score, but presented a slightly longer disease duration (9.67 ± 7.21 , $p < 0.001$) (Table 13).

Baseline Characteristics	Selected (N=2207)	Not selected (N=181)
Female gender (n (%))	1609 (73.10)	127 (70.90)
Age (mean±SD)	41.73±10.19	41.06±9.98
Duration of disease since diagnosis, years		
mean±SD	7.54±6.11	9.67±7.21
median (IQR)	6.00 (3.00, 11.00)	8.00 (1.00, 19.00)
EDSS score		
mean±SD	2.43±1.57	2.68±1.73
median (IQR)	2.00 (1.00, 5.50)	2.00 (1.00, 5.50)

Table 13: Comparison of baseline characteristics of included and excluded patients for the validation process.

4.1.2.2 Reliability

I assessed the reliability mainly as test-re-test reliability in a stable subgroup of patients. Overall, 1192 of 2207 (54.0%) patients fulfilled the criteria of presenting a stable MS within that period. The ICC for that group was 0.828 over a course of three months. In addition, Guttman's Lambda-2 ($\lambda_2=0.679$) and Lambda-6 ($\lambda_6=0.694$) indicated an acceptable consistency between the standardized monetarized items of the health-resource model (excluding DMT costs).

4.1.2.3 Validity

I analyzed the construct validity for total costs of the MS-HRS via known-groups (excluding DMT costs). In all four EDSS groups, I found significantly different MS-related total costs and sub costs (Table 14). Further, all pairwise comparisons indicated significant differences in the direction as expected before ($p<0.001$ for total costs, $p<0.05$ for sub costs).

Total costs		Direct medical (excluding DMTs)		Direct non-medical		Indirect	
Mean ±SD	Median	Mean ±SD	Median	Mean ±SD	Median	Mean ±SD	Median
EDSS 0-1.0							
1099 ±2211	150	309 ±910	99	12 ±108	0	777 ±1822	0
EDSS 1.5-2.5							
2295 ±3647	585	541 ±1724	165	20 ±154	0	1734 ±2820	0
EDSS 3.0-4.0							
3112 ±3979	1334	789 ±1998	249	45 ±152	0	2280 ±3178	127
EDSS 4.5-5.5							
4733 ±4639	4158	1003 ±2295	411	158 ±428	0	3572 ±3572	3637

Table 14: Direct (non-)medical and indirect costs (excl. DMTs) stratified by EDSS scores. Costs differed significantly between EDSS strata. Reproduced with layout modifications from Ness et al. (2020a).

Convergent validity analyses showed that total costs correlated positively with an increased (patient-reported) disability (UKNDS sum score: $r=0.411$), the lost ability to participate in daily routines and activities (PRIMUS Activities: $r=0.423$) and negatively with health-related quality of life (HRQoL; PRIMUS QoL: $r=0.350$; EQ-5D: $r=-0.342$) (all $p<0.001$).

For discriminant validity, correlations of total costs with TSQM scores ranged from -0.006 (convenience) to -0.216 (effectiveness). As expected, the inclusion of DMT costs lowered the correlations with all PROs (by 0.07 on average) but the relations between correlations were maintained (Table 15).

	Correlation Coefficient with total costs (excl. DMTs)	Correlation Coefficient with total costs (incl. DMTs)
Convergent validity		
UKNDS sum score	0.411	0.348
PRIMUS activity	0.423	0.346
Primus Quality of life	0.350	0.290
EQ5D	-0.342	-0.274
Discriminant validity		
TSQM-9		
Effectiveness	-0.006	-0.031
Convenience	-0.216	-0.187

Table 15: Correlation coefficients for convergent and discriminant validity. Total costs correlated positively with disability and negatively with HRQoL.

4.1.3 Resource utilization

The majority of patients stated that they used direct medical services in the past three months. In contrast, less than every second patient (44%) caused indirect medical costs and 16% claimed direct non-medical costs. Besides DMTs, indirect costs were the main cost drivers ahead of direct medical and direct non-medical costs.

Patients were mainly treated in outpatient practices (85%) and less often during inpatient hospital stays (6%) and daycare admissions (3%). Among the direct medical costs, inpatient treatment caused the highest costs (315.06€±1587.09€), followed by consultations in the primary sector (209.87±292.73€) and day admissions in hospitals (32.22€±311.68€). Total costs per quarter averaged on 2462±3650€ (median: 631€) without DMTs and 7126€±3697€ (median: 5871€) including DMTs. Therefore, annual costs of a mild to moderate RRMS ranged between 9528±14,603€ (without DMTs) and 28,203±14,808€ (with DMTs) (Table 16).

	Users	Mean±SD	Median (IQR)
Direct medical costs (without DMT)	2068 (93.70%)	601.30±1708.55	174.00 (75.00, 417.50)
Inpatient care	138 (6.25%)	315.06±1587.09	0.00 (0.00, 0.00)
Day admission	65 (2.95%)	32.22±311.68	0.00 (0.00, 0.00)
Consultations	1817 (82.33%)	209.87±292.73	109.50 (44.72, 268.32)
Examinations	1429 (64.75%)	32.34±35.63	30.05 (0.28, 45.77)
OTC Medication	693 (31.40%)	15.02±46.43	0.00 (0.00, 10.00)
Professional care	115 (5.21%)	7.02 (44.79)	0.00 (0.00, 0.00)
Disease modifying therapies	2185 (99.00%)	4733.24±820.30	4629.11 (4280.54, 5656.61)
Direct non-medical costs	352 (15.95%)	44.65±229.51	0.00 (0.00, 0.00)
Investments	38 (1.72%)	9.14±165.80	0.00 (0.00, 0.00)
Informal care and community service	337 (15.27%)	37.25±158.57	0.00 (0.00, 0.00)
Indirect costs	977 (44.27%)	1816.50±2880.74	0.00 (0.00, 3037.00)
Short-term absence	320 (14.50%)	441.53±1782.18	0.00 (0.00, 0.00)
Disability pension	352 (15.95%)	852.98±2062.98	0.00 (0.00, 0.00)
Presenteeism	475 (21.52%)	522.00±1333.18	0.00 (0.00, 0.00)

Table 16: Health resource utilization in persons with MS per quarter (N=2207).

Abbreviation: over-the-counter (OTC).

Reproduced with layout modifications from Ness et al. (2020a).

4.2 Societal economic impact of disability worsening

4.2.1 Study population

In total, 1959 patients had an EDSS score <6 and were therefore included to analyze the economic impact of disability worsening. Patients reported an average age of 41.62 ± 10.04 years and were mostly female (72.82%). The mean disease duration of the total population was 7.30 ± 5.95 years with a baseline mean EDSS value of 2.26 ± 1.37 . In terms of relapses in the year before baseline, patients were balanced between active (43.56%) and inactive (56.44%).

Of the total population, 985 patients (50.28%) were classified achieving DAF status after two years, 166 (8.47%) fell in the group PIRA, 150 (7.66%) into RAW and 407 (20.78%) had RWW (Table 17). Due to an insufficient number of EDSS assessments, 251 (12.81%) patients did not meet the definition of any aforementioned groups. For all subsequent analyses, these patients were summarized under the label “others”.

PIRA patients were older (44.48 ± 9.80 years of age) and longer affected by the disease (8.48 ± 7.13 years) compared to the other groups of interest. Furthermore, the rate of women was highest in patients with RAW (78.67%).

In the year prior baseline, patients grouped in RAW and RWW were significantly more often relapse-active (51.68% and 57.18%, respectively) than patients with PIRA (30.91%) ($p < 0.001$).

Nearly two third (62.17%) of all patients were employed at baseline with the lowest rate reported for PIRA patients (57.23%). RAW and RWW patients had a higher sick leave rate (17.33% and 17.44%, respectively) compared to PIRA (10.24%).

Baseline Characteristics	Total (N=1959)	DAF (N=985)	PIRA (N=166)	RAW (N=150)
Socio-demographic				
Female gender (n (%))	1425 (72.82)	689 (70.02)	115 (69.28)	118 (78.67)
Age (mean $\pm$ SD)	41.62 $\pm$ 10.04	41.88 $\pm$ 9.93	44.48 $\pm$ 9.80	42.09 $\pm$ 10.17
18-29 years (n(%))	271 (13.83)	125 (12.69)	13 (7.83)	22 (14.67)
30-49 years (n(%))	1233 (62.94)	624 (63.35)	96 (57.83)	90 (60.00)
50 years and older (n(%))	455 (23.23)	236 (23.96)	57 (34.34)	38 (25.33)
Patients in working age (n(%))	1920 (98.01)	965 (97.97)	162 (97.59)	147 (98.00)
Employed (n(%))	1218 (62.17)	624 (63.35)	95 (57.23)	92 (61.33)
Employed at full time (n(%))	722 (36.86)	385(39.09)	58 (34.94)	54 (36.00)
Employed with sick leave (past 3 months) (n(%))	265 (13.53)	101 (10.25)	17 (10.24)	26 (17.33)
Employed with disabili- ty pension (n(%))	289 (14.75)	136 (13.81)	28 (16.87)	31 (20.67)
Not living alone (n(%))	1494 (81.42)	746 (80.56)	123 (79.87)	109 (80.74)
Disease Characteris- tics				
Disease duration (mean $\pm$ SD)	7.30 $\pm$ 5.95	7.43 $\pm$ 5.94	8.48 $\pm$ 7.13	7.29 $\pm$ 5.42
EDSS (mean $\pm$ SD)	2.26 $\pm$ 1.37	2.15 $\pm$ 1.39	2.16 $\pm$ 1.30	2.17 $\pm$ 1.32
EDSS<4 (n(%))	1613 (82.34)	814 (82.64)	143 (86.14)	128 (85.33)
EDSS $\geq$ 4 (n(%))	346 (17.66)	171 (17.36)	23 (13.86)	22 (14.67)
At least one relapse prior to baseline (n(%))	849 (43.56)	352 (35.85)	51 (30.91)	77 (51.68)

Table 17: Baseline characteristics of the study population for the analysis of associations between disability worsening and societal economic costs (N=1959).
Reproduced with layout modifications from Ness et al. (2019a).

4.2.2 Descriptive cost analyses

Total mean annual costs (inclusive DMTs) amounted to 27,958€±10,139€ per patient, which corresponds to quarterly costs of 6929€±2886€, averaged over a mean follow up time of 20.68±6.07 months. Excluding DMTs, overall mean quarterly costs were 2029€±2765€. Indirect costs accounted for 83.7% (1699€±2444€) of resource utilization, followed by direct medical costs (14.59%; 296€±888€) and direct non-medical costs (1.72%; 35€±214€) (Table 18). The proportion of patients claiming resources in two years varied within cost categories: While 97.35% of the patients utilized at least once direct medical resources (without DMTs), 25.54% used resources leading to direct non-medical costs and 73.46% affirmed causing indirect costs.

Female patients utilized slightly more resources compared to males. Total societal costs increased with age from an average of 984€±1922€ for patients younger than 30 years to 2519€±2933€ for patients older than 50 years significant ($p<0.05$), mainly due to higher expenditures for indirect and direct medical costs.

A significant monetary difference occurred in all cost categories between patients with mild (1707€±2536€) and moderate disability (3556€±3254€) ($p<0.001$). This was mainly due differences in indirect costs (3010€±2877€ (EDSS <4) vs. 1423€±2248€ (EDSS ≥4)). PIRA patients consistently incurred higher costs (2391€±2897€) compared to stable patients achieving DAF (1703€±2489€) ($p<0.001$). RAW status led to highest total resource utilization (2812€±3426€) but did not statistically significant differ from PIRA.

	Total costs (excl. DMTs)	Direct medical costs	Direct non-medical costs	Indirect costs	DMT costs
	Mean±SD	Mean±SD	Mean±SD	Mean±SD	Mean±SD
Whole Sample	2029 ±2765	296 ±888	35 ±214	1699 ±2444	4900 ±746
Age					
18-29 years	984 ±1922	210 ±655	5 ±41	770 ±1707	4939 ±761
30-49 years	2059 ±2790	300 ±924	36 ±216	1723 ±2453	4933 ±763
50+ years	2520 ±2933	330 ±897	49 ±258	2140 ±2618	4789 ±678
Gender					
Male	1984 ±2811	268 ±829	25 ±146	1691 ±2509	4937 ±753
Female	2046 ±2746	307 ±910	39 ±236	1701 ±2417	4885 ±742
EDSS					
<4	1707 ±2536	261 ±816	22 ±171	1423 ±2248	4865 ±731
≥4	3566 ±3254	461 ±1157	95 ±348	3010 ±2877	5064 ±790
Disability					
DAF	1703 ±2489	201 ±609	26 ±201	1475 ±2284	4916 ±739
RWW	2316 ±2837	397 ±1014	31 ±124	1887 ±2488	4877 ±764
PIRA	2392 ±2897	329 ±956	55 ±278	2008 ±2614	4773 ±700
RAW	2812 ±3426	549 ±1499	76 ±365	2188 ±2704	4861 ±745
Others	2440 ±3311	421 ±1217	45 ±181	1978 ±2883	5116 ±764

Table 18: Mean±SD quarterly cost in Euro (€) within two years of study time (N=1959).
Reproduced with layout modifications from Ness et al. (2019a).

4.2.3 Associations between disability worsening and costs

PIRA caused on average 29% higher total costs compared to patients achieving DAF status after controlling for the effects of increased disability, age and female gender (IRR=1.29, 95% CI 1.08-1.53, $p<0.05$) (Figure 8). The effect of PIRA was most striking in direct medical costs. These were factored by 1.48 (95% CI: 1.13-1.95, $p<0.01$) compared to their stable counterparts. Indirect costs increased by 25% (95% CI: 1.03-1.52, $p<0.05$).

Figure 8: Effects (IRR, 95% CI) of PIRA and RAW on the quarterly economic costs compared to DAF.

Results were adjusted for age, gender, EDSS, time and data source.

Abbreviation: Progression independent of relapse activity (PIRA); Relapse associated worsening (RAW); Disease activity free (DAF).

*: $p < 0.05$; **: $p < 0.01$

Reproduced without modifications from Ness et al. (2019a).

In comparison to PIRA, the effect of RAW was stronger in all cost categories. Total costs increased by factor 1.56 (95% CI: 1.30-1.87, $p < 0.001$) in comparison to stable patients. Direct medical costs increased by factor 2.25 (95% CI: 1.72-2.94, $p < 0.001$) and indirect costs by factor 1.43 (95% CI: 1.17-1.75, $p < 0.001$) compared to patients achieving DAF. For both, PIRA and RAW, the increases in direct medical costs were in particular due to higher rates of inpatient and outpatient hospitalizations (Figure 9).

These findings support the first hypothesis that societal economic costs are significantly higher in patients with disability worsening compared to patients being free of disease activity.

Figure 9: Comparison of patients with in- and outpatient hospitalizations within two years. Reproduced without modifications from Ness et al. (2019a).

4.3 Societal economic impact of relapses

4.3.1 Study population

Out of 1959 principal eligible subjects, we excluded 77 persons who had less than three visits so that 1882 patients were included in the analysis. Of these, 607 (32%) presented at least one relapse, while 1275 (68%) patients were free of relapse activity during the two-year study period (Table 20). Compared with relapse inactive patients, relapse active patients were younger (40.46 ± 9.98 vs. 42.09 ± 10.01 , $p < 0.01$), more often female (76% vs. 70%, $p < 0.01$), more severely disabled (2.41 ± 1.30 and 2.17 ± 1.39 , $p < 0.001$) and more often employed (64% vs. 62%, $p = 0.712$). The cohorts did not differ in BMI with average scores of 25.45 ± 5.14 and 25.66 ± 5.11 between the relapse inactive and relapse active population, respectively. After PS matching, 597 out of 607 patients per cohort were retained for further analysis. The PS matching resulted in an alignment of the relapse inactive to the relapse active population (Figure 10 and Figure 11). There were no statistically significant differences between the matched cohorts (Table 20). One quarter (25%) of the relapse active patients presented with RAW.

Baseline Characteristics	Selected (N=1882)	Not selected (N=77)
Female gender (n (%))	1360 (72.30)	65 (85.53)
Age (mean $\pm$ SD)	41.56 $\pm$ 10.03	42.73 $\pm$ 10.40
Duration of disease since diagnosis, years		
mean $\pm$ SD	7.29 $\pm$ 5.95	7.55 $\pm$ 5.83
median (IQR)	6.00 (3.00, 12.00)	6.00 (3.00, 11.00)
EDSS score		
mean $\pm$ SD	2.25 $\pm$ 1.36	2.53 $\pm$ 1.40
median (IQR)	2.00 (1.00, 3.50)	2.50 (1.50, 4.00)

Table 19: Comparison of baseline characteristics of included and excluded patients for the relapse costs analyses.

	Unmatched population			Matched set		
Baseline Characteristics	No Relapse (n=1275)	Relapsed (n=607)	P-Value	No Relapse (n=597)	Relapsed (n=597)	P-Value
Socio-demographic						
Female gender (n (%))	896 (70)	464 (76)	0.006a	459 (77)	454 (76)	0.733 ^a
Age (mean±SD)	42.09 ±10.01	40.46 ±9.98	0.002b	41.14 ± 9.74	40.68 ±9.85	0.431 ^b
18-29 years (n(%))	492 (39)	266 (44)		248 (42)	257 (43)	
30-49 years (n(%))	464 (36)	225 (37)		215 (36)	224 (38)	
50 years and older (n(%))	319 (25)	116 (19)		134 (22)	116 (19)	
BMI (mean±SD)	25.45 ±5.14	25.13 ±4.97	0.239b	25.23 ±5.02	25.11 ±4.97	0.050 ^b
Under-/normal weight (BMI <25) (n(%))	687 (56)	340 (58)		312 (54)	335 (58)	
Overweight (BMI 25-<30) (n(%))	334 (27)	159 (27)		164 (29)	157 (27)	
Obesity (BMI 30+) (n(%))	202 (17)	92 (16)		99 (17)	89 (15)	
Disease Characteristics						
Disease duration (mean±SD)	7.53 ±6.16	6.78 ±5.47	0.027b	6.82 ±5.60	6.83 ±5.47	0.957 ^b
EDSS (mean±SD; median)	2.17 ±1.39; 2	2.41 ±1.3; 2	<0.001b	2.33 ±1.31; 2	2.40 ±1.31; 2	0.503 ^b
0-1.5 (n(%))	519 (41)	185 (30)		179 (30)	185 (31)	
2.0-3.5 (n(%))	535 (42)	320 (53)		316 (53)	312 (52)	
4-5.5 (n(%))	221 (17)	102 (17)		102 (17)	100 (17)	
Relapse associated worsening						<0.001 ^a
Yes (n(%))	0 (0)	150 (25)		0 (0)	149 (25)	
No (n(%))	1275 (100)	457 (75)		597 (100)	448 (75)	
Therapy						0.77
BRACE (n(%))	958 (75)	493 (81)		479 (80)	483 (81)	
Fingolimod (n(%))	317 (25)	114 (19)		118 (20)	114 (19)	

Table 20: Characteristics of the study population for the analysis of relapse associated societal economic costs (N=1882).

^a P-values were derived from Chi Square test.

^b P-values were derived from Mann-Whitney U test.

Reproduced with modifications from Ness et al. (2020b).

Figure 10: Love plot summarizing the covariate balance before and after PS matching. All covariates were balanced using PS matching. The vertical line represents the threshold within balance is considered to be achieved. Reproduced without modifications from Ness et al. (2020b).

Figure 11: Histogram comparing the density of PS before and after matching. Reproduced without modifications from Ness et al. (2020b).

4.3.2 Relapse associated costs

The relapse active cohort incurred significantly higher costs during relapses compared to the inactive cohort ($p < 0.001$) (Figure 12). More specifically, total relapse costs amounted to 1253€ consisting of 556€ (44%) direct and 697€ (56%) indirect costs. Differences in direct costs arose essentially from inpatient treatment (relapse inactive: 93€ vs. relapse active: 497€) day admissions (relapse inactive: 9€ vs. relapse active: 75€) physician consultations (relapse inactive: 152€ vs. relapse active: 232€) while costs of absenteeism (relapse inactive: 137€ vs. relapse active: 750€) were mainly responsible for indirect relapse costs.

Figure 12: Resource costs (3 months) for relapse inactive (n=597) and relapse active (RAW: n=149; not RAW n=448) patients (before, during and after relapse). Relapse costs, defined as the difference in inactive and active patients during relapses, were significant (Δ 1253€, $p<0.001$). A significant increase was also observed between prior relapse and during relapse quarter. Abbreviations: prior relapse= three months before relapse quarter; during relapse= relapse quarter; post relapse= three months after relapse quarter; RAW= relapse associated worsening Reproduced with modifications from Ness et al. (2020b).

There were noticeable patterns in relapse costs regarding disease and patient characteristics (Table 21). Total relapse costs were higher in patients with shorter disease duration (<5 years: 1729€ vs. >10 years: 758€; $p<0.001$) and those being mildly disabled (EDSS 0-1.5: 1506€ vs. EDSS 4.5-5.5: 987€; $p<0.001$). In mildly disabled patients and those with a short disease duration, indirect relapse costs (range: 971€-1093€) constantly outweighed direct costs (535€-636€). Men incurred higher total relapse costs compared to women (1451€ vs. 1193€), which is mostly due to higher indirect work related relapse costs (920€ vs. 628€). However, differences were not statistically significant. Furthermore, patients on fingolimod incurred lower relapse costs throughout all cost categories than patients on BRACE therapy (total relapse costs: 1067€ vs. 1296€) with noticeable differences in direct relapse costs (316€ vs. 611€). A detailed breakdown of these costs is provided in Table 22.

	Total Mean quarterly costs in Euro±SD		Direct Mean quarterly costs in Euro±SD		Indirect Mean quarterly costs in Euro±SD	
	Cost without relapse	Relapse costs (Δ)	Cost without relapse	Relapse costs (Δ)	Cost without relapse	Relapse costs (Δ)
Gender						
Male (a)	2003±2014	1451	311±474	531	1692±1897	920
Female (b)	2086±2321	1193	351±723	564	1735±2063	628
Age						
18-29 (a)	1466±1844	1405 ^c	230±432	686 ^{b,c}	1236±1707	718
30-49 (b)	2319±2278	1437	424±705	505 ^a	1895±2055	932
50+ (c)	2777±2604	690 ^a	419±919	382 ^a	2358±2296	308
BMI						
<25 (a)	1880±2213	1029	318±740	465	1562±1963	564
25-30 (b)	2317±2295	1510	378±648	720	1938±2086	789
>30 (c)	2324±2280	1653	386±520	586	1938±2119	1067
Disease Duration						
<5 years (a)	1559±1869	1729 ^{b,c}	298±519	636	1261±1710	1093 ^{b,c}
5-10 years (b)	2435±2362	713 ^a	373±687	463	2062±2140	250 ^a
>10 years (c)	2822±2630	758 ^a	409±928	480	2413±2288	278 ^a
EDSS						
0-1.5 (a)	806±1203	1506 ^{b,c}	137±198	535	669±1137	971 ^{b,c}
2-3.5 (b)	2310±2126	1239 ^a	367±582	566	1943±1968	672 ^a
4-5.5 (c)	3528±2849	987 ^a	625±1181	594	2903±2509	393 ^a
Therapy						
BRACE (a)	2055±2252	1296	331±657	611	1724±2045	685
FINGO (b)	2114±2262	1067	387±739	316	1728±1947	751

Table 21: Total, direct and indirect relapse costs (Δ) stratified by patient and disease characteristics.

Relapse costs (Δ) were defined as the difference in costs between relapse inactive and relapse active patients. Letters indicate significant statistical differences ($p < 0.05$) between subgroups. P-Values were derived from GLM. The GLM included the respective cost category and covariates as well as a variable indicating the relapse status and the interaction of both.

Abbreviations: Body mass index (BMI); Beta interferons and glatiramer acetate (BRACE); Fingolimod (FINGO).

Reproduced with modifications from Ness et al. (2020b).

	Inpatient stays Mean quarterly costs in Euro (SD)		Day admissions Mean quarterly costs in Euro (SD)		Physician consultations Mean quarterly costs in Euro (SD)		Informal care Mean quarterly costs in Euro (SD)		Sick leave Mean quarterly costs in Euro (SD)		Presenteeism Mean quarterly costs in Euro (SD)	
	Cost without relapse	Relapse costs (Δ)	Cost without relapse	Relapse costs (Δ)	Cost without relapse	Relapse costs (Δ)	Cost with- out relapse	Relapse costs (Δ)	Cost with- out relapse	Relapse costs (Δ)	Cost with- out relapse	Relapse costs (Δ)
Gender												
Male (a)	79± 237	385	17± 166	79	154± 264	64	25± 139	14	100± 365	776	878± 1201	105
Female (b)	97± 519	409	7± 46	60	152± 218	85	32± 117	24	149± 612	551	600± 1042	215
Age												
18-29 (a)	50± 283	468	15± 132	102	108± 153	95	18± 95	21	98± 336	645	656± 1049	168
30-49 (b)	126± 456	391	4± 21	52	189± 308	59	40± 150	16	176± 715	644	695± 1098	224
50+ (c)	120± 706	287	7± 45	6	176± 182	94	36± 117	36	149± 625	436	631± 1141	174
BMI												
<25 (a)	121± 614	278	3± 22	76	134± 184	86	20± 89	25	130± 586	420	645± 1053	165
25-30 (b)	50± 173	655	19± 161	38	186± 317	63	51± 183	-7	125± 387	867	680± 1136	318
>30 (c)	80± 254	404	14± 57	79	168± 199	98	31± 92	56	188± 768	850	653± 1077	100
Disease Duration												
<5 years (a)	93± 362	434	7± 48	98	135± 200	89	22± 103	21	133± 520	767	674± 1070	253

5-10 years (b)	86± 420	353	16± 156	59	177± 296	44	32± 130	20	219± 802	417	718± 1190	304
>10 years (c)	103± 697	393	7± 46	-2	164± 200	103	48± 150	25	54± 200	458	576± 997	-60
EDSS												
0-1.5 (a)	24± 104	283	4± 25	128	70± 81	106	2± 12	21	24± 90	608	494± 812	489
2-3.5 (b)	105± 397	450	12± 120	31	157± 196	82	32± 128	22	198± 647	698	847± 1228	42
4-5.5 (c)	176± 879	499	9± 33	53	283± 387	37	73± 183	28	149± 727	301	397± 936	113
Therapy												
BRACE (a)	87± 466	443	9± 96	60	152± 212	93	26± 103	28	141± 596	632	671± 1091	109
FINGO (b)	117± 486	231	10± 55	85	152± 290	24	48± 180	-3	124± 414	475	637± 1074	539

Table 22: Detailed breakdown of relapse costs (Δ) stratified by patient and disease characteristics .

Relapse costs (Δ) were defined as the difference in costs between relapse inactive and relapse active patients. The table lists resources that were increasingly used during relapse periods. Certain resources, such as complementary medicine or direct non-medical investments, are utilized at a lower degree during relapse periods compared to relapse-free periods..

Abbreviations: BMI= Body mass index; BRACE= Beta interferons and glatiramer acetate; EDSS= Expanded disability status scale

Reproduced with modifications from Ness et al. (2020b).

I assessed relapse costs without PS-matching as a sensitivity analysis. The matched and unmatched total relapse cost estimates differed by up to 23% due to the heterogeneity in patient and disease characteristics in the respective populations (Table 23).

	Total Mean quarterly costs in Euro±SD		Direct Mean quarterly costs in Euro±SD		Indirect Mean quarterly costs in Euro±SD	
	Cost without relapse	Relapse costs (Δ)	Cost without relapse	Relapse costs (Δ)	Cost without relapse	Relapse costs (Δ)
Gender						
Male (a)	2001±2150	1453	244±376	599	1758±2031	854
Female (b)	1997±2331	1289	317±648	598	1680±2074	690
Age						
18-29 (a)	1410±1935	1473	221±526	687	1189±1729	786
30-49 (b)	2238±2351	1532	335±566	603	1903±2157	929
50+ (c)	2560±2449	907	352±670	449	2208±2211	457
BMI						
<25 (a)	1740±2091	1175	265±550	522	1475±1899	654
25-30 (b)	2190±2397	1632	308±565	796	1882±2193	836
>30 (c)	2432±2488	1548	393±737	555	2039±2160	993
Disease Duration						
<5 years (a)	1541±2012	1758	258±538	670	1283±1827	1088
5-10 years (b)	2218±2322	930	303±569	533	1915±2105	397
>10 years (c)	2599±2499	981	353±660	536	2246±2247	445
EDSS						
0-1.5 (a)	1006±1506	1305	137±211	535	870±1453	771
2-3.5 (b)	2284±2154	1247	330±500	597	1954±1996	649
4-5.5 (c)	3640±2861	916	582±1050	648	3058±2512	268
Therapy						
BRACE (a)	1915±2253	1441	287±541	655	1628±2048	786
FINGO (b)	2253±2332	929	321±690	382	1932±2085	547

Table 23: Sensitivity analysis for relapse costs (Δ) stratified by patient and disease characteristics.

Relapse costs (Δ) were defined as the difference in costs between relapse inactive and relapse active patients.

Abbreviations: Body mass index (BMI); Beta interferons and glatiramer acetate (BRACE); Fingolimod (FINGO).

Reproduced with modifications from Ness et al. (2020b).

A higher economic burden for patients with short disease duration and those that are mildly disabled was also observed when assessing the increase in resource costs from prior relapse to relapse quarter in relapse active patients. However, relapse associated costs were lower throughout all subgroups in this predisposed active only population.

4.3.3 Longitudinal analysis of relapse associated costs

In the relapse active cohort, costs increased significantly from prior relapse to the relapse period (total 1042€; $p<0.001$). In detail, direct and indirect costs increased by 527€ ($p<0.001$) and 515€ ($p<0.001$), respectively. The increase was strongest for inpatient stays (+364%, 390€; $p<0.001$) and day admissions (+241%, 53€; $p<0.001$) in direct costs, while in indirect costs the increase was higher in costs of absenteeism (+127%, 419€; $p<0.001$) than in those of presenteeism (+10%, 88€; $p=0.002$). Costs mostly decreased in the post relapse quarter close to the initial level. However, in patients with RAW direct costs and costs of absenteeism remained elevated (Figure 13).

The results presented in chapter 4.3 support the hypothesis that young patients and those with a shorter disease duration have higher societal economic relapse costs than older patients and those with longer disease duration.

Figure 13: Quarterly costs (in €) of direct and indirect resource items before during and post relapse.

Abbreviations: Three months before relapse quarter (prior relapse); Relapse quarter (during relapse); Three months after relapse quarter (post relapse); Relapse associated worsening (RAW).

Reproduced with modifications from Ness et al. (2020b).

5 Discussion

The following chapters are excerpts of manuscripts submitted to or published in peer review journals:

- *Chapters 5.2 is based on an article titled “Differentiating societal costs of disability worsening in multiple sclerosis” by Ness et al. (2019a).*
- *Chapter 5.3 is based on an article titled “Real-World Evidence on the Societal Economic Relapse Costs in Patients with Multiple Sclerosis” by Ness et al. (2020b).*
- *Chapter 5.4 is based on an article titled “Multiple Sclerosis Health Resource Utilization Survey (MS-HRS): Development and Validation Study” by Ness et al. (2020a).*

5.1 Summary of results

There is an urgent need to reduce the societal economic burden in chronic diseases like MS. Therefore, I investigated how relapses and disability worsening affect the societal economic costs. The results showed that disability worsening is highly associated with a substantial societal economic burden. In particular, two-year real-world data revealed that not only RAW but also PIRA were associated with a significantly increase in costs. Hence, I confirmed the hypothesis that societal economic costs are significantly higher in patients with disability worsening compared to patients being free of disease activity. Furthermore, I confirmed the hypothesis that younger patients and those with a shorter disease duration have higher societal economic relapse costs than older patients and those with longer disease duration. This study provides preliminary evidence that delaying disability worsening and preventing relapses should be prioritized in medical research to reduce the societal economic burden in patients with MS.

The analyses provide valuable information about the diverse economic impact of MS and thus might help decision makers to allocate scarce resources. Furthermore, in the context of economic evaluation, the results have the potential to increase the validity of cost-effectiveness assessments of innovative therapies in relapsing MS. Given the large number of available and upcoming MS therapies of varying effectiveness and costs, economic evaluations become even more relevant.

5.2 Societal economic impact of disability worsening

The economic burden of MS including its direct, indirect and intangible costs has been widely examined in cross-sectional analyses (Ernstsson et al., 2016; Naci et al., 2010). However, to the best of my knowledge, no previous studies on societal economic costs used a longitudinal design comparable to mine. I assigned patients to either clinical DAF, PIRA, RAW or RWW through up to five consecutive EDSS and eight relapse assessments and estimated quarterly societal economic costs. I found that RAW and PIRA lead to significantly higher societal economic costs than DAF.

Both, RAW and PIRA led to increased utilization of direct medical resources. Nevertheless, the monetary effect was stronger for RAW. The proportion of participants with inpatient hospitalizations was almost twice as high in RAW compared to PIRA within two years. However, the average number of inpatient days did not differ between the groups. In contrast, both the proportion of patients with outpatient admissions and the average days of outpatient stays in the RAW group were more than twofold compared to the PIRA population. Beyond disease activity, I observed further parameters influencing the economic burden of MS. Higher age was associated with increased costs. Higher age (≥ 50 years) was significantly more frequent in patients with PIRA compared to DAF and RAW, which could be explained by the underlying progressive course that preferentially occurs at later stages of the disease and thus, in older patients (Lublin et al., 2014). Increased mean disease duration for patients with PIRA supports this suggestion. Lastly, total costs were slightly higher in female patients compared to their male counterparts. RAW status was associated with female gender. This observation can be attributed to the predisposition of relapses in women (Kalincik et al., 2013). To receive valid results, I adjusted the multivariate analysis for age, gender, EDSS.

In the present analyses, 985 patients neither showed disability worsening nor relapses (herein defined as DAF), which corresponds to 50.28% of patients. This is in line with a comprehensive DAF analysis within a real-world cohort of 306 mild to moderately disabled MS patients in Germany, in which DAF was achieved by 45% of patients after two years (Huhn et al., 2019). Although the population I analyzed consisted exclusively of patients with RRMS, a substantial proportion of accumulated disability was due to a progressive clinical course: 8.47% of patients in the study had a PIRA event within two years which corresponds to findings from a recent study (7.00%-9.66%) (Kappos et al., 2018b). On the contrary, the incidence of RAW was slightly higher in the investigated population (7.66% vs. 2.0%-4.7%) which might be explained by differences in the methodology in the assessment of RAW events (Kappos et al., 2018b). EDSS baseline scores were

comparable between the subgroups. A large number of recent studies conducting research on PIRA and RAW in a clinical context, supporting the validity and importance of the approach (Kappos et al., 2018a; Kappos et al., 2018b; Koch-Henriksen et al., 2019).

The mean annualized economic burden including DMTs was 26,482€±9107€ for patients with EDSS scores <4. This finding is comparable to results of Flachenecker et al. (2017) who found mean annual costs of 28,200€ (EDSS <4; price level 2015) in a German substudy with self-assessed EDSS and a multinational survey by Karampampa et al. (2012) in which mean annual costs of 25,270€ (price level 2009) were reported for Germany (Flachenecker et al., 2017; Karampampa et al., 2012b). Furthermore, I found a cost ratio of 1.32 between EDSS strata (0-3.5 vs. 4-5.5) while Flachenecker et al. reported a cost ratio of 1.56 (Flachenecker et al., 2017). Taking into account that in Flachenecker EDSS strata of 0-3.5 and 4.0-6.0 were compared, the ratios can be regarded as comparable. I also confirmed that a progressive clinical course increases societal economic costs as it was stated before (Gyllensten et al., 2019). Nevertheless, I found differences in the composition of total costs that depart from those reported in literature (Flachenecker et al., 2017). Particularly noticeable are lower expenses for hospital stays, examinations and informal care in this analysis, which could be explained by the milder disease profile in this analysis.

5.3 Societal economic impact of relapses

Relapse costs have been investigated repeatedly in cross-sectional studies with highly variable methodologies and results ranging from 503€-8862€ (Flachenecker et al., 2017; O'Connell et al., 2014). Most commonly, relapse costs were calculated as the difference between costs of patients with and without relapses without presenting characteristics of these distinct populations (Flachenecker et al., 2017; Hawton and Green, 2016). Less frequently, relapse associated costs were directly enquired in interviews (O'Connell et al., 2014). This methodology can be critically questioned because patients are required to differentiate between routinely utilized resources and relapse related resources. I extended the first mentioned approach by applying a PS matching. Each relapse active patient was assigned to a statistical twin, which was relapse inactive. I reported relapse costs stratified by patient and disease characteristics with estimated relapse costs ranging between 690€ (age: 50+ years) and 1729€ (disease duration: <5 years). The matched and unmatched total relapse cost estimates differed by up to 23% due to the heterogeneity in patient and disease characteristics in the respective populations. The wide range of relapse costs suggests that precision of cost estimates in future economic evaluations in MS will increase if patient and disease characteristics are considered.

Former studies hypothesized that relapses may have a stronger effect in patients with limited permanent disability, however, a systematic investigation of this hypothesizes remained unfulfilled (Kobelt et al., 2006a). With regard to the analysis, I can confirm that patients with a less severe burden of disease (mildly disabled, short disease duration) are more likely to incur high relapse costs predominantly due to higher indirect costs. These patients are more frequently employed resulting in higher costs due to sick leave. The results complement the findings of O'Brien who showed that relapse costs varied with relapse severity (O'Brien et al., 2003).

A recent systematic review found that the most notable increase in relapse costs was associated with inpatient treatments, physician consultations, informal care and sick leave (Naci et al., 2010). The present analyses confirmed these findings.

Inpatient treatment during a relapse might be indicative of a severe relapse (O'Brien et al., 2003; O'Connell et al., 2014). In the investigated population inpatient care costs were lower than previously assumed (Flachenecker et al., 2017). This could partly be explained by the inclusion criteria, which require patients to be treated with DMTs. DMTs can affect the disease course by slowing disease worsening, reducing the frequency and the severity of relapses (Giovannoni et al., 2016; Haas et al., 2019; Sanchirico et al., 2019; Vercellino et al., 2009). As confirmed by the present results, fingolimod has shown to reduce healthcare resource utilization during relapses (Haas et al., 2019; Raimundo et al., 2013).

Despite the increase in resource costs during relapses, I found that resource cost utilization decreases in the post relapse phase. However, costs partially remained above the initial cost level. Interestingly, this was particularly observed in patients with RAW for resources that are notably increased during periods of relapses (inpatient stays, physician consultations, informal care, and sick leave). This might indicate an incomplete recovery with residual disability (Kalincik, 2015). This result is in line with recent findings, showing the high frequency of post-relapse residual disability (Achiron et al., 2018).

5.4 Discussion of the Multiple Sclerosis Health Resource Survey

A prerequisite for the Col analyses was an instrument for resource use measurement with good psychometric properties. Even though Col studies have a long tradition in MS, there were no validated instruments public available (DIRUM, 2019). Therefore, I developed and psychometrically assessed the MS-HRS. The results confirm that the MS-HRS represents a reliable, valid and easy to administer questionnaire providing a holistic assessment of resource utilization for patients with MS. Health-resources were derived from all pathways of pwMS into an adapted health-resource model for MS.

Some very respectable efforts have already been taken to research the health economic footprint of MS (Ernstsson et al., 2016; Karampampa et al., 2012a; Kobelt et al., 2006a; Kobelt et al., 2017b). In Europe, Kobelt et al. repeatedly assessed costs and burden of MS in a cross-sectional survey approach including direct, indirect and intangible costs (Kobelt et al., 2006a; Kobelt et al., 2017b). Concepts and definitions of sub costs may differ over time with respect to scope and style of reporting, for example on intangible costs where a clear line is recommended (Wundes et al., 2010). For the model approach, I gave strong emphasis on the direct and indirect costs as Karampampa et al. and Reese et al. did with their models (Karampampa et al., 2012a; Reese et al., 2011).

I increased the depth of the assessment of indirect costs by adding a quantification of costs due to presenteeism in addition to costs of absenteeism (sick leave, disability pension).

Part of the questionnaire development was a validation process that confirmed reliability and validity in a way that no previous approach did within the MS domain (Figure 14). Kobelt et al. also recognized the need for validation, but a systematic approach beyond the aspects of practicability and face validity remained a task to fulfill (Kobelt et al., 2019b). Even though I did not develop a one-dimensional questionnaire for a latent construct, I was able to demonstrate essential psychometric qualities in the health economic assessment of MS. As of today, a multi-domain open-access database of resource-use questionnaires did not contain a published instrument for MS (DIRUM, 2019). Most of the cost assessments for MS were not developed to be published for general use but for the application in certain cross-sectional frameworks. A very recent analysis of real-life cost outcomes underlined the rising interest in longitudinal assessment of health-resource utilization (Kobelt et al., 2019a).

Figure 14: Development process of the MS-HRS.
Reproduced without modifications from Ness et al. (2020a).

Compared to the results of Flachenecker et al. (2017), the proportion of patients claiming certain health services was slightly lower than in the present population. Particularly noticeable was the lower proportion of patients taking over-the-counter (OTC) medication and investments. This may be explained by the less severe disease progression beyond a RRMS profile. Taking this into account, the average annualized burden of disease lied within the range expected from previous publications (Karampampa et al., 2012a; Kobelt et al., 2006a; Reese et al., 2011).

Whereas Kobelt et al. have recently reported precise price tags for most unit costs, other authors have at least provided an indirect description of the valuation process and its exact results (Karampampa et al., 2012a; Kobelt et al., 2017a). Differences were found for the valuation of sub-costs, like inpatient hospitalization, due to different sources of valuation or different definitions for health-resource units. In any case, a precise and fully transparent reporting of per-unit costs is highly recommended, especially for main cost drivers.

Event-related costing can also be considered of interest but a clear separation from costs being assessed via routine data collection is necessary (Karampampa et al., 2012a). Otherwise, costs may double due to double assessment. I discussed in an expert group whether to collect information both on sick days and on missed working hours as described by Reilly et al (Reilly et al., 1993). Furthermore, asking for sick hours would have

meant switching to a recall period of seven days while the remaining questionnaire covers the last six months. Due to higher usability, I decided to assess sick days only.

Claims data were considered for the assessment of criterion validity. However, they have the disadvantage that only billing-relevant data is available preventing taking a societal perspective. Information on informal care, OTC medication, presenteeism and partially on investments and medical consumables are not recorded. In addition, claims data do not provide the number of physician visits of a patient in Germany. In consequence, claims data were not suitable for the validation process.

Health economic studies often require both clinical and economic data. The MS-HRS can easily be used as part of clinical interventional and non-interventional studies to collect economic data. In a large population of two non-interventional multicenter studies the questionnaire was easy to administer and has good psychometric properties. These characteristics provide the prerequisites for high-quality health economic studies, e.g. cost-effectiveness analyses.

5.5 Generalizability of the study results

The study population represented typical patient characteristics for a clinical RRMS population: patients were about 41 years old with a mean disease duration of 7 years. About three quarter of patients were female. To ensure representativeness, I used data from two large observational, prospective studies in which patients were documented homogeneously across Germany. Representativeness towards a real-world setting was supported by the fact that no explicit exclusion criteria, except contraindications associated with current DMTs, had been stated for the (sub)studies PEARL and PANGAEA. High data quality was further achieved through standardized case report forms and the use of external validation checks at the time of data entry. Therefore, it might be assumed that the study results can be generalized to the wider MS population in Germany. Caution is required when transferring results to other countries in which healthcare systems have different reimbursement and treatment regimes.

5.6 Limitations of the study

For all analyses, I excluded patients with baseline EDSS ≥ 6 to ensure a balanced mild to moderate disabled study population. Corresponding data on severely disabled patients are not yet available but consecutive studies addressing this population are already being conducted (Haase et al., 2018a). The study results are based on patient-reported

data on resource utilization, which may not be completely free of biases. To avoid potential conflicts, data were collected with the validated MS-HRS in a longitudinal format.

Focusing on the disability worsening analyses, definitions of PIRA and RAW are solely based on EDSS, which means that the limitation of the EDSS also applies to the analysis. Particularly noteworthy is the focus on functional mobility and insensitivity on MS-related impairments like cognition. To address this problem, novel composite measures additionally include Nine-Hole Peg Test and Timed 25-Foot Walk to detect composite PIRA and RAW events (Kappos et al., 2018b). Nevertheless, as the definitions are solely based on EDSS, the analyses are less demanding and easier to implement in non-interventional studies.

A small proportion of patients utilized direct non-medical resources with a high variability of costs, which might result from the low disability level in the study population. In consequence, confidence intervals were conspicuously wide. Former analyses have shown that utilization of direct non-medical resources radically increase in more severely disabled patients (Flachenecker et al., 2017). To better understand the effect of PIRA and RAW on direct non-medical costs, further analyses in more disabled populations may be helpful.

In the relapse cost analyses, I compensated for variation between patients with and without relapses by applying a PS matching. PS matching is a powerful technique to reduce imbalances in observable outcomes between populations. However, it cannot address unknown or unmeasured differences. Furthermore, I did not include corticosteroid medication costs, which could have led to an underestimation of relapse costs.

The MS-HRS is a patient-centered questionnaire and therefore subject to notable limitations. Reliable recall periods of health-resource use are time-limited, especially in patients that are cognitively more affected by the disease. Furthermore, patients may not want to disclose socioeconomic and health economic information because it is considered too confidential. Further resource items were discussed to include into the analysis like crowd-funded therapies or cost-related voluntary work loss. However, these costs are unlikely to affect the economic burden on a societal level. In future studies, price tags for cost components have to be updated and adapted to local levels. For patients with progressive MS and severely disabled patients, further studies have to confirm the given psychometric properties.

6 Conclusion

For the case of MS, disability worsening and relapses might be focused in medical research to reduce the societal economic burden. Disability worsening is associated with a tremendous societal economic burden in patients with MS. Both PIRA and RAW lead to significant increases in direct medical and indirect work-related costs compared to patients achieving DAF status. Accordingly, it is highly important from a societal perspective to delay or even prevent the transition into the progressive phase of MS. That is another argument for treating MS patients early with appropriate DMTs.

Furthermore, relapses significantly increase the societal economic burden of MS. Particularly young patients and those with a recent diagnosis incur high relapse costs. The prevention of relapses in these patients should be a particular focus to reduce the economic burden. In addition, it should be ensured that patients are diagnosed quickly to get the best available treatment.

Further studies on the incidence and prevalence of disability worsening and relapses are demanded to assess financial risks for health systems and optimizing health service planning in patients with MS.

As part of the study, I psychometrically assessed and published the MS-HRS. The MS-HRS provides a holistic cross-sectional and longitudinal assessment of resource utilization in patients with MS in the German health care setting. Further country-specific cost-weights will facilitate the transparent estimation of MS-related costs across multiple regions.

7 Summary

7.1 English version

Background

Multiple sclerosis (MS) is a chronic, incurable disease of the central nervous system (CNS) and a potentially severe cause of neurological disability throughout adult life. Typically diagnosed between the ages of 20 and 40, the disease is associated with a severely impaired health-related quality of life and a high societal economic impact.

About 85% of patients are initially diagnosed with relapsing-remitting MS (RRMS), which is characterized by the unpredictable occurrence of relapses. Clinically, a relapse is defined as a worsening or appearance of new neurological symptoms followed by a period of partial or complete recovery (relapse without worsening, RWW). Up to 80% of patients transit to the secondary progressive course (SPMS) within 20 years. SPMS is marked by an accumulation of irreversible disability either resulting from relapse-associated worsening (RAW) or progression independent of relapse activity (PIRA).

Even though health economic assessments are well established in MS, implications of relapses and disability worsening on societal economic costs have scarcely been studied. Regarding disability worsening it is known that annual costs increase with increasing disability from 21,174–28,200€ (EDSS 0–3) to 39,923–44,000€ (EDSS 4–6.5) (in 2015 values). However, the impact of PIRA and RAW on societal costs remains not well understood.

Research to date has shown that the societal economic burden of MS increases in periods of relapses with most recent estimates at 2468€ per relapse (in 2015 values). The results of relapse cost analyses are inconsistent between studies since relapses and associated costs can vary across and within individuals depending on disease severity, disease duration, age, gender or therapy. However, a comprehensive test of this hypothesis is still needed.

Research aims

I investigated the societal economic costs of disability worsening and relapses in patients with MS. First, I determined the societal economic costs of patients achieving disease activity free status (DAF, neither relapses nor disability worsening) and compared them with those having PIRA and RAW events. Second, I estimated direct and indirect relapse costs and determined differences in costs related to patient characteristics. Furthermore, I described the pharmacoeconomic impact during the relapse follow-up.

Methods

Two-year clinical and pharmacoeconomic data were derived from two single arm, prospective, multicenter, non-interventional long term studies conducted in Germany applying the Multiple Sclerosis Health Resource Survey (MS-HRS). The MS-HRS estimates the MS-associated direct and indirect costs from a societal perspective using the most accurate form of bottom-up microcosting. Costs are presented per quarter in 2011 Euros. PIRA and RAW events were indicated through a roving confirmed disability worsening (CDW) measure and corresponding relapse assessments. CDW is based on a predefined increase on the Expanded Disability Status Scale (EDSS) sustained over at least 6 months.

The effects of PIRA and RAW on total, direct medical, direct non-medical and indirect societal economic costs are adjusted for age, gender and EDSS. Due to the right skewness of cost data, I applied a Generalized Linear Mixed Model (GLMM) with a negative binomial distribution. Patients achieving DAF were assigned as the reference group.

I calculated relapse costs as the difference in quarterly costs between propensity score (PS) matched patients with and without relapses (1:1 ratio). Statistical differences in relapse costs between patient and disease strata were derived from a Generalized Linear Model (GLM) with a negative binomial distribution. For relapse active patients, I additionally calculated the difference between quarterly costs prior- and during relapse and determined costs in the post-relapse quarter.

Results

Overall, 1959 patients were analyzed with typical patient characteristics for a clinical RRMS population. Patients reported an average age of 41.62 ± 10.04 years and were mostly female (73%). The mean disease duration was 7.30 ± 5.95 with a mean baseline EDSS value of 2.26 ± 1.37 .

Total mean quarterly societal economic costs including disease-modifying therapies (DMTs) were $6929\text{€} \pm 2886\text{€}$ per patient averaged over two years. Excluding DMTs, patients achieving DAF status had total mean quarterly costs of $1703\text{€} \pm 2489\text{€}$. PIRA caused 29% (IRR: 1.29; 95% CI 1.06–1.50, $p < 0.05$) higher total costs compared to DAF. On the contrary, RAW increased total costs by factor 1.56 (95% CI 1.30–1.87, $p < 0.001$). The effect of PIRA and RAW was striking for direct medical costs which increased by factor 1.48 (95% CI 1.13–1.95, $p < 0.01$) and 2.25 (95% CI 1.72–2.94, $p < 0.001$), respectively.

In the relapse cost analyses, a total of 1882 patients were included encompassing 607 (32%) relapse-active patients. After PS-matching, 597 active and inactive patients were

retained. Relapse costs ranged between 690€ (age: 50+ years) and 1729€ (disease duration: <5 years). In mildly disabled and recently diagnosed patients, indirect relapse costs (range: 971€-1093€) constantly outweighed direct costs (range: 535€-636€). The increase from prior- to relapse quarter was strongest for inpatient stays (+364%, 390€; $p<0.001$), day admissions (+241%, 53€; $p<0.001$) and absenteeism (+127%, 419€; $p<0.001$). In the post-relapse quarter direct costs and costs of absenteeism remained elevated for patients with relapse associated worsening.

Conclusion

For the case of MS, disease activity should be a focus in medical research to reduce the societal economic burden of the disease. Disability worsening is associated with significantly higher societal economic costs compared to DAF status in patients with MS. Accordingly, it is highly important from a societal perspective to delay or even prevent transition into the progressive phase of MS. That is another argument for treating MS patients early with appropriate DMTs. Furthermore, relapses significantly increase the societal economic burden of MS. From a health economic societal perspective, the prevention of relapses in young patients and those with a recent diagnosis should be a particular focus.

7.2 German version

Hintergrund

Multiple Sklerose (MS) ist eine chronische, unheilbare Krankheit des zentralen Nervensystems (ZNS) und eine potenziell schwerwiegende Ursache für neurologische Behinderungen im Erwachsenenalter. Typischerweise wird die Krankheit im Alter zwischen 20 und 40 Jahren diagnostiziert und ist mit einer stark beeinträchtigten gesundheitlichen Lebensqualität und hohen ökonomischen Auswirkungen für die Gesellschaft verbunden. Bei etwa 85% der Patienten wird zunächst eine schubförmig-remittierende MS (RRMS) diagnostiziert, die durch das unvorhersehbare Auftreten von Schüben gekennzeichnet ist. Klinisch ist ein Schub als Verschlechterung oder Auftreten neuer neurologischer Symptome definiert, die von einer Periode der teilweisen oder vollständigen Rückbildung (RWW) gefolgt ist. Bis zu 80% der Patienten gehen innerhalb von 20 Jahren in den sekundär progredienten Verlauf (SPMS) über. Dieser ist durch eine Akkumulation von irreversiblen Behinderungen gekennzeichnet, die entweder aus einer schubassoziierten Verschlechterung (RAW) oder einer schubunabhängigen Progression (PIRA) resultieren.

Obwohl gesundheitsökonomische Analysen bei MS etabliert sind, wurden die Auswirkungen von Schüben und Behinderungsprogression auf die sozioökonomischen Kosten bisher wenig untersucht. Hinsichtlich des Behinderungsstatus ist bekannt, dass die jährlichen Kosten mit zunehmender Einschränkung von 21.174-28.200€ (EDSS 0-3) auf 39.923-44.000€ (EDSS 4-6,5) steigen (in Beträgen des Jahres 2015). Der Einfluss von PIRA und RAW auf die sozioökonomischen Kosten ist jedoch unklar.

Darüber hinaus hat die bisherige Forschung gezeigt, dass die sozioökonomische Belastung durch MS während der Schubperiode steigt. Die aktuellsten Schätzungen liegen bei 2468 € pro Schub (in Beträgen des Jahres 2015). Jedoch schwanken die Ergebnisse, da Schübe und die damit verbundenen Kosten je nach Schwere der Krankheit, der Krankheitsdauer, dem Alter, dem Geschlecht oder der Therapie bei ein und demselben Patienten und zwischen Patienten variieren können. Diese Hypothese bedarf jedoch noch einer genaueren Untersuchung.

Forschungsziel

Ziel der vorliegenden Arbeit ist es, die sozioökonomischen Kosten der Behinderungsprogression und der Schübe bei Patienten mit MS zu untersuchen. Zuerst werden die sozioökonomischen Kosten für Patienten geschätzt, die frei von Krankheitsaktivität (DAF) sind. Diese werden anschließend mit den Kosten von Patienten verglichen, welche

Krankheitsaktivität in Form von PIRA und RAW vorweisen. In einem zweiten Schritt werden die direkten und indirekten Schubkosten analysiert und Unterschiede in Bezug auf Patienten- und Krankheitseigenschaften aufgezeigt. Im letzten Schritt wird der ökonomische Einfluss von Schüben in der Post-Phase des Schubes analysiert.

Methodik

Datengrundlage bilden klinische und pharmakoökonomische Daten aus zwei einarmigen, prospektiven, multizentrischen, nicht-interventionellen, deutschen Langzeitstudien, in denen der Multiple Sclerosis Health Resource Survey (MS-HRS) verwendet wurde. Der MS-HRS ermittelt die MS-assoziierten direkten und indirekten Kosten aus gesellschaftlicher Perspektive mit hoher Genauigkeit (Bottom-up Microcosting). Die Kosten werden per Quartal in Preisen des Jahres 2011 berichtet.

PIRA und RAW wurden auf Grundlage bestätigter Behinderungsprogression (CDW) und zugehörigen Schubevaluationen identifiziert. CDW basiert auf einem vordefinierten Anstieg auf der Expanded Disability Status Scale (EDSS), der nach 6 Monaten bestätigt wird.

Der Effekt von PIRA und RAW auf die totalen, direkten medizinischen, direkten nicht-medizinischen sowie indirekten sozioökonomischen Kosten wurde für das Alter, Geschlecht und EDSS-Wert der Patienten adjustiert. Aufgrund der rechtsschiefen Verteilung der Kosten wurden die Effekte mit einem generalisierten gemischten linearen Model (GLMM) geschätzt. Patienten, die frei von Krankheitsaktivität sind, wurden als Referenzgruppe verwendet.

Schubkosten wurden als Differenz der vierteljährlichen Kosten zwischen gematchten Patienten (Propensity Score Matching) mit und ohne Schub berechnet. Die statistischen Unterschiede der Schubkosten zwischen Patienten- und Krankheitseigenschaften wurden mit einem generalisierten linearen Model (GLM) mit einer negativen Binomialverteilung berechnet. Zusätzliche wurden für die schubaktiven Patienten die Differenz der Kosten zwischen dem Prä-Schubquartal und Schubquartal sowie die Kosten in dem Post-Schubquartal ermittelt.

Ergebnisse

Insgesamt wurden 1959 Patienten mit RRMS-typischen Patientencharakteristika analysiert: Die Patienten waren im Durchschnitt $41,62 \pm 10,04$ Jahre alt und überwiegend weiblich (73%). Im Mittel betrug die Krankheitsdauer $7,30 \pm 5,95$ Jahre und der durchschnittliche EDSS-Wert lag bei $2,26 \pm 1,37$ Punkten.

Die mittleren vierteljährlichen sozioökonomischen Kosten einschließlich krankheitsmodifizierender Therapien (DMTs) betrugen 6929€±2886€ pro Patient im Zweijahresdurchschnitt. Patienten, die keine Krankheitsaktivität zeigten, verursachten durchschnittliche Gesamtkosten von 1703€±2489€ (exkl. DMTs) pro Quartal. PIRA bewirkte eine signifikante Steigerung der totalen Kosten um durchschnittlich 29% (IRR: 1,29; 95% CI 1,06-1,50, $p<0,05$). Hingegen führte RAW zu einer Steigerung der totalen Kosten um den Faktor 1,56 (95% CI 1,30-1,87, $p<0,001$). Besonders tragend war der Effekt von PIRA und RAW bei den direkten medizinischen Kosten, die um den Faktor 1,48 (95% CI 1,13-1,95, $p<0,01$) bzw. 2,25 (95% CI 1,72-2,94, $p<0,001$) stiegen.

In die Schubkostenanalyse wurden insgesamt 1882 Patienten einbezogen, von denen 607 (32%) schubaktiv waren. Nach dem PS-Matching wurden 597 aktive und inaktive Patienten für die weitere Analyse beibehalten. Die totalen Schubkosten lagen zwischen 690€ (Alter: 50+ Jahre) und 1729€ (Krankheitsdauer: <5 Jahre). Bei minimal bis mäßiggradiger Behinderung und Patienten mit kurzer Erkrankungsdauer überstiegen die indirekten Schubkosten (Spannweite: 971€-1093€) stets die direkten Schubkosten (Spannweite: 535€-636€). Der Kostenanstieg vom Prä-Schubquartal zum Schubquartal war am stärksten bei stationären Aufenthalten (+364%, 390€; $p<0,001$), ambulanten Aufenthalten (+241%, 53€; $p<0,001$) sowie Krankheitstagen (+127%, 419€; $p<0,001$). Bei Patienten mit RAW blieben die direkten Kosten sowie Kosten für Krankheitstage erhöht.

Fazit

Um die wirtschaftliche Belastung der MS zu verringern, sollte in der medizinischen Forschung die Krankheitsaktivität fokussiert werden. Behinderungsprogression führt zu signifikant erhöhten sozioökonomischen Kosten im Vergleich zu Patienten, die frei von Krankheitsaktivität sind. Dementsprechend ist es aus sozioökonomischer Perspektive besonders wichtig, den Übergang in die progressive Phase der MS zu verzögern oder sogar zu verhindern. Dies ist ein weiteres Argument für eine frühzeitige Behandlung mit geeigneten DMTs. Weiterhin erhöhen Schübe die sozioökonomische Belastung erheblich. Eine besondere Relevanz gilt Schüben bei jungen Patienten und solchen Patienten mit kurzer Erkrankungsdauer.

Bibliography

- Achiron A, Sarova-Pinhas I, Magalashvili D, Stern Y, Gal A, Dolev M, Menascu S, Harari G, Gurevich M. 2018. Residual disability after severe relapse in people with multiple sclerosis treated with disease-modifying therapy. *Mult Scler* 25(13):1746–1753.
- Aharoni R. 2013. The mechanism of action of glatiramer acetate in multiple sclerosis and beyond. *Autoimmun Rev* 12(5):543-553.
- Alsop J, Medin J, Cornelissen C, Vormfelde SV, Ziemssen T. 2017. Two studies in one: A propensity-score-matched comparison of fingolimod versus interferons and glatiramer acetate using real-world data from the independent German studies, PANGAEA and PEARL. *PLoS One* 12(5):e0173353.
- Amato MP, Derfuss T, Hemmer B, Liblau R, Montalban X, Soelberg Sørensen P, Miller DH. 2017. Environmental modifiable risk factors for multiple sclerosis: Report from the 2016ECTRIMS focused workshop. *Mult Scler* 24(5):590-603.
- Ascherio A, Munger KL. 2016. Epidemiology of Multiple Sclerosis: From Risk Factors to Prevention-An Update. *Semin Neurol* 36(2):103-114.
- Atkinson MJ, Kumar R, Cappelleri JC, Hass SL. 2005. Hierarchical construct validity of the treatment satisfaction questionnaire for medication (TSQM version II) among outpatient pharmacy consumers. *Value Health* 8(Suppl_1):S9-S24.
- Atkinson MJ, Sinha A, Hass SL, Colman SS, Kumar RN, Brod M, Rowland CR. 2004. Validation of a general measure of treatment satisfaction, the Treatment Satisfaction Questionnaire for Medication (TSQM), using a national panel study of chronic disease. *Health Qual Life Outcomes* 2(12).
- Beecham J, Knapp M. 1992. Costing psychiatric interventions. In: Thornicroft G, Brewin CR, Wing J (Eds) *Measuring mental health needs*. Royal College of Psychiatrists, London. p. 179-190.
- Bock JO, Brettschneider C, Seidl H, Bowles D, Holle R, Greiner W, König HH. 2015. Calculation of standardised unit costs from a societal perspective for health economic evaluation. *Gesundheitswesen* 77(1):53-61.
- BPB. Bundeszentrale für politische Bildung. 2011. Geleistete Arbeitsstunden je Erwerbstätigem pro Jahr, Produktivität je Erwerbstätigenstunde, 1970-2012 [accessed: 15.01.2020] URL: <http://www.bpb.de/nachschlagen/zahlen-und-fakten/soziale-situation-in-deutschland/61711/arbeitszeit>.
- Breuer J, Loser K, Mykicky N, Wiendl H, Schwab N. 2019. Does the environment influence multiple sclerosis pathogenesis via UVB light and/or induction of vitamin D? *J Neuroimmunol* 329:1-8.
- Byford S, Leese M, Knapp M, Seivewright H, Cameron S, Jones V, Davidson K, Tyrer P. 2007. Comparison of alternative methods of collection of service use data for the economic evaluation of health care interventions. *Health Econ* 16(5):531-536.
- Calabresi PA, Radue EW, Goodin D, Jeffery D, Rammohan KW, Reder AT, Vollmer T, Agius MA, Kappos L, Stites T, Li B, Capiello L, von Rosenstiel P, Lublin FD.

2014. Safety and efficacy of fingolimod in patients with relapsing-remitting multiple sclerosis (FREEDOMS II): a double-blind, randomised, placebo-controlled, phase 3 trial. *Lancet Neurol* 13(6):545-556.
- Chen J, Taylor B, Palmer AJ, Kirk-Brown A, van Dijk P, Simpson S, Blizzard L, van der Mei I. 2018. Estimating MS-related work productivity loss and factors associated with work productivity loss in a representative Australian sample of people with multiple sclerosis. *Mult Scler* 25(7):994-1004.
- Chrüzander C, Tinghog P, Ytterberg C, Widen Holmqvist L, Alexanderson K, Hillert J, Johansson S. 2016. Longitudinal changes in sickness absence and disability pension, and associations between disability pension and disease-specific and contextual factors and functioning, in people with multiple sclerosis. *J Neurol Sci* 367:319-325.
- Chun J, Hartung HP. 2010. Mechanism of action of oral fingolimod (FTY720) in multiple sclerosis. *Clin Neuropharmacol* 33(2):91-101.
- Cohen JA, Barkhof F, Comi G, Hartung H-P, Khatri BO, Montalban X, Pelletier J, Capra R, Gallo P, Izquierdo G, Tiel-Wilck K, de Vera A, Jin J, Stites T, Wu S, Aradhye S, Kappos L. 2010. Oral Fingolimod or Intramuscular Interferon for Relapsing Multiple Sclerosis. *N Engl J Med* 362(5):402-415.
- Cohen JA, Reingold SC, Polman CH, Wolinsky JS. 2012. Disability outcome measures in multiple sclerosis clinical trials: current status and future prospects. *Lancet Neurol* 11(5):467-76.
- Compston A, Coles A. 2008. Multiple sclerosis. *Lancet* 372:1502-1517.
- Confavreux C, Aimard G, Devic M. 1980. Course and prognosis of multiple sclerosis assessed by the computerized data processing of 349 patients. *Brain* 103(2):281-300.
- Correale J, Gaitan MI, Ysraelit MC, Fiol MP. 2017. Progressive multiple sclerosis: from pathogenic mechanisms to treatment. *Brain* 140(3):527-546.
- Cunnam L, Sinanovic E, Ramma L, Foster N, Berrie L, Stevens W, Molapo S, Marokane P, McCarthy K, Churchyard G, Vassall A. 2016. Using Top-down and Bottom-up Costing Approaches in LMICs: The Case for Using Both to Assess the Incremental Costs of New Technologies at Scale. *Health Econ* 25(Suppl_1):S53-S66.
- D'Amico E, Haase R, Ziemssen T. 2019. Review: Patient-reported outcomes in multiple sclerosis care. *Mult Scler Relat Disord* 33:61-66.
- Dagenais S, Caro J, Haldeman S. 2008. A systematic review of low back pain cost of illness studies in the United States and internationally. *Spine J* 8(1):8-20.
- De Angelis F, John NA, Brownlee WJ. 2018. Disease-modifying therapies for multiple sclerosis. *BMJ* 363:k4674.
- Dendrou CA, Fugger L, Friese MA. 2015. Immunopathology of multiple sclerosis. *Nat Rev Immunol* 15:545.

- Derfuss T, Ontaneda D, Nicholas J, Meng X, Hawker K. 2016. Relapse rates in patients with multiple sclerosis treated with fingolimod: Subgroup analyses of pooled data from three phase 3 trials. *Mult Scler Relat Disord* 8:124-130.
- DeVon HA, Block ME, Moyle-Wright P, Ernst DM, Hayden SJ, Lazzara DJ, Savoy SM, Kostas-Polston E. 2007. A psychometric toolbox for testing validity and reliability. *J Nurs Scholarsh* 39(2):155-164.
- DIRUM. Database of Instruments for Resource Use Measurement. 2019. Database of Instruments for Resource Use Measurement - All Instruments [accessed: 15.01.2020] URL: <http://www.dirum.org/instruments/all>.
- Doward LC, McKenna SP, Meads DM, Twiss J, Eckert BJ. 2009. The development of patient-reported outcome indices for multiple sclerosis (PRIMUS). *Mult Scler* 15(9):1092-1102.
- Drost EA. 2011. Validity and reliability in social science research. *Educ Res Perspect* 38(1):105.
- Drummond M. 1992. Cost-of-illness studies. *Pharmacoeconomics* 2(1):1-4.
- Drummond MF. 2015. Methods for the economic evaluation of health care programmes. 4. Ed. Oxford University Press, Oxford.
- Ernstsson O, Gyllenstein H, Alexanderson K, Tinghog P, Friberg E, Norlund A. 2016. Cost of Illness of Multiple Sclerosis - A Systematic Review. *PLoS One* 11(7):e0159129.
- Fasbender P, Kölmel HW. 2008. Incidence of Multiple Sclerosis in the Urban Area of Erfurt, Thuringia, Germany. *Neuroepidemiology* 30(3):147-151.
- Fisk JD, Brown MG, Sketris IS, Metz LM, Murray TJ, Stadnyk KJ. 2005. A comparison of health utility measures for the evaluation of multiple sclerosis treatments. *J Neurol Neurosurg Psychiatry* 76(1):58-63.
- Flachenecker P, Kobelt G, Berg J, Capsa D, Gannedahl M, European Multiple Sclerosis P. 2017. New insights into the burden and costs of multiple sclerosis in Europe: Results for Germany. *Mult Scler* 23(Suppl_2):S78-S90.
- Fowler FJ. 1995. Improving survey questions: Design and evaluation. Sage, Thousand Oaks.
- Fowler J, Floyd J. 1993. Survey research methods. In: Bickman L, Rog D (Eds) *Applied social research methods series*. Sage, Thousand Oaks. p. 1-156.
- Franklin M, Thorn J. 2019. Self-reported and routinely collected electronic healthcare resource-use data for trial-based economic evaluations: the current state of play in England and considerations for the future. *BMC Med Res Methodol* 19(8).
- Gasperini C, Prosperini L, Tintoré M, Sormani MP, Filippi M, Rio J, Palace J, Rocca MA, Ciccarelli O, Barkhof F, Sastre-Garriga J, Vrenken H, Frederiksen JL, Yousry TA, Enzinger C, Rovira A, Kappos L, Pozzilli C, Montalban X, De Stefano N. 2019. Unraveling treatment response in multiple sclerosis: A clinical and MRI challenge. *Neurology* 92(4):180-192.

- Ghasemi N, Razavi S, Nikzad E. 2017. Multiple Sclerosis: Pathogenesis, Symptoms, Diagnoses and Cell-Based Therapy. *Cell journal* 19(1):1-10.
- Giovannoni G, Butzkueven H, Dhib-Jalbut S, Hobart J, Kobelt G, Pepper G, Sormani MP, Thalheim C, Traboulsee A, Vollmer T. 2016. Brain health: time matters in multiple sclerosis. *Mult Scler Relat Disord* 9(Suppl_1):S5-S48.
- Graf von der Schulenburg JM, Claes C, Greiner W, Uber A. 1998. Die deutsche Version des EuroQol-Fragebogens. *Z Gesundh Wiss* 6(1):3-20.
- Graf von der Schulenburg JM, Greiner W, Jost F, Klusen N, Kubin M, Leidl R, Mittendorf T, Rebscher H, Schöffski O, Vauth C, Volmer T, Wahler S, Wasem J, Weber C. 2007. Deutsche Empfehlungen zur gesundheitsökonomischen Evaluation - dritte und aktualisierte Fassung des Hannoveraner Konsens. *Gesundh ökon Qual manag* 12(05):285-290.
- Greiner W, Claes C, Busschbach JJ, von der Schulenburg JM. 2005. Validating the EQ-5D with time trade off for the German population. *Eur J Health Econ* 6(2):124-30.
- Grigoriadis N, van Pesch V. 2015. A basic overview of multiple sclerosis immunopathology. *Eur J Neurol* 22(Suppl_2):S3-S13.
- Grytten Torkildsen N, Lie SA, Aarseth JH, Nyland H, Myhr KM. 2008. Survival and cause of death in multiple sclerosis: results from a 50-year follow-up in Western Norway. *Mult Scler* 14(9):1191-1198.
- Gyllenstein H, Kavaliunas A, Murley C, Alexanderson K, Hillert J, Tinghög P, Friberg E. 2019. Costs of illness progression for different multiple sclerosis phenotypes: a population-based study in Sweden. *Mult Scler J Exp Transl Clin* 5(2).
- Haas J, Jeffery D, Silva D, Meier DP, Meinert R, Cohen J, Hartung H-P. 2019. Early switch to fingolimod reduces rates of severe relapses over the long term: Post hoc analysis from the FREEDOMS, FREEDOMS II, and TRANSFORMS studies. *Mult Scler Relat Disord* 36:101335.
- Haase R, Kern R, K. A, Wunderlich M, Dominik M, Ziemssen T. 2018a. Development of the adaptive, bilingual multiple sclerosis Health Resource Survey (MS-HRS). *Mult Scler* 24(Suppl_2):S738-S980.
- Haase R, Wunderlich M, Dillenseger A, Kern R, Akgün K, Ziemssen T. 2018b. Improving multiple sclerosis management and collecting safety information in the real world: the MSDS3D software approach. *Expert Opin Drug Saf* 17(4):369-378.
- Hawton AJ, Green C. 2016. Multiple sclerosis: relapses, resource use, and costs. *Eur J Health Econ* 17(7):875-884.
- He A, Spelman T, Jokubaitis V, Havrdova E, Horakova D, Trojano M, Lugaresi A, Izquierdo G, Grammond P, Duquette P, Girard M, Pucci E, Iuliano G, Alroughani R, Oreja-Guevara C, Fernandez-Bolaños R, Grand'Maison F, Sola P, Spitaleri D, Granella F, Terzi M, Lechner-Scott J, Van Pesch V, Hupperts R, Sánchez-Menoyo JL, Hodgkinson S, Rozsa C, Verheul F, Butzkueven H, Kalincik T, Group ftMS. 2015. Comparison of Switch to Fingolimod or Interferon

- Beta/Glatiramer Acetate in Active Multiple Sclerosis. *JAMA Neurol* 72(4):405-413.
- Healy BC, Degano IR, Schreck A, Rintell D, Weiner H, Chitnis T, Glanz BI. 2012. The impact of a recent relapse on patient-reported outcomes in subjects with multiple sclerosis. *Qual Life Res* 21(10):1677-1684.
- Heesen C, Reich C, Borchers C, Gold SM, Schulz KH. 2007. Patientenbasierte Scoringssysteme bei Multipler Sklerose - Untersuchung zur Validierung der United Kingdom Disability Scale (UNDS) und des Patient Generated Index (PGI). *Neurology and Rehabilitation* 13(1):24-36.
- Heslin M, Babalola O, Ibrahim F, Stringer D, Scott D, Patel A. 2018. A Comparison of Different Approaches for Costing Medication Use in an Economic Evaluation. *Value Health* 21(2):185-192.
- Huhn K, Senger D, Utz KS, Schmidt M, Fröhlich K, Waschbisch A, Seifert F, Dörfler A, Lee D-H, Linker RA. 2019. No evidence of disease activity status over 3 years in a real-world cohort of relapsing remitting MS patients in Germany. *Mult Scler Relat Disord* 27:133-138.
- Hyland M, Rudick RA. 2011. Challenges to clinical trials in multiple sclerosis: outcome measures in the era of disease-modifying drugs. *Curr Opin Neurol* 24(3):255-61.
- IAB. Institut für Arbeitsmarkt- und Berufsforschung. 2011. Daten zur kurzfristigen Entwicklung von Wirtschaft und Arbeitsmarkt Pressemitteilung vom 15.09.2016 – 324/16 [accessed: 15.01.2020] URL: <https://www.iab.de/de/publikationen/jahresbericht/jb2011.aspx>.
- IQWiG. Institut für Qualität und Wirtschaftlichkeit im Gesundheitswesen. 2008. Technischer Anhang Kostenbestimmung [accessed: 15.01.2020] URL: <https://www.iqwig.de/de/methoden/methodenpapier/methoden-knb-fruehere-versionen.3023.html>.
- IQWiG. Institut für Qualität und Wirtschaftlichkeit im Gesundheitswesen. 2017. Allgemeine Methoden Version 5.0 vom 10.07.2017 [General Methods Version 5.0 from 10.07.2017] [accessed: 15.01.2020] URL: <https://www.iqwig.de/de/methoden/methodenpapier.3020.html>.
- Jimenez-Martin S, Prieto CV. 2012. The trade-off between formal and informal care in Spain. *Eur J Health Econ* 13(4):461-490.
- Jo C. 2014. Cost-of-illness studies: concepts, scopes, and methods. *Clin Mol Hepatol* 20(4):327-337.
- Johnston K, Buxton MJ, Jones DR, Fitzpatrick R. 1999. Assessing the costs of healthcare technologies in clinical trials. *Health Technol Assess* 3(6):1-76.
- Kalincik T. 2015. Multiple Sclerosis Relapses: Epidemiology, Outcomes and Management. A Systematic Review. *Neuroepidemiology* 44(4):199-214.
- Kalincik T, Cutter G, Spelman T, Jokubaitis V, Havrdova E, Horakova D, Trojano M, Izquierdo G, Girard M, Duquette P. 2015. Defining reliable disability outcomes in multiple sclerosis. *Brain* 138(11):3287-3298.

- Kalincik T, Vivek V, Jokubaitis V, Lechner-Scott J, Trojano M, Izquierdo G, Lugaresi A, Grand'maison F, Hupperts R, Oreja-Guevara C, Bergamaschi R, Iuliano G, Alroughani R, Van Pesch V, Amato MP, Slee M, Verheul F, Fernandez-Bolanos R, Fiol M, Spitaleri DL, Cristiano E, Gray O, Cabrera-Gomez JA, Shaygannejad V, Herbert J, Vucic S, Needham M, Petkovska-Boskova T, Sirbu CA, Duquette P, Girard M, Grammond P, Boz C, Giuliani G, Rio ME, Barnett M, Flechter S, Moore F, Singhal B, Bacile EA, Saladino ML, Shaw C, Skromne E, Poehlau D, Vella N, Spelman T, Liew D, Kilpatrick TJ, Butzkueven H. 2013. Sex as a determinant of relapse incidence and progressive course of multiple sclerosis. *Brain* 136(12):3609-3617.
- Kappos L, Butzkueven H, Wiendl H, Spelman T, Pellegrini F, Chen Y, Dong Q, Koendgen H, Belachew S, Trojano M. 2018a. Greater sensitivity to multiple sclerosis disability worsening and progression events using a roving versus a fixed reference value in a prospective cohort study. *Mult Scler* 24(7):963-973.
- Kappos L, European Study Group on Interferon β -1b in Secondary Progressive MS. 1998. Placebo-controlled multicentre randomised trial of interferon beta-1b in treatment of secondary progressive multiple sclerosis. European Study Group on interferon beta-1b in secondary progressive MS. *Lancet* 352(9139):1491-7.
- Kappos L, Radue E-W, O'Connor P, Polman C, Hohlfeld R, Calabresi P, Selmaj K, Agoropoulou C, Leyk M, Zhang-Auberson L, Burtin P. 2010. A Placebo-Controlled Trial of Oral Fingolimod in Relapsing Multiple Sclerosis. *N Engl J Med* 362(5):387-3401.
- Kappos L, Weinshenker B, Pozzilli C, Thompson AJ, Dahlke F, Beckmann K, Polman C, McFarland H. 2004. Interferon beta-1b in secondary progressive MS: a combined analysis of the two trials. *Neurology* 63(10):1779-1787.
- Kappos L, Wolinsky J, Giovannoni G, Arnold D, Wang Q, Bernasconi C, Model F, Manfrini M, Belachew S, Hauser S. 2018b. Relapse-associated worsening and progression independent of relapse activity in patients with relapsing multiple sclerosis in the Phase III OPERA I and OPERA II studies [P547]. *Mult Scler* 24(Suppl_2):S121-S327.
- Karampampa K, Gustavsson A, Miltenburger C, Eckert B. 2012a. Treatment experience, burden and unmet needs (TRIBUNE) in MS study: results from five European countries. *Mult Scler* 18(Suppl_2):S7-S15.
- Karampampa K, Gustavsson A, Miltenburger C, Neidhardt K, Lang M. 2012b. Treatment experience, burden and unmet needs (TRIBUNE) in MS study: results from Germany. *Mult Scler* 18(Suppl_2):S23-S27.
- KBV. Kassenärztliche Bundesvereinigung. 2011. Einheitlicher Bewertungsmaßstab [accessed: 15.01.2020] URL: https://www.kvsa.de/fileadmin/user_upload/PDF/Praxis/Honorarverteilung/EBM2011.pdf.
- Khan O, Rieckmann P, Boyko A, Selmaj K, Zivadinov R, Group GS. 2013. Three times weekly glatiramer acetate in relapsing–remitting multiple sclerosis. *Ann Neurol* 73(6):705-713.
- Kieseier BC. 2011. The mechanism of action of interferon-beta in relapsing multiple sclerosis. *CNS Drugs* 25(6):491-502.

- Killestein J, Polman CH. 2011. Determinants of interferon β efficacy in patients with multiple sclerosis. *Nat Rev Neurol* 7(4):221-228.
- Kind P. 1990. Measuring valuations for health states: a survey of patients in general practice. University of York, Discussion Paper.
- Kingwell E, van der Kop M, Zhao Y, Shirani A, Zhu F, Oger J, Tremlett H. 2012. Relative mortality and survival in multiple sclerosis: findings from British Columbia, Canada. *J Neurol Neurosurg Psychiatry* 83(1):61-66.
- Kobelt G, Berg J, Lindgren P, Fredrikson S, Jonsson B. 2006a. Costs and quality of life of patients with multiple sclerosis in Europe. *J Neurol Neurosurg Psychiatry* 77(8):918-926.
- Kobelt G, Berg J, Lindgren P, Jonsson B. 2006b. Costs and quality of life in multiple sclerosis in Europe: method of assessment and analysis. *Eur J Health Econ* 7(Suppl_2):S5-S13.
- Kobelt G, Eriksson J, Phillips G, Berg J. 2017a. The burden of multiple sclerosis 2015: Methods of data collection, assessment and analysis of costs, quality of life and symptoms. *Mult Scler* 23(Suppl_2):S4-S16.
- Kobelt G, Jönsson L, Miluse P, Havrdov EK. 2019a. Real-Life Outcome in Multiple Sclerosis in the Czech Republic. *Multiple Sclerosis International* 2019:8.
- Kobelt G, Teich V, Cavalcanti M, Canzonieri AM. 2019b. Burden and cost of multiple sclerosis in Brazil. *PLoS One* 14(1):e0208837.
- Kobelt G, Thompson A, Berg J, Gannedahl M, Eriksson J, Group MS, European Multiple Sclerosis P. 2017b. New insights into the burden and costs of multiple sclerosis in Europe. *Mult Scler* 23(8):1123-1136.
- Koch-Henriksen N, Thygesen LC, Sørensen PS, Magyari M. 2019. Worsening of disability caused by relapses in multiple sclerosis: a different approach. *Mult Scler Relat Disord* 32:1-8.
- Koo TK, Li MY. 2016. A Guideline of Selecting and Reporting Intraclass Correlation Coefficients for Reliability Research. *J Chiropr Med* 15(2):155-163.
- Kurtzke JF. 1955. A new scale for evaluating disability in multiple sclerosis. *Neurology* 5(8):580-580.
- Kurtzke JF. 1983. Rating neurologic impairment in multiple sclerosis: an expanded disability status scale (EDSS). *Neurology* 33(11):1444-1152.
- La Mantia L, Di Pietrantonj C, Rovaris M, Rigon G, Frau S, Berardo F, Gandini A, Longobardi A, Weinstock-Guttman B, Vaona A. 2016. Interferons-beta versus glatiramer acetate for relapsing-remitting multiple sclerosis. *Cochrane Database Syst Rev* 11:Cd009333.
- Lambert J, Hansen BB, Arnould B, Grataloup G, Guillemin I, Højbjerg L, Strandberg-Larsen M, Reilly MC. 2014. Linguistic validation into 20 languages and content validity of the rheumatoid arthritis-specific Work Productivity and Activity Impairment Questionnaire. *PATIENT* 7(2):171-176.

- Landfeldt E, Zethraeus N, Lindgren P. 2019. Standardized Questionnaire for the Measurement, Valuation, and Estimation of Costs of Informal Care Based on the Opportunity Cost and Proxy Good Method. *Appl Health Econ Health Policy* 17(1):15-24.
- Larg A, Moss JR. 2011. Cost-of-illness studies. *Pharmacoeconomics* 29(8):653-671.
- Lassmann H, van Horssen J, Mahad D. 2012. Progressive multiple sclerosis: pathology and pathogenesis. *Nat Rev Neurol* 8(11):647-656.
- Leggett LE, Khadaroo RG, Holroyd-Leduc J, Lorenzetti DL, Hanson H, Wagg A, Padwal R, Clement F. 2016. Measuring Resource Utilization: A Systematic Review of Validated Self-Reported Questionnaires. *Medicine (Baltimore)* 95(10):e2759.
- Li H, Hu F, Zhang Y, Li K. 2019. Comparative efficacy and acceptability of disease-modifying therapies in patients with relapsing–remitting multiple sclerosis: a systematic review and network meta-analysis. *J Neurol*.
- Ljungqvist O, Thanh NX, Nelson G. 2017. ERAS—Value based surgery. *J Surg Oncol* 116(5):608-612.
- Lorscheider J, Buzzard K, Jokubaitis V, Spelman T, Havrdova E, Horakova D, Trojano M, Izquierdo G, Girard M, Duquette P, Prat A, Lugaresi A, Grand'Maison F, Grammond P, Hupperts R, Alroughani R, Sola P, Boz C, Pucci E, Lechner-Scott J, Bergamaschi R, Oreja-Guevara C, Iuliano G, Van Pesch V, Granella F, Ramo-Tello C, Spitaleri D, Petersen T, Slee M, Verheul F, Ampapa R, Amato MP, McCombe P, Vucic S, Sanchez Menoyo JL, Cristiano E, Barnett MH, Hodgkinson S, Olascoaga J, Saladino ML, Gray O, Shaw C, Moore F, Butzkueven H, Kalincik T. 2016. Defining secondary progressive multiple sclerosis. *Brain* 139(9):2395-2405.
- Lublin FD, Reingold SC, Cohen JA, Cutter GR, Sørensen PS, Thompson AJ, Wolinsky JS, Balcer LJ, Banwell B, Barkhof F, Bebo B, Calabresi PA, Clanet M, Comi G, Fox RJ, Freedman MS, Goodman AD, Inglese M, Kappos L, Kieseier BC, Lincoln JA, Lubetzki C, Miller AE, Montalban X, O'Connor PW, Petkau J, Pozzilli C, Rudick RA, Sormani MP, Stüve O, Waubant E, Polman CH. 2014. Defining the clinical course of multiple sclerosis: The 2013 revisions. *Neurology* 83(3):278-286.
- Luce BR, Manning WG, Siegel JE, Lipscomb J. 1996. Estimating costs in cost-effectiveness analysis. In: Gold MR, Siegel JE, Russell LB, Weinstein MC (Eds) *Cost-effectiveness in health and medicine*. Oxford University Press, New York. p. 176-213.
- Martin A, Jones A, Mugford M, Shemilt I, Hancock R, Wittenberg R. 2012. Methods used to identify and measure resource use in economic evaluations: a systematic review of questionnaires for older people. *Health Econ* 21(8):1017-1022.
- McCrone P, Heslin M, Knapp M, Bull P, Thompson A. 2008. Multiple sclerosis in the UK: service use, costs, quality of life and disability. *Pharmacoeconomics* 26(10):847-860.

- McDonald WI, Compston A, Edan G, Goodkin D, Hartung H-P, Lublin FD, McFarland HF, Paty DW, Polman CH, Reingold SC, Sandberg-Wollheim M, Sibley W, Thompson A, Van Den Noort S, Weinshenker BY, Wolinsky JS. 2001. Recommended diagnostic criteria for multiple sclerosis: Guidelines from the international panel on the diagnosis of multiple sclerosis. *Ann Neurol* 50(1):121-127.
- McKenna SP, Doward LC, Twiss J, Hagell P, Oprandi NC, Fisk J, Grand'Maison F, Bhan V, Arbizu T, Brassat D, Kohlmann T, Meads DM, Eckert BJ. 2010. International Development of the Patient-Reported Outcome Indices for Multiple Sclerosis (PRIMUS). *Value Health* 13(8):946-951.
- Meuth SG, Simon OJ, Grimm A, Melzer N, Herrmann AM, Spitzer P, Landgraf P, Wiendl H. 2008. CNS inflammation and neuronal degeneration is aggravated by impaired CD200-CD200R-mediated macrophage silencing. *J Neuroimmunol* 194(1-2):62-69.
- Mihaylova B, Briggs A, O'Hagan A, Thompson SG. 2011. Review of Statistical Methods for Analysing Healthcare Resources and Costs. *Health Econ* 20(8):897-916.
- Miller DH, Leary SM. 2007. Primary-progressive multiple sclerosis. *Lancet Neurol* 6(10):903-912.
- Montalban X, Gold R, Thompson AJ, Otero-Romero S, Amato MP, Chandraratna D, Clanet M, Comi G, Derfuss T, Fazekas F, Hartung HP, Havrdova E, Hemmer B, Kappos L, Liblau R, Lubetzki C, Marcus E, Miller DH, Olsson T, Pilling S, Selmaj K, Siva A, Sorensen PS, Sormani MP, Thalheim C, Wiendl H, Zipp F. 2018.ECTRIMS/EAN Guideline on the pharmacological treatment of people with multiple sclerosis. *Mult Scler* 24(2):96-120.
- MSIF. Multiple Sclerosis International Federation. 2013. Atlas of MS 2013: Mapping Multiple Sclerosis around the World [accessed: 15.01.2020] URL: <http://www.msif.org/wp-content/uploads/2014/09/Atlas-of-MS.pdf>.
- Naci H, Fleurence R, Birt J, Duhig A. 2010. Economic burden of multiple sclerosis: a systematic review of the literature. *Pharmacoeconomics* 28(5):363-379.
- Ness N-H, Haase R, Kern R, Schriefer D, Ettl B, Cornelissen C, Akgün K, Ziemssen T. 2020a. Multiple Sclerosis Health Resource Utilization Survey (MS-HRS): Development and Validation Study. *J Med Internet Res* 22(3):e17921.
- Ness N-H, Schriefer D, Haase R, Ettl B, Cornelissen C, Ziemssen T. 2019a. Differentiating societal costs of disability worsening in multiple sclerosis. *J Neurol* 267(4):1035-1042.
- Ness N-H, Schriefer D, Haase R, Ettl B, Wunderlich M, Akgün K, Ziemssen T. 2019b. Real-World Evidence on the influence of relapses and relapse associated progression on societal economic costs in patients with multiple sclerosis ECTRIMS 2019 – Oral Presentations. *Mult Scler* 25(Suppl_2):3-130.
- Ness N-H, Schriefer D, Haase R, Ettl B, Ziemssen T. 2020b. Real-World Evidence on the Societal Economic Relapse Costs in Patients with Multiple Sclerosis. *Pharmacoeconomics* (online first).

- Novartis. 2011. Dossier zur Nutzenbewertung gemäß § 35a SGB V: Modul 3 A
[accessed: 15.01.2020] URL: https://www.g-ba.de/downloads/92-975-41/2011-09-21_Modul3_Fingolimod.pdf.
- O'Brien JA, Ward AJ, Patrick AR, Caro J. 2003. Cost of managing an episode of relapse in multiple sclerosis in the United States. *BMC Health Serv Res* 3(17).
- O'Connell K, Kelly SB, Fogarty E, Duggan M, Buckley L, Hutchinson M, McGuigan C, Tubridy N. 2014. Economic costs associated with an MS relapse. *Mult Scler Relat Disord* 3(6):678-683.
- O'Gorman C, Lucas R, Taylor B. 2012. Environmental risk factors for multiple sclerosis: a review with a focus on molecular mechanisms. *Int J Mol Sci* 13(9):11718-11752.
- O'Sullivan AK, Thompson D, Drummond MF. 2005. Collection of health-economic data alongside clinical trials: is there a future for piggyback evaluations? *Value Health* 8(1):67-79.
- Oliva-Moreno J, Peña-Longobardo LM, García-Mochón L, del Río Lozano M, Mosquera Metcalfe I, García-Calvente MdM. 2019. The economic value of time of informal care and its determinants (The CUIDARSE Study). *PLoS One* 14(5):e0217016.
- Oliva-Moreno J, Trapero-Bertran M, Peña-Longobardo LM, del Pozo-Rubio R. 2017. The valuation of informal care in cost-of-illness studies: a systematic review. *Pharmacoeconomics* 35(3):331-345.
- Patzold U, Pocklington PR. 1982. Course of multiple sclerosis. First results of a prospective study carried out of 102 MS patients from 1976-1980. *Acta Neurol Scand* 65(4):248-266.
- Petersen G, Wittmann R, Arndt V, Gopffarth D. 2014. Epidemiology of multiple sclerosis in Germany: regional differences and drug prescription in the claims data of the statutory health insurance. *Nervenarzt* 85(8):990-998.
- Pike J, Grosse SD. 2018. Friction Cost Estimates of Productivity Costs in Cost-of-Illness Studies in Comparison with Human Capital Estimates: A Review. *Appl Health Econ Health Pol* 16(6):765-778.
- Pugliatti M, Harbo HF, Holmoy T, Kampman MT, Myhr KM, Riise T, Wolfson C. 2008. Environmental risk factors in multiple sclerosis. *Acta Neurol Scand Suppl* 188:34-40.
- Raftery J. 2000. Costing in economic evaluation. *BMJ* 320(7249):1597.
- Ragonese P, Aridon P, Salemi G, D'Amelio M, Savettieri G. 2008. Mortality in multiple sclerosis: a review. *Eur J Neurol* 15(2):123-127.
- Raimundo K, Tian H, Zhou H, Zhang X, Kahler KH, Agashivala N, Kim E. 2013. Resource utilization, costs and treatment patterns of switching and discontinuing treatment of MS patients with high relapse activity. *BMC Health Serv Res* 13(131).
- Raulinajtys-Grzybek M, Świdarska GK. 2015. Payment by results vs. costs of 24-hour standby in hospitals: Evidence from Poland. *Argum Oecon* 2(35):85-105.

- Reese JP, John A, Wienemann G, Wellek A, Sommer N, Tackenberg B, Balzer-Geldsetzer M, Dodel R. 2011. Economic burden in a German cohort of patients with multiple sclerosis. *Eur Neurol* 66(6):311-321.
- Reilly M, Bracco A, Ricci JF, Santoro J, Stevens T. 2004. The validity and accuracy of the Work Productivity and Activity Impairment questionnaire–irritable bowel syndrome version (WPAI: IBS). *Aliment Pharmacol Ther* 20(4):459-467.
- Reilly MC, Gooch KL, Wong RL, Kupper H, Van der Heijde D. 2010. Validity, reliability and responsiveness of the Work Productivity and Activity Impairment Questionnaire in ankylosing spondylitis. *Rheumatology* 49(4):812-819.
- Reilly MC, Zbrozek AS, Dukes EM. 1993. The validity and reproducibility of a work productivity and activity impairment instrument. *Pharmacoeconomics* 4(5):353-365.
- Rice DP. 1967. Estimating the cost of illness. *Am J Public Health Nations Health* 57(3):424-440.
- Ridyard CH, Hughes DA. 2010. Methods for the collection of resource use data within clinical trials: a systematic review of studies funded by the UK Health Technology Assessment program. *Value Health* 13(8):867-872.
- Ridyard CH, Hughes DA, Team D. 2015. Taxonomy for Methods of Resource Use Measurement. *Health Econ* 24(3):372-378.
- Roick C, Kilian R, Matschinger H, Bernert S, Mory C, Angermeyer MC. 2001. Die deutsche Version des Client Sociodemographic and Service Receipt Inventory. *Psychiatr Prax* 28(Sup. 2):84-90.
- Rommer PS, Milo R, Han MH, Satyanarayan S, Sellner J, Hauer L, Illes Z, Warnke C, Laurent S, Weber MS, Zhang Y, Stuve O. 2019. Immunological Aspects of Approved MS Therapeutics. *Front Immunol* 10(1564).
- Rossier P, Wade DT. 2002. The Guy's Neurological Disability Scale in patients with multiple sclerosis: a clinical evaluation of its reliability and validity. *Clin Rehabil* 16(1):75-95.
- Ruger JP, Reiff M. 2016. A checklist for the conduct, reporting, and appraisal of microcosting studies in health care: protocol development. *JMIR Res Protoc* 5(4):e195.
- Russell LB, Siegel JE, Daniels N, Gold MR, Luce BR, Mandelblatt JS. 1996. Cost-effectiveness analysis as a guide to resource allocation in health: roles and limitations. In: Gold MR, Siegel JE, Russell LB, Weinstein MC (Eds) *Cost-effectiveness in health and medicine*. Oxford University Press, New York. p. 3-24.
- Sanchirico M, Caldwell-Tarr A, Mudumby P, Hashemi L, Dufour R. 2019. Treatment Patterns, Healthcare Resource Utilization, and Costs Among Medicare Patients with Multiple Sclerosis in Relation to Disease-Modifying Therapy and Corticosteroid Treatment. *Neurol Ther* 8(1):121-133.
- Scalfari A, Neuhaus A, Degenhardt A, Rice GP, Muraro PA, Daumer M, Ebers GC. 2010. The natural history of multiple sclerosis: a geographically based study 10: relapses and long-term disability. *Brain* 133(7):1914-1929.

- Schweikert B, Hahmann H, Leidl R. 2008. Development and first assessment of a questionnaire for health care utilization and costs for cardiac patients. *BMC Health Serv Res* 8(187).
- Segel JE. 2006. Cost-of-illness studies—a primer. RTI International Center of Excellence in Health Promotion Economics.
- Sharrack B, Hughes RA. 1999. The Guy's Neurological Disability Scale (GNDS): a new disability measure for multiple sclerosis. *Mult Scler* 5(4):223-233.
- Simoens S. 2009. Health Economic Assessment: A Methodological Primer. *Int J Environ Res Public Health* 6(12):2950-2966.
- Simpson S, Jr., Blizzard L, Otahal P, Van der Mei I, Taylor B. 2011. Latitude is significantly associated with the prevalence of multiple sclerosis: a meta-analysis. *J Neurol Neurosurg Psychiatry* 82(10):1132-1141.
- Smith KA, Rudmik L. 2013. Cost collection and analysis for health economic evaluation. *Otolaryngol Head Neck Surg* 149(2):192-199.
- Stuke K, Flachenecker P, Zettl UK, Elias W, Freidel M, Haas J, Pitschnau-Michel D, Schimrigk S, Rieckmann P. 2008. MS-Register in Deutschland 2008 – Symptomatik der MS [accessed: 15.01.2020] URL: https://www.dmsg.de/dokumentearchiv/dgn2008_msregister.pdf.
- Stull DE, Leidy NK, Parasuraman B, Chassany O. 2009. Optimal recall periods for patient-reported outcomes: challenges and potential solutions. *Curr Med Res Opin* 25(4):929-942.
- Tan SS. 2009. Microcosting in economic evaluations: Issues of accuracy, feasibility, consistency and generalisability. Erasmus University Rotterdam, Dissertation.
- Tan SS, Rutten FFH, van Ineveld BM, Redekop WK, Hakkaart-van Roijen L. 2009. Comparing methodologies for the cost estimation of hospital services. *Eur J Health Econ* 10(1):39-45.
- Tarricone R. 2006. Cost-of-illness analysis: What room in health economics? *Health Policy* 77(1):51-63.
- The North American Study Group on Interferon beta-1b in Secondary Progressive MS. 2004. Interferon beta-1b in secondary progressive MS: Results from a 3-year controlled study. *Neurology* 63(10):1788-1795.
- Thompson S, Wordsworth S. 2001. An annotated cost questionnaire for completion by patients. University of York, Discussion Paper.
- Thorn JC, Brookes ST, Ridyard C, Riley R, Hughes DA, Wordsworth S, Noble SM, Thornton G, Hollingworth W. 2018. Core Items for a Standardized Resource Use Measure: Expert Delphi Consensus Survey. *Value Health* 21(6):640-649.
- Thorn JC, Coast J, Cohen D, Hollingworth W, Knapp M, Noble SM, Ridyard C, Wordsworth S, Hughes D. 2013. Resource-Use Measurement Based on Patient Recall: Issues and Challenges for Economic Evaluation. *Appl Health Econ Health Pol* 11(3):155-161.

- Tremlett H, Paty D, Devonshire V. 2006. Disability progression in multiple sclerosis is slower than previously reported. *Neurology* 66(2):172-177.
- Turner B, Cree BAC, Kappos L, Montalban X, Papeix C, Wolinsky JS, Buffels R, Fiore D, Garren H, Han J, Hauser SL. 2019. Ocrelizumab efficacy in subgroups of patients with relapsing multiple sclerosis. *J Neurol* 266(5):1182-1193.
- Twork S, Wiesmeth S, Spindler M, Wirtz M, Schipper S, Pohlau D, Klewer J, Kugler J. 2010. Disability status and quality of life in multiple sclerosis: non-linearity of the Expanded Disability Status Scale (EDSS). *Health Qual Life Outcomes* 8(55).
- Uitdehaag BMJ. 2018. Disability Outcome Measures in Phase III Clinical Trials in Multiple Sclerosis. *CNS Drugs* 32(6):543-558.
- van den Hout WB. 2010. The value of productivity: human-capital versus friction-cost method. *Ann Rheum Dis* 69(Suppl_1):S89-S91.
- van Lier LI, Bosmans JE, van Hout HPJ, Mokkink LB, van den Hout WB, de Wit GA, Dirksen CD, Nies H, Hertogh C, van der Roest HG. 2018. Consensus-based cross-European recommendations for the identification, measurement and valuation of costs in health economic evaluations: a European Delphi study. *Eur J Health Econ* 19(7):993-1008.
- van Munster CEP, Uitdehaag BMJ. 2017. Outcome Measures in Clinical Trials for Multiple Sclerosis. *CNS Drugs* 31(3):217-236.
- Vercellino M, Romagnolo A, Mattioda A, Masera S, Piacentino C, Merola A, Chiò A, Mutani R, Cavalla P. 2009. Multiple sclerosis relapses: a multivariable analysis of residual disability determinants. *Acta Neurol Scand* 119(2):126-130.
- VFA. Verband Forschender Arzneimittelhersteller. 2010. Register nicht-interventioneller Studien - Prospective phArmaceutic cohoRt evaluation [accessed: 15.01.2020] URL: <https://www.vfa.de/de/arzneimittel-forschung/datenbanken-zu-arzneimitteln/nisdb/nis-details/500>.
- VFA. Verband Forschender Arzneimittelhersteller. 2011. PANGAEA Post-authorization non-interventional German safety study of Gilenya® in MS patients [accessed: 15.01.2020] URL: <https://www.vfa.de/de/arzneimittel-forschung/datenbanken-zu-arzneimitteln/nisdb/nis-details/596>.
- Vickrey BG, Hays RD, Harooni R, Myers LW, Ellison GW. 1995. A health-related quality of life measure for multiple sclerosis. *Qual Life Res* 4(3):187-206.
- Vormfelde SV, Ortler S, Ziemssen T. 2016. Multiple Sclerosis Therapy With Disease-Modifying Treatments in Germany: The PEARL (Prospective phArmaceutic cohoRt evaluation) Noninterventional Study Protocol. *JMIR Res Protoc* 5(1):e23.
- Wallin MT, Culpepper WJ, Nichols E, Bhutta ZA, Gebrehiwot TT, Hay SI, Khalil IA, Krohn KJ, Liang X, Naghavi M, Mokdad AH, Nixon MR, Reiner RC, Sartorius B, Smith M, Topor-Madry R, Werdecker A, Vos T, Feigin VL, Murray CJL. 2019. Global, regional, and national burden of multiple sclerosis 1990-2013;2016: a systematic analysis for the Global Burden of Disease Study 2016. *Lancet Neurol* 18(3):269-285.

- WHO. World Health Organization. 1995. Physical status: the use and interpretation of anthropometry physical status. World Health Organization, Geneva.
- WHO. World Health Organization. 2009. Guide to identifying the economic consequences of disease and injury. World Health Organization, Geneva.
- Wickstrom A, Dahle C, Vrethem M, Svenningsson A. 2014. Reduced sick leave in multiple sclerosis after one year of natalizumab treatment. A prospective ad hoc analysis of the TYNERGY trial. *Mult Scler* 20(8):1095-1101.
- Wiendl H, Meuth SG. 2015. Pharmacological Approaches to Delaying Disability Progression in Patients with Multiple Sclerosis. *Drugs* 75(9):947-977.
- Wundes A, Brown T, Bienen EJ, Coleman CI. 2010. Contribution of intangible costs to the economic burden of multiple sclerosis. *J Med Econ* 13(4):626-632.
- Zeller J. 2012. Immuntherapeutika und Zytostatika. In: Schwabe U, Paffrath D, Ludwig W-D, Klauber J (Eds) *Arzneiverordnungs-Report 2012: Aktuelle Daten, Kosten, Trends und Kommentare*. Springer, Heidelberg. p. 679-696.
- Ziemssen T, De Stefano N, Sormani MP, Van Wijmeersch B, Wiendl H, Kieseier BC. 2015a. Optimizing therapy early in multiple sclerosis: An evidence-based view. *Mult Scler Relat Disord* 4(5):460-469.
- Ziemssen T, Kern R, Cornelissen C. 2015b. The PANGAEA study design - a prospective, multicenter, non-interventional, long-term study on fingolimod for the treatment of multiple sclerosis in daily practice. *BMC Neurol* 15(93).

Supplement

Multiple Sclerosis Health Resource Utilization Survey (MS-HRS)

The purpose of this questionnaire is to determine the total cost of a multiple sclerosis therapy. Due to the lack of complete data, it is often difficult to estimate the costs. Your contribution to our efforts to better understand these costs is therefore of particular importance.

When answering the questions, please note that the period for cost collection refers to the last 6 months. Please answer the questions one by one, any shortcuts between the questions will be announced.

Thank you very much.

I. Core questions (1 to 12)

1. What is your current job?

- ☐ Employed
- ☐ Self-employed
- ☐ Looking for work
- ☐ Housewife/man
- ☐ In study/training
- ☐ Retired due to MS to a degree of ____%
- ☐ Retired for other reasons
- ☐ Other

2. How many hours per week do you currently work? (If unemployed, fill in „0“)

_____ hours

3. How many days have you been on sick leave due to MS in the last 6 months?

(If unemployed, fill in „0“)

_____ days

4. Due to my MS, my state of health has reduced my productivity at work (e.g. concentration or speed at work) in the last half year. (Choose a high number if MS affected your work a lot)

☐ 0 ☐ 1 ☐ 2 ☐ 3 ☐ 4 ☐ 5 ☐ 6 ☐ 7 ☐ 8 ☐ 9 ☐ 10

No effect
on my work

Completely
prevented me
from working

5. Due to my MS, I have been hospitalized in a hospital, rehabilitation clinic or nursing home (including overnight stay) in the last 6 months.

☐ No → Proceed to question 6.

☐ Yes, in detail:

Of which _____ were days in a hospital, and thereof _____ days in neurology.

Of which _____ were days in a rehabilitation clinic.

Of which _____ were days in a nursing home.

6. Due to my MS, I have been treated in a hospital, a rehabilitation clinic or a nursing home, i.e. in a day clinic (without an overnight stay) or a night clinic (without a day stay) in the last 6 months.

☐ No → Proceed to question 7.

☐ Yes, in detail:

Of which _____ were days in a hospital, and thereof _____ days in neurology.

Of which _____ were days in a rehabilitation clinic.

Of which _____ were days in a nursing home.

7. Due to my MS, I have had contact with the following specialists or professional groups on an outpatient basis in a practice or hospital in the last 6 months. Please only list those contacts that have not already been reported in question 5 or 6. (Multiple entries of different specialists are possible)

() I did not consult any of the professionals listed below. → Proceed to question 8.

() Yes, in detail:

General practitioner / family doctor _____ times
Neurologist _____ times
Other specialist (e.g. urologist, ophthalmologist, psychologist) _____ times
MS nurse _____ times
Physiotherapist _____ times
Occupational therapist _____ times
Optician _____ times
Speech therapist _____ times
Acupuncture _____ times
Other alternative healthcare _____ times
Other _____ times

8. Due to my MS, I have been in one of the following examinations for the last 6 months: Magnetic Resonance Imaging (MRI), Computer Tomography (CT), Lumbar Puncture or Blood Examinations.

() No → Proceed to question 9.

() Yes, in detail:

_____ Magnetic Resonance Imaging (MRI) (count)
_____ Computer Tomography (CT) (count)
_____ Lumbar Puncture (count)
_____ Blood Examinations (count)
_____ Others (count)

9. Due to my MS, I have needed help in the household or with everyday tasks within the last 6 months.

() No → Proceed to question 10.

() Yes, by the following persons:

Professional care specialist (nursing service) _____ hours per week on average

Domestic help or personal advisor _____ hours per week on average

Friends or family members _____ hours per week on average

10. How much (in Euro) have you spent on over-the-counter medicines to treat MS symptoms over the last 6 months?

_____ Euro

11. How much (in Euro) have you spent on medical consumables (e.g. bandages, patches) approximately due to your MS during the last 6 months?

_____ Euro

12. Have you made any changes to your house, apartment or car in the past 6 months, or have you needed special equipment or aids due to your MS?

() No (zero costs) → Proceed to question 13.

() Yes, in detail:

Changes to your home due to problems related to your MS: _____ Euro

Changes to your car due to problems related to your MS: _____ Euro

For walking aids, wheelchair (manual or electric), electric scooter: _____ Euro

For a special hospital bed: _____ Euro

Other investments: _____ Euro

II. Additional questions (13 to 24)

13. By how many hours per week did you have to permanently reduce your workload due to MS? (If not applicable, then „0“)
____ hours
14. Due to my MS, my earnings in the last 6 months have reduced by ... €. (If not applicable, then „0“)
____ Euro
15. Due to my MS, I had to change the type of work or my job in the last 6 months.
☐ No
☐ Yes
16. Have your relatives reduced their working hours during the last 6 months to support you due to your MS?
☐ No
☐ Yes, from ____ hours per week to ____ hours per week
17. Are you responsible for caring for other family members (e.g. relatives or children in need of care - including children in education)?
☐ No
☐ Yes
18. What everyday tasks do you need help with? (Multiple answers possible).
☐ Movement
☐ Body care
☐ Transport
☐ Other
19. What level of education have you acquired?
☐ Secondary school
☐ University degree
☐ Completed vocational training
☐ Without school leaving certificate
☐ Not specified

20. Do you live ...?

- ☐ Alone
- ☐ Family/ partner
- ☐ In a nursing home
- ☐ Other

21. Your health insurance status:

- ☐ Statutory health insurance
- ☐ Private health insurance

22. I'm classified as

- ☐ Not handicapped → Proceed to question 23.
- ☐ Handicapped
- ☐ Severely handicapped

Degree of disability:

_____ %

23. Your level of care:

- ☐ None
- ☐ _____ degree (number)
- ☐ I don't know, but applicable

24. Do you currently use one of the following devices due to your MS?

(Multiple answers possible)

- ☐ Sticks/Rollator
- ☐ Wheelchair
- ☐ Hospital bed
- ☐ Other

Acknowledgement

An dieser Stelle möchte ich meinen Dank den nachstehenden Personen aussprechen, die die vorliegende Promotion unterstützt haben.

Mein Dank gilt zunächst meinem Doktorvater Prof. Dr. Tjalf Ziemssen für die hervorragende Betreuung dieser Arbeit. Die zahlreichen Treffen in Dresden, Nürnberg und auf Kongressen waren stets konstruktiv und auch persönlich bereichernd. Besonders hervorheben möchte ich die tatkräftige Unterstützung und Ermutigung, meinen ersten wissenschaftlichen Vortrag auf einem internationalen Kongress zu halten. Ferner danke ich den wissenschaftlichen Mitarbeitern am MS-Zentrum für deren Unterstützung. Dabei sind besonders Dirk Schriefer und Rocco Haase zu nennen, die tatkräftig bei statistischen Fragestellungen zur Seite standen. Die Zeit als Doktorand am MS-Zentrum bleibt mir in herausragender Erinnerung.

Ferner danke ich den aktuellen und ehemaligen Kollegen bei Novartis. Besonders hervorzuheben sind Dr. Christian Cornelissen, der dieses Promotionskonstrukt in die Wege geleitet hat sowie Dr. Benjamin Etle, Hannes Kauper, Kathrin Jeckel und Dr. Simone Hiltl, die die außergewöhnliche Finanzierung der Promotion ermöglicht haben. Dr. Benjamin Etle danke ich weiterhin für seine akademische Unterstützung bei der Anfertigung von Publikationen als auch für sein persönliches Mentoring. Mein weiterer Dank gilt meiner Büronachbarin Dr. Carolin Baufeld für die vielfachen spontanen medizinischen Konsultationen sowie Marvin Heyne für statistische Diskussionen. Weiterhin danke ich dem gesamten Neuroscience-Team, die meine Erkenntnisse in mehreren Vorträgen kritisch gewürdigt haben.

Mein größter Dank gebührt meiner Partnerin und meiner Familie. Mit unermüdlicher Geduld hat mich Johanna während der Doktorarbeit unterstützt und mir in Zeiten hoher Belastung vollstes Verständnis entgegengebracht. Meinen Eltern, Astrid und Roland, danke ich für deren unablässige emotionale sowie finanzielle Unterstützung während meiner Studienzeit. Ihr habt mir meinen Lebensweg ermöglicht und euch widme ich die vorliegende Arbeit.

Anlage 1

Technische Universität

Medizinische Fakultät Carl Gustav Carus

Promotionsordnung vom 24.10.2014

Erklärung zur Eröffnung des Promotionsverfahrens

1. Hiermit versichere ich, dass ich die vorliegende Arbeit ohne unzulässige Hilfe Dritter und ohne Benutzung anderer als der angegebenen Hilfsmittel angefertigt habe; die aus fremden Quellen direkt oder indirekt übernommenen Gedanken sind als solche kenntlich gemacht.

2. Bei der Auswahl und Auswertung des Materials sowie bei der Herstellung des Manuskripts habe ich Unterstützungsleistungen von folgenden Personen erhalten:

Prof. Dr. Tjalf Ziemssen, Dirk Schriefer, Rocco Haase

3. Weitere Personen waren an der geistigen Herstellung der vorliegenden Arbeit nicht beteiligt. Insbesondere habe ich nicht die Hilfe eines kommerziellen Promotionsberaters in Anspruch genommen. Dritte haben von mir weder unmittelbar noch mittelbar geldwerte Leistungen für Arbeiten erhalten, die im Zusammenhang mit dem Inhalt der vorgelegten Dissertation stehen.

4. Die Arbeit wurde bisher weder im Inland noch im Ausland in gleicher oder ähnlicher Form einer anderen Prüfungsbehörde vorgelegt.

5. Die Inhalte dieser Dissertation wurden in folgender Form veröffentlicht:

- Ness N-H, Haase R, Kern R, Schriefer D, Etle B, Cornelissen C, Akgün K, Ziemssen T. 2020. The Multiple Sclerosis Health Resource Utilization Survey (MS-HRS): Development and Validation Study JMIR 22(3):e17921.
- Ness N-H, Schriefer D, Haase R, Etle B, Cornelissen C, Ziemssen T. 2019. Differentiating societal costs of disability worsening in multiple sclerosis. J Neurol 267(4):1035-42.
- Ness N-H, Schriefer D, Haase R, Etle B, Ziemssen T. 2020. Real-World Evidence on the Societal Economic Relapse Costs in Patients with Multiple Sclerosis. Pharmacoeconomics (online first).
- Ness N-H, Schriefer D, Haase R, Etle B, Wunderlich M, Akgün K, Ziemssen T. 2019. Real-World Evidence on the influence of relapses and relapse associated progression on societal economic costs in patients with multiple sclerosis ECTRIMS 2019 – Oral Presentations. Mult Scler 25(Suppl_2):3-130.

- Ness N-H, Schriefer D, Haase R, Ettl B, Wunderlich M, Akgün K, Ziemssen T. 2019. Societal economic impact of disability progression independent of relapse activity: a real world longitudinal analysis in patients with multiple sclerosis ECTRIMS 2019 – Poster Presentation. Mult Scler 25(Suppl_2):357-580.

6. Ich bestätige, dass es keine zurückliegenden erfolglosen Promotionsverfahren gab.

7. Ich bestätige, dass ich die Promotionsordnung der Medizinischen Fakultät der Technischen Universität Dresden anerkenne.

8. Ich habe die Zitierrichtlinien für Dissertationen an der Medizinischen Fakultät der Technischen Universität Dresden zur Kenntnis genommen und befolgt.

Dresden, den 12.05.2020

Nils-Henning Neß

Anlage 2

Hiermit bestätige ich die Einhaltung der folgenden aktuellen gesetzlichen Vorgaben im Rahmen meiner Dissertation:

- ☐ Das zustimmende Votum der Ethikkommission bei Klinischen Studien, epidemiologischen Untersuchungen mit Personenbezug oder Sachverhalten, die das Medizinproduktegesetz betreffen

Aktenzeichen der zuständigen Ethikkommissionen:

Technische Universität Dresden: EK97032011

Technische Universität Dresden: EK242082010

- ☐ Die Einhaltung der Bestimmung des Tierschutzgesetzes
Aktenzeichen der Genehmigungsbehörde zum Vorhaben/zur Mitwirkung
entfällt

- ☐ Die Einhaltung des Gentechnikgesetzes
entfällt

- ☐ Die Einhaltung von Datenschutzbestimmungen der Medizinischen Fakultät und des Universitätsklinikums Carl Gustav Carus.

Dresden, den 12.05.2020

Nils-Henning Neß